

Contents

Volume 107 (4) 1989

ESSAY IN DEVELOPMENT

Smith, L. D.

The induction of oocyte maturation: transmembrane signaling events and regulation of the cell cycle

685

CONTRIBUTED PAPERS

Sharpe, C. R., Pluck, A. and Gurdon, J. B.

XIF3, a *Xenopus* peripherin gene, requires an inductive signal for enhanced expression in anterior neural tissue

701

Simcox, A. A., Roberts, I. J. H., Hersperger, E.,

Gribbin, M. C., Shearn, A. and Whittle, J. R. S.

Imaginal discs can be removed from cultured embryos mutant for the segment-polarity genes *engrailed*, *naked* and *patched* but not from *wingless*

715

Basler, K. and Hafen, E.

Dynamics of *Drosophila* eye development and temporal requirements of *sevenless* expression

723

Kobayashi, S. and Okada, M.

Restoration of pole-cell-forming ability to u.v.-irradiated *Drosophila* embryos by injection of mitochondrial lrRNA

733

Duxson, M. J., Usson, Y. and Harris, A. J.

The origin of secondary myotubes in mammalian skeletal muscles: ultrastructural studies

743

Harris, A. J., Fitzsimons, R. B. and McEwan, J. C.

Neural control of the sequence of expression of myosin heavy chain isoforms in foetal mammalian muscles

751

Harris, A. J., Duxson, M. J., Fitzsimons, R. B. and Rieger, F.

Myonuclear birthdates distinguish the origins of primary and secondary myotubes in embryonic mammalian skeletal muscles

771

Jones, E. A. and Woodland, H. R.

Spatial aspects of neural induction in *Xenopus laevis*

785

Van Straaten, H. W. M., Hekking, J. W. M., Beurgens, J. P. W. M., Terwindt-Rouwenhorst, E. and Drukker, J.

Effect of the notochord on proliferation and differentiation in the neural tube of the chick embryo

793

Richardson, M. K., Hornbruch, A. and Wolpert, L.

Pigment pattern expression in the plumage of the quail embryo and the quail-chick chimaera

805

Barton, P. J. R., Harris, A. J. and Buckingham, M. E.

Myosin light chain gene expression in developing and denervated fetal muscle in the mouse

819

Cameron-Curry, P., Dulac, C. and Le Douarin, N. M.
Expression of the SMP antigen by oligodendrocytes in the developing avian central nervous system

825

Mohun, T. J., Garrett, N. and Taylor, M. V.

Temporal and tissue-specific expression of the proto-oncogene *c-fos* during development in *Xenopus laevis*

835

Lawrence, P. A. and Johnston, P.

Analysis of function of the pair-rule genes *hairy*, *even-skipped* and *fushi tarazu* in mosaic *Drosophila* embryos

847

Tix, S., Bate, M. and Technau, G. M.

Pre-existing neuronal pathways in the leg imaginal discs of *Drosophila*

855

Eichele, G.

Retinoic acid induces a pattern of digits in anterior half wing buds that lack the zone of polarizing activity

863

Lorenz, L. J., Hall, J. C. and Rosbash, M.

Expression of a *Drosophila* mRNA is under circadian clock control during pupation

869

Boocock, C. A.

Unidirectional displacement of cells in fibrillar matrices

881

Wilcox, M., DiAntonio, A. and Leptin, M.

The function of PS integrins in *Drosophila* wing morphogenesis

891

Kloc, M., Miller, M., Carrasco, A. E., Eastman, E. and Etkin, L.

The maternal store of the *xlgv7* mRNA in full-grown oocytes is not required for normal development in *Xenopus*

899

Levi, G. and Teichberg, V. I.

Patterns of expression of a 15K β -D-galactosidase-specific lectin during early development of the avian embryo

909

Kam, E. and Pitts, J. D.

Tissue-specific regulation of junctional communication in the skin of mouse fetuses homozygous for the repeated epilation (*Er*) mutation

923

Bagnall, K. M., Higgins, S. J. and Sanders, E. J.

The contribution made by cells from a single somite to tissues within a body segment and assessment of their integration with similar cells from adjacent segments

931

Dooley, T. P., Miranda, M., Jones, N. C. and DePamphilis, M. L.

Transactivation of the adenovirus EIIa promoter in the absence of adenovirus E1a protein is restricted to mouse oocytes and preimplantation embryos

945

Index of Authors and Titles

957

Subject Index

961