

# Contents

Volume 111 (1) 1991

<b>Young, P. E., Pesacreta, T. C. and Kiehart, D. P.</b> Dynamic changes in the distribution of cytoplasmic myosin during <i>Drosophila</i> embryogenesis	1	<b>Cunha, G. R., Young, P., Higgins, S. J. and Cooke, P. S.</b> Neonatal seminal vesicle mesenchyme induces a new morphological and functional phenotype in the epithelia of adult ureter and ductus deferens	145
<b>Ranscht, B. and Bronner-Fraser, M.</b> T-cadherin expression alternates with migrating neural crest cells in the trunk of the avian embryo	15	<b>Levi, G., Gumbiner, B. and Thiery, J. P.</b> The distribution of E-cadherin during <i>Xenopus laevis</i> development	159
<b>Condic, M. L., Fristrom, D. and Fristrom, J. W.</b> Apical cell shape changes during <i>Drosophila</i> imaginal leg disc elongation: a novel morphogenetic mechanism	23	<b>Sun, F. Z. and Moor, R. M.</b> Nuclear-cytoplasmic interactions during ovine oocyte maturation	171
<b>Maden, M., Hunt, P., Eriksson, U., Kuroiwa, A., Krumlauf, R. and Summerbell, D.</b> Retinoic acid-binding protein, rhombomeres and the neural crest	35	<b>de Pomerai, D. I., Ip, W. K., McLaughlin, M. and Perry, K. C.</b> Expression in non-lens tissues of an enzyme activity related to the 'lens-specific' protein, $\delta$ crystallin	181
<b>Ruberte, E., Dollé, P., Chambon, P. and Morriss-Kay, G.</b> Retinoic acid receptors and cellular retinoid binding proteins. II. Their differential pattern of transcription during early morphogenesis in mouse embryos	45	<b>Linsenmayer, T. F., Chen, Q., Gibney, E., Gordon, M. K., Marchant, J. K., Mayne, R. and Schmid, T. M.</b> Collagen types IX and X in the developing chick tibiotarsus: analyses of mRNAs and proteins	191
<b>Murphy, P. and Hill, R. E.</b> Expression of the mouse <i>labial</i> -like homeobox-containing genes, <i>Hox 2.9</i> and <i>Hox 1.6</i> , during segmentation of the hindbrain	61	<b>Cooke, J. and Wong, A.</b> Growth-factor-related proteins that are inducers in early amphibian development may mediate similar steps in amniote (bird) embryogenesis	197
<b>Sobis, H., Verstuyf, A. and Vandeputte, M.</b> Endodermal origin of yolk-sac-derived teratomas	75	<b>Trautman, M. S., Kimelman, J. and Bernfield, M.</b> Developmental expression of syndecan, an integral membrane proteoglycan, correlates with cell differentiation	213
<b>Prokop, A. and Technau, G. M.</b> The origin of postembryonic neuroblasts in the ventral nerve cord of <i>Drosophila melanogaster</i>	79	<b>Bloch-Gallego, E., Huchet, M., El M'Hamdi, H., Xie, F.-K., Tanaka, H. and Henderson, C. E.</b> Survival <i>in vitro</i> of motoneurons identified or purified by novel antibody-based methods is selectively enhanced by muscle-derived factors	221
<b>Bang, A. G., Hartenstein, V. and Posakony, J. W.</b> <i>Hairless</i> is required for the development of adult sensory organ precursor cells in <i>Drosophila</i>	89	<b>Lyons, G. E., Buckingham, M. E., Tweedie, S. and Edwards, Y. H.</b> Carbonic anhydrase III, an early mesodermal marker, is expressed in embryonic mouse skeletal muscle and notochord	233
<b>Ayer-Le Lievre, C., Ståhlbom, P.-A. and Sara, V. R.</b> Expression of IGF-I and -II mRNA in the brain and craniofacial region of the rat fetus	105	<b>Smith, S. M. and Eichele, G.</b> Temporal and regional differences in the expression pattern of distinct retinoic acid receptor- $\beta$ transcripts in the chick embryo	245
<b>Schmid, P., Cox, D., Bilbe, G., Maier, R. and McMaster, G. K.</b> Differential expression of TGF $\beta$ 1, $\beta$ 2 and $\beta$ 3 genes during mouse embryogenesis	117		
<b>Millan, F. A., Denhez, F., Kondaiah, P. and Akhurst, R. J.</b> Embryonic gene expression patterns of TGF $\beta$ 1, $\beta$ 2 and $\beta$ 3 suggest different developmental functions <i>in vivo</i>	131		