

# Contents

Volume 111 (2) 1991

- Fisher, C., Byers, M. R., Iadarola, M. J. and Powers, E. A.**  
Patterns of epithelial expression of Fos protein suggest important role in the transition from viable to cornified cell during keratinization 253
- Wiles, M. V. and Keller G.**  
Multiple hematopoietic lineages develop from embryonic stem (ES) cells in culture 259
- Mackenzie, A., Leeming, G. L., Jowett, A. K., Ferguson, M. W. J. and Sharpe, P. T.**  
The homeobox gene Hox 7.1 has specific regional and temporal expression patterns during early murine craniofacial embryogenesis, especially tooth development *in vivo* and *in vitro* 269
- Davis, C. A., Holmyard, D. P., Millen, K. J. and Joyner, A. L.**  
Examining pattern formation in mouse, chicken and frog embryos with an *En*-specific antiserum 287
- Fernandez, D. E., Turner, F. R. and Crouch, M. L.**  
*In situ* localization of storage protein mRNAs in developing meristems of *Brassica napus* embryos 299
- Ginsberg, D., DeSimone, D. and Geiger, B.**  
Expression of a novel cadherin (EP-cadherin) in unfertilized eggs and early *Xenopus* embryos 315
- Takahashi, Y. and Nogawa, H.**  
Branching morphogenesis of mouse salivary epithelium in basement membrane-like substratum separated from mesenchyme by the membrane filter 327
- Shi, X., Lu, L., Qiu, Z., He, W. and Frankel, J.**  
Microsurgically generated discontinuities provoke heritable changes in cellular handedness of a ciliate, *Stylonychia mytilus* 337
- Boulter, C. A., Aguzzi, A., Williams, R. L., Wagner, E. F., Evans, M. J. and Beddington, R.**  
Expression of *v-src* induces aberrant development and twinning in chimaeric mice 357
- Eldon, E. D. and Pirrotta, V.**  
Interactions of the *Drosophila* gap gene *giant* with maternal and zygotic pattern-forming genes 367
- Yasuda, G. K., Baker, J. and Schubiger, G.**  
Independent roles of centrosomes and DNA in organizing the *Drosophila* cytoskeleton 379
- Boulet, A. M., Lloyd, A. and Sakonju, S.**  
Molecular definition of the morphogenetic and regulatory functions and the *cis*-regulatory elements of the *Drosophila Abd-B* homeotic gene 393
- Irvine, K. D., Helfand, S. L. and Hogness, D. S.**  
The large upstream control region of the *Drosophila* homeotic gene *Ultrabithorax* 407
- Swalla, B. J., Badgett, M. R. and Jeffery, W. R.**  
Identification of a cytoskeletal protein localized in the myoplasm of ascidian eggs: localization is modified during anural development 425
- Sánchez-Herrero, E.**  
Control of the expression of the bithorax complex genes abdominal-A and Abdominal-B by *cis*-regulatory regions in *Drosophila* embryos 437
- Lyons, G. E., Buckingham, M. E. and Mannherz, H. G.**  
 $\alpha$ -actin proteins and gene transcripts are colocalized in embryonic mouse muscle 451
- Wanaka, A., Milbrandt, J. and Johnson, E. M.**  
Expression of FGF receptor gene in rat development 455
- Drysdale, T. A. and Elinson, R. P.**  
Development of the *Xenopus laevis* hatching gland and its relationship to surface ectoderm patterning 469
- Perkins, A. S., Mercer, J. A., Jenkins, N. A. and Copeland, N. G.**  
Patterns of *Evi-1* expression in embryonic and adult tissues suggest that *Evi-1* plays an important regulatory role in the mouse development 479
- Brown, R. and Brockes, J. P.**  
Identification and expression of a regeneration-specific homeobox gene in the newt limb blastema 489
- Ferretti, P., Brockes, J. P. and Brown, R.**  
A new type II keratin restricted to normal and regeneration limbs and tails is responsive to retinoic acid 497
- Minkoff, R., Parker, S. B. and Hertzberg, E. L.**  
Analysis of distribution patterns of gap junctions during development of embryonic chick facial primordia and brain 509
- Godsave, S. F. and Slack, J. M. W.**  
Single cell analysis of mesoderm formation in the *Xenopus* embryo 523

Contents

<b>Jones, C. M., Lyons, K. M. and Hogan, B. L. M.</b> Involvement of <i>Bone Morphogenetic Protein-4</i> (BMP-4) and <i>Vgr-1</i> in morphogenesis and neurogenesis in the mouse	531	<b>Perris, R., Krotoski, D., Lallier, T., Domingo, C., Sorrell, J. M. and Bronner-Fraser, M.</b> Spatial and temporal changes in the distribution of proteoglycans during avian neural crest development	583
<b>Miwa, Y., Atsumi, T., Imai, N. and Ikawa, Y.</b> Primitive erythropoiesis of mouse teratocarcinoma stem cells PCC3/A/1 in serum- free medium	543	<b>Kraut, R. and Levine, M.</b> Spatial regulation of the gap gene <i>giant</i> during <i>Drosophila</i> development	601
<b>Hopwood, N. D., Pluck, A. and Gurdon, J. B.</b> <i>Xenopus</i> Myf-5 marks early muscle cells and can activate muscle genes ectopically in early embryos	551	<b>Kraut, R. and Levine, M.</b> Mutually repressive interactions between the gap genes <i>giant</i> and <i>Krüppel</i> define middle body regions of the <i>Drosophila</i> embryo	611
<b>Mazarakis, N. D., Nelki, D., Lyon, M. F., Ruddy, S., Evans, E. P., Freemont, P. and Dudley, K.</b> Isolation and characterisation of a testis-expressed developmentally regulated gene from the distal inversion of the mouse <i>t</i> -complex	561	<b>Kelso-Winemiller, L. C. and Winkler, M. M.</b> 'Unmasking' of stored maternal mRNAs and the activation of protein synthesis at fertilization in sea urchins	623
<b>Sasaki, H., Hamada, T., Ueda, T., Seki, R., Higashinakagawa, T. and Sakaki, Y.</b> Inherited type of allelic methylation variations in a mouse chromosome region where an integrated transgene shows methylation imprinting	573	<b>Stocker, K. M., Sherman, L., Rees, S. and Ciment, G.</b> Basic FGF and TGF- $\beta$ 1 influence commitment to melanogenesis in neural-crest-derived cells of avian embryos	635