

Contents

Volume 111 (2) 1991

- Fisher, C., Byers, M. R., Iadarola, M. J. and Powers, E. A.**
Patterns of epithelial expression of Fos protein suggest important role in the transition from viable to cornified cell during keratinization 253
- Wiles, M. V. and Keller G.**
Multiple hematopoietic lineages develop from embryonic stem (ES) cells in culture 259
- Mackenzie, A., Leeming, G. L., Jowett, A. K., Ferguson, M. W. J. and Sharpe, P. T.**
The homeobox gene Hox 7.1 has specific regional and temporal expression patterns during early murine craniofacial embryogenesis, especially tooth development *in vivo* and *in vitro* 269
- Davis, C. A., Holmyard, D. P., Millen, K. J. and Joyner, A. L.**
Examining pattern formation in mouse, chicken and frog embryos with an *En*-specific antiserum 287
- Fernandez, D. E., Turner, F. R. and Crouch, M. L.**
In situ localization of storage protein mRNAs in developing meristems of *Brassica napus* embryos 299
- Ginsberg, D., DeSimone, D. and Geiger, B.**
Expression of a novel cadherin (EP-cadherin) in unfertilized eggs and early *Xenopus* embryos 315
- Takahashi, Y. and Nogawa, H.**
Branching morphogenesis of mouse salivary epithelium in basement membrane-like substratum separated from mesenchyme by the membrane filter 327
- Shi, X., Lu, L., Qiu, Z., He, W. and Frankel, J.**
Microsurgically generated discontinuities provoke heritable changes in cellular handedness of a ciliate, *Stylonychia mytilus* 337
- Boulter, C. A., Aguzzi, A., Williams, R. L., Wagner, E. F., Evans, M. J. and Beddington, R.**
Expression of *v-src* induces aberrant development and twinning in chimaeric mice 357
- Eldon, E. D. and Pirrotta, V.**
Interactions of the *Drosophila* gap gene *giant* with maternal and zygotic pattern-forming genes 367
- Yasuda, G. K., Baker, J. and Schubiger, G.**
Independent roles of centrosomes and DNA in organizing the *Drosophila* cytoskeleton 379
- Boulet, A. M., Lloyd, A. and Sakonju, S.**
Molecular definition of the morphogenetic and regulatory functions and the *cis*-regulatory elements of the *Drosophila Abd-B* homeotic gene 393
- Irvine, K. D., Helfand, S. L. and Hogness, D. S.**
The large upstream control region of the *Drosophila* homeotic gene *Ultrabithorax* 407
- Swalla, B. J., Badgett, M. R. and Jeffery, W. R.**
Identification of a cytoskeletal protein localized in the myoplasm of ascidian eggs: localization is modified during anural development 425
- Sánchez-Herrero, E.**
Control of the expression of the bithorax complex genes abdominal-A and Abdominal-B by *cis*-regulatory regions in *Drosophila* embryos 437
- Lyons, G. E., Buckingham, M. E. and Mannherz, H. G.**
 α -actin proteins and gene transcripts are colocalized in embryonic mouse muscle 451
- Wanaka, A., Milbrandt, J. and Johnson, E. M.**
Expression of FGF receptor gene in rat development 455
- Drysdale, T. A. and Elinson, R. P.**
Development of the *Xenopus laevis* hatching gland and its relationship to surface ectoderm patterning 469
- Perkins, A. S., Mercer, J. A., Jenkins, N. A. and Copeland, N. G.**
Patterns of *Evi-1* expression in embryonic and adult tissues suggest that *Evi-1* plays an important regulatory role in the mouse development 479
- Brown, R. and Brockes, J. P.**
Identification and expression of a regeneration-specific homeobox gene in the newt limb blastema 489
- Ferretti, P., Brockes, J. P. and Brown, R.**
A new type II keratin restricted to normal and regeneration limbs and tails is responsive to retinoic acid 497
- Minkoff, R., Parker, S. B. and Hertzberg, E. L.**
Analysis of distribution patterns of gap junctions during development of embryonic chick facial primordia and brain 509
- Godsave, S. F. and Slack, J. M. W.**
Single cell analysis of mesoderm formation in the *Xenopus* embryo 523

Contents

Jones, C. M., Lyons, K. M. and Hogan, B. L. M. Involvement of <i>Bone Morphogenetic Protein-4</i> (BMP-4) and <i>Vgr-1</i> in morphogenesis and neurogenesis in the mouse	531	Perris, R., Krotoski, D., Lallier, T., Domingo, C., Sorrell, J. M. and Bronner-Fraser, M. Spatial and temporal changes in the distribution of proteoglycans during avian neural crest development	583
Miwa, Y., Atsumi, T., Imai, N. and Ikawa, Y. Primitive erythropoiesis of mouse teratocarcinoma stem cells PCC3/A/1 in serum- free medium	543	Kraut, R. and Levine, M. Spatial regulation of the gap gene <i>giant</i> during <i>Drosophila</i> development	601
Hopwood, N. D., Pluck, A. and Gurdon, J. B. <i>Xenopus</i> Myf-5 marks early muscle cells and can activate muscle genes ectopically in early embryos	551	Kraut, R. and Levine, M. Mutually repressive interactions between the gap genes <i>giant</i> and <i>Krüppel</i> define middle body regions of the <i>Drosophila</i> embryo	611
Mazarakis, N. D., Nelki, D., Lyon, M. F., Ruddy, S., Evans, E. P., Freemont, P. and Dudley, K. Isolation and characterisation of a testis-expressed developmentally regulated gene from the distal inversion of the mouse <i>t</i> -complex	561	Kelso-Winemiller, L. C. and Winkler, M. M. 'Unmasking' of stored maternal mRNAs and the activation of protein synthesis at fertilization in sea urchins	623
Sasaki, H., Hamada, T., Ueda, T., Seki, R., Higashinakagawa, T. and Sakaki, Y. Inherited type of allelic methylation variations in a mouse chromosome region where an integrated transgene shows methylation imprinting	573	Stocker, K. M., Sherman, L., Rees, S. and Ciment, G. Basic FGF and TGF- β 1 influence commitment to melanogenesis in neural-crest-derived cells of avian embryos	635