

Ectopic expression of *seven-up* causes cell fate changes during ommatidial assembly

Yasushi Hiromi^{1,2}, Marek Mlodzik^{1,3}, Steven R. West², Gerald M. Rubin¹ and Corey S. Goodman¹

¹Howard Hughes Medical Institute and Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA

²Department of Molecular Biology, Princeton University, Princeton, NJ 08544-1014, USA

³Differentiation Programme, EMBL, Heidelberg, D-6900 Germany

SUMMARY

During *Drosophila* ommatidial development, a single cell is selected within the ommatidial cluster to become the R7 photoreceptor neuron. The *seven-up* gene has been shown to play a role in this process by preventing four other photoreceptor precursors, R3/R4/R1/R6, from adopting the R7 cell fate. The *seven-up* gene encodes a steroid receptor-like molecule that is expressed only in those four cells that require *seven-up* function in the developing *Drosophila* ommatidium. We have examined the functional significance of the spatially restricted expression of *seven-up* by misexpressing seven-up isoforms. As expected from the function that seven-up performs in R3/R4/R1/R6, ubiquitous expression of seven-up causes transformation of the R7 cell to an R1-R6 cell fate. In addition, depending on the timing and spatial pattern of expression, various other phenotypes are produced including the loss of the R7 cell and the

formation of extra R7 cells. Ubiquitous expression of *seven-up* close to the morphogenetic furrow interferes with R8 differentiation resulting in failure to express the boss protein, the ligand for the sevenless receptor tyrosine kinase, and the R7 cell is lost consequently. Extra R7 cells are formed by recruiting non-neuronal cone cells as photoreceptor neurons in a *sevenless* and *bride of sevenless* independent way. Thus, the spatiotemporal pattern of *seven-up* expression plays an essential role in controlling the number and cellular origin of the R7 neuron in the ommatidium. Our results also suggest that *seven-up* controls decisions not only between photoreceptor subtypes, but also between neuronal and non-neuronal fates.

Key words: *Drosophila*, *seven-up*, steroid receptor, ommatidial assembly, cell fate

INTRODUCTION

How the diversity of neurons is generated during neurogenesis is one of the central questions in developmental biology. The compound eye of *Drosophila* offers an excellent model system for studying the genetic control of specification of neuronal identities. The *Drosophila* eye is a hexagonal array of ~800 ommatidia, or unit eyes, each containing 8 photoreceptor neurons, R1 through R8, and 12 non-neuronal accessory cells. Individual photoreceptor neurons can be uniquely identified by their morphology and the stereotyped position that they occupy (reviewed by Tomlinson, 1988; Ready, 1989). Since no lineage restrictions exist within the 20 cells that constitute an ommatidium, cells acquire their identity by responding to signals from neighboring cells within the ommatidium (Ready et al., 1976; Lawrence and Green, 1979; Wolf and Ready, 1991). A number of genes have been identified that are required for correct specification of cell fates during ommatidial assembly. Through mosaic analysis one can identify the cells in which a gene activity is required, and deduce its role in the cell-cell interactions that mediate cell fate decisions. Molecular features of the genes isolated so far are consis-

tent with the proposed cell-cell interactions and induction mechanisms (reviewed by Banerjee and Zipursky, 1990; Hafen, 1991; Rubin, 1991)

The specification of the R7 neuron has been studied in the most detail and is at present the best understood (reviewed by Rubin, 1991). R7 is the UV-sensitive photoreceptor that synapses in a layer of the optic lobe distinct from all other photoreceptor neurons in each ommatidium (reviewed by Hardie, 1986). In normal development, the R7 neuron differentiates from a cell that occupies a fixed position in the ommatidial cluster, between R1 and R6. Several genes have been identified that are involved in selecting a single photoreceptor precursor as the R7 photoreceptor. Loss-of-function alleles of *sevenless* (*sev*), *bride of sevenless* (*boss*) and *seven in absentia* (*sina*) all transform the R7 precursor cell to a non-neuronal cone cell (Tomlinson and Ready, 1986, 1987b; Reinke and Zipursky, 1988; Carthew and Rubin, 1990). The *sevenless* gene encodes a receptor tyrosine kinase (Hafen et al., 1987; Bowtell et al., 1988; Basler and Hafen, 1988; Simon et al., 1989), whereas *boss* encodes a transmembrane protein that is expressed in the R8 cell and is the ligand for *sev* (Hart et al., 1990; Kramer et al., 1991). Genetic analyses indicate that the *sev* tyrosine

kinase acts through activation of the ras pathway (Simon et al., 1991; Fortini et al., 1992; Rogge et al., 1991; Bonfini et al., 1992; Gaul et al., 1992). Increased levels of ras activity in the cone cells achieved by either a ligand-independent allele of *sev* (Basler et al., 1991), ectopic expression of *boss* (Van Vactor et al., 1991), expression of activated *Ras1* (Fortini et al., 1992), or by reduction of the activity of *Gap1* (Gaul et al., 1992; Rogge et al., 1992; Buckles et al., 1992) all result in transformation of cone cells to R7 neurons. In contrast, loss-of-function alleles of *rough* (Tomlinson et al., 1988; Heberlein et al., 1991; Van Vactor et al., 1991) and *seven-up* (*svp*) (Mlodzik et al., 1990b) cause more than one photoreceptor precursor to adopt the R7 fate without affecting cone cell differentiation. In particular, loss of *svp*⁺ function results in the cell autonomous transformation of four outer photoreceptor cells, R3/R4/R1/R6, towards R7-like cells (Mlodzik et al., 1990b).

The predicted *svp* protein shares homology with members of the steroid receptor family (reviewed by Evans, 1988; Green and Chambon, 1988), suggesting that it acts as a ligand-responsive transcription factor (Mlodzik et al., 1990b). Two human homologues of *svp* have been identified that share extensive homology in both the DNA-binding domain and the ligand-binding domain (Miyajima et al., 1988; Wang et al., 1989; Ladias and Karathanasis, 1991). A striking aspect of *svp* expression is that, despite its apparent structure as a receptor, there is complete coincidence between the cells that express *svp* and those that require its function (Mlodzik et al., 1990b). This is in contrast to the *sev* receptor tyrosine kinase, which is required only in the R7 cell but is expressed in most photoreceptor precursors as well as cone cells (Tomlinson et al., 1987; Banerjee et al., 1987). The expression pattern of *sev* does not play a major role in restricting R7-forming potential, since ectopic expression of *sev* in all cells under heat-shock promoter does not cause other cells to adopt the R7 cell fate (Basler and Hafen, 1989a; Bowtell et al., 1989a). The restriction of *sev* activity is achieved by local presentation of its ligand, the *boss* protein by the R8 cell. There is also a restriction in the ability of *sev*-expressing cells to internalize the *boss* protein to the R7 precursor, but the significance of this restriction in controlling R7-forming potential is not clear (Kramer et al., 1990; Van Vactor et al., 1991; Cagan et al., 1992). The ligand for *svp* is not identified, nor its distribution known.

Here we have tested the functional significance of the *svp* expression pattern by analyzing the consequences of ectopic expression of *svp*. We observe a variety of cell fate transformations within an ommatidium including both the loss and gain of R7 cells. Our results indicate that the spatially restricted expression of *svp* plays an essential role in controlling the number of R7 cells that form within an ommatidium.

MATERIALS AND METHODS

Plasmid construction and P-element-mediated transformation

P-element constructs containing *hs-svp1* and *hs-svp2* genes were made by cloning a 1.7 kb *EagI* fragment of pc162.1 and a 2.4 kb *EagI-ClaI* fragment of pc162.2 (Mlodzik et al., 1990b), respectively, into the polylinker region of the CaSpeRhs vector

(Thummel and Pirrotta, 1991). P-elements containing *sev-svp1* and *sev-svp2* genes were made by first inserting a 0.57 kb *BamHI* fragment of CaSpeR-hs containing the trailer sequence of the *hsp70* gene into the *BamHI* site of SE8/DM30 (Bowtell et al., 1989b) and then inserting the 2.9 kb *EcoRI-ClaI* fragment of pc162.1 and the 2.7 kb *EcoRI-ClaI* fragment of pc162.2, respectively, into the *ClaI* site located upstream of the *BamHI* site in the SE8/DM30 vector. *sev-svp Mlu* has an insertion of a stop codon linker (New England Biolabs) at the *MluI* site of *sev-svp2*, truncating the protein at residue 273. *sev-svp Sal* has a deletion of a 1 kb *SalI-ClaI* fragment truncating the protein 17 residues before the divergence point of the two isoforms.

Germ-line transformation was done using *ry*⁵⁰⁶ and *w*¹¹¹⁸ as host strains and p 25.7wc (Karess and Rubin, 1984) as a helper plasmid. Secondary jumps to new locations were made using a strain carrying a genomic source of transposase activity (Robertson et al., 1988).

Histology

Antibody stainings of imaginal discs were performed as described (Tomlinson and Ready, 1987a) except that in most cases the peripodial membrane was not removed. Affinity-purified rabbit antibody against BarH1/BarH2 proteins (Higashijima et al., 1992) was a kind gift of K. Saigo. Monoclonal antibody anti-*boss1* (Kramer et al., 1991) was a generous gift of L. Zipursky. Monoclonal antibody against β -galactosidase was purchased from Promega. Monoclonal antibody against elav protein was made in the Rubin laboratory monoclonal antibody facility. Sections of adult retinae were made according to Tomlinson and Ready (1987a).

Marker strains

The following enhancer trap marker lines were used as cell-type-specific markers; BB02 (Hart et al., 1990) and rO156 (U. Gaul unpublished), which express β -galactosidase late in R8 differentiation, H214, which expresses β -galactosidase strongly in the R7 cell (Mlodzik et al., 1992), and rI533, an insertion in the *Gap1* gene (Gaul et al., 1992).

Generation of *svp* mutant clones by FLP-FRT mediated mitotic recombination

Clones homozygous for a null allele of *svp*, *svp*^{e22} (Mlodzik et al., 1990b), were made by mitotic recombination catalyzed by the FLP-FRT system (Golic and Lindquist, 1991). A 24 hour egg collection was taken from the crosses between males of *w*¹¹¹⁸ *hsFLP1/Y*; *75AE svp*^{e22}/*TM6B Hu, Tb* and females of *w*¹¹¹⁸; *75A M(3) w*¹²⁴/*TM6B Hu Tb*. *hsFLP1* has an insertion of the P[ry⁺; *hsFLP*] element on the X-chromosome (Golic and Lindquist, 1989). *75A* is an insertion of P[>*w*^{hs}>] element (> denotes an FRT site) at the base of 3R (Golic and Lindquist, 1991; K. Golic personal communication). *75AE* has an excision of the *w*^{hs} gene catalyzed by FLP-mediated recombination from the *75A* insertion and has one copy of FRT left. 40 hours after the end of the egg collection period, the vials containing larvae were submerged in a 37°C waterbath for 2 hours. Female *Tubby*⁺ larvae were selected as wandering third instar and their discs were processed for antibody staining. *Tubby*⁺ male siblings were used as controls that do not produce *svp* mutant clones. The *svp* mutant clones were visualized using an anti-BarH1/BarH2 antiserum as a molecular probe.

RESULTS

Ubiquitous expression of *svp* causes a variety of cell fate changes

Neuronal differentiation of the eye starts in the eye imaginal disc of the third instar larva as an indentation of the disc,




Fig. 1. Schematic structure and expression patterns of *svp* ectopic expression constructs. (A) cDNAs encoding the two isoforms of *svp*, as well as their truncated versions, were fused to the heat-inducible *hsp70* promoter, and to the *sev* enhancer/promoter. DNA, region showing homology to the DNA-binding domain of steroid receptors; ligand, the putative ligand-binding domain. (B) Schematic representation of ommatidia with eight photoreceptor precursors (numbered 1 through 8) and four cone cells (AC, anterior cone cell; EQC, equatorial cone cell; PC, posterior cone cell; PLC, polar cone cell). Expression patterns of the endogenous *svp* gene (*svp*) and the two promoter fusions (*hs-svp* and *sev-svp*) are indicated by shading. Light shading of R1/R6 in *sev-svp* shows lower levels of expression compared to R3/R4/R7 and cone cells.

called the morphogenetic furrow, moves in a posterior-to-anterior direction. Pattern formation starts in the furrow as cells are recruited to form ommatidial clusters containing photoreceptor precursors (Tomlinson and Ready, 1987a; Wolf and Ready, 1991). From the morphogenetic furrow, rosettes of cells consisting of four to five core cells surrounded by a ring of 10-15 cells emerge with regular spacing. Each group of cells is then transformed into a precluster containing five postmitotic photoreceptor precursors, R8, R2, R5, R3 and R4. After a wave of mitosis, three more photoreceptor precursors R1/R6/R7 and four cone cells join the cluster successively. Each of the eight photoreceptor precursors and the cone cells occupies a stereotyped position within the cluster. Based on expression of neuron specific antigens, it has been inferred that photoreceptor precursors initiate neuronal differentiation in an invariant sequence; R8 initiates differentiation first, followed by R2/R5, R3/R4, R1/R6 and finally R7 (Tomlinson and Ready, 1987).

Since an eye disc contains ommatidial clusters at different stages of differentiation, it should be possible to identify the developmental stage that is sensitive to the ectopic expression of *svp* by applying a pulse of *svp* expression throughout the eye disc. Two classes of cDNAs, called type 1 and type 2, have been identified from the *svp* locus. Type 1 cDNA encodes a protein that shares homology with both the DNA-binding domain and the ligand-binding domain of steroid receptors, whereas type 2 cDNA diverges from type 1 in the middle of the putative ligand-binding domain (Mlodzik et al., 1990b). To express these two *svp* isoforms in all cells in the imaginal disc, we generated transformant lines that carry P elements containing *svp* cDNAs that were placed under the control of the heat-inducible *hsp70* promoter (Fig. 1). Fusion genes employing type 1 and type 2 cDNAs will be called *hs-svp1* and *hs-svp2*, respectively.

When late third instar larvae carrying these fusion genes

were exposed to a brief (1-2 hours) heat pulse, 5 to 20% of the animals failed to pupate or died as early pupae. Those that survived to adulthood had no obvious defects in external morphology, except that their eyes had a stripe of rough region running dorsoventrally. In retinal sections of such eyes, a stripe of ommatidia with abnormal numbers of photoreceptor cells was found, the stripe often being wider than the region of the rough exterior. The width of stripe with abnormal ommatidia was often more than 10 rows. Since a new row of ommatidia is produced approximately every 2 hours (Basler and Hafen, 1989b), if ubiquitous expression of *svp* affected a single developmental stage, we would have expected to see a relatively narrow stripe of abnormal ommatidia, e. g. one or two rows. The broadness of the affected region suggests either that the *svp* protein expressed under heat-shock control has a long perdurance, or that ectopically expressed *svp* interferes with multiple differentiation steps.

In retinal sections of wild-type eyes, three classes of photoreceptor cells can be distinguished by their morphology. The outer photoreceptor cells R1 through R6 have rhabdomeres of large diameter that project throughout the depth of the retina, whereas the two classes of central photoreceptor cells, R7 and R8, have rhabdomeres of small diameter, the former located in the apical retina, the latter in the basal retina. Within the stripe of abnormal ommatidia in heat-shocked *hs-svp* flies, ommatidia with different phenotypes were observed: these include loss of outer photoreceptor cells, loss of central photoreceptor cells, appearance of extra central photoreceptor cells and appearance of extra outer photoreceptor cells (Figs 2, 3). Ommatidia with different phenotypes appeared ordered in an anterior-to-posterior manner, forming narrow substripes within the broader stripe of abnormally constructed ommatidia (Fig. 3). For example, in a *hs-svp1* eye, loss of one or two outer photoreceptor cells was seen in the anteriormost region of the




Fig. 2. Retinal phenotypes of heat-shocked *hs-svp* animals. (A) Wild type, (B) *hs-svp1*, (C) *hs-svp2*. Animals shown in B and C were heat shocked at 37°C for 2 hours as wandering third instar larvae. (A-C) Apical sections at the R7 level; (E-L) individual ommatidium corresponding to the labeled ommatidia in B and C, in basal sections at the R8 level. E lacks two outer photoreceptor cells without affecting R7 or R8; F and K lack both R7 and R8 yet contain six outer photoreceptor cells; G has an extra R7 and has a concomitant loss of an outer photoreceptor cell (class 2 in Fig. 3); H has an extra R7 with normal number of outer photoreceptor (class 1 in Fig. 3); I and L have a transformation of R7 to outer photoreceptor fate and J has reduced number of outer photoreceptor cell and contain extra central photoreceptor cells (class 2, in Fig. 3). A section of a wild-type ommatidia at the R8 level is shown in D.

stripe, partially overlapping a region of ommatidia that lacked the central photoreceptor cells R7 and R8. In the more posterior region of the affected stripe, ommatidia with extra central photoreceptor cells were present. This was followed by a row that contained mostly normal ommatidia, and further posterior was a region that lacked R7 but had an extra photoreceptor with the morphology of an outer photoreceptor cell (Figs 2B, 3D). Formation of substripes of ommatidia with specific phenotypes in a defined order suggests that each phenotype is caused by affecting a specific differentiation process that takes place in a stereotyped position in the developing eye imaginal disc. *hs-svp2* retinæ also contained substripes of ommatidia with mutant phenotypes similar to those seen in *hs-svp1* retinæ (Figs 2C,

3E-H). The order of specific substripes in *hs-svp2* retinæ, however, differed from that in *hs-svp1* retinæ (see for example the position of extra R7 phenotype, relative to that of the loss of R7 phenotype), indicating that superficially similar phenotypes are not necessarily caused by the same cellular mechanisms.

Ectopic expression of *svp* causes transformation of R7 to R1-R6 subtype

Previous analysis of loss-of-function phenotype of *svp* showed that *svp* prevents R3/R4/R1/R6 cells assuming the R7 cell fate (Mlodzik et al., 1990b). It is thus possible that ectopic expression of *svp* in the R7 precursor would prevent its differentiation and either transform it towards an




Fig. 3. A graphic representation of the *hs-svp* phenotypes. (A-C and D-F) Histograms representing specific phenotypes seen in the *hs-svp1* retina shown in Fig. 2B and the *hs-svp2* retina shown in Fig. 2C, respectively. Vertical axes show frequencies in percentage of ommatidia showing particular phenotypes among ommatidia in a given row. The position of row 0 is arbitrary. (A,D) Ommatidia with normal morphology. We define the stripe of affected ommatidia as rows 3 to 12 and rows 8 to 18 in retinas represented in A-C and D-F, respectively. Within this region, substripes of different phenotypes are observed. (B,E) Ommatidia lacking central photoreceptor cells (R7⁻, R8⁻, R7[→]outer). R7⁻ (loss of R7) class and R8⁻ (loss of R8) class are scored independently. These classes and R7[→]outer (transformation of R7 to outer photoreceptor cells) are mutually exclusive. Ommatidia exhibiting other phenotypes are depicted in C and F. Ommatidia that have extra R7-like cells are divided into two classes, class 1 and class 2. Class 1 ommatidia contain normal number

(6) of outer photoreceptor cells, whereas class 2 ommatidia have reduced numbers of outer photoreceptor cells. Class 1 ommatidia were not found in *hs-svp2* retinas. Ommatidia that lack outer photoreceptor cells with one or no R7 are classified as outer⁻. Note that both outer⁻ class and class 2 ommatidia have reduced numbers of outer photoreceptor cells. Ommatidia that show loss of central and outer photoreceptor cells are scored independently, as outer⁻ and R7⁻, thus the total percentage does not necessarily add up to 100%. Anterior to the major stripe of affected ommatidia, *hs-svp2* retinas usually have a small number of ommatidia that lack outer photoreceptor cells with or without R7. This region (corresponding to rows 1 to 5 in D-F), although reproducibly seen, was not included in the row counts of the affected stripe. For each construct, we generated histograms from four to five animals that received identical heat-shock treatment. Although the peak height of specific phenotypes varied from eye to eye, the relative order of specific phenotypes was invariant among retinas carrying a given construct.

R3/R4/R1/R6 cell fate or cause the loss of the R7 cell. Indeed, we found substripes of ommatidia that show such phenotypes in heat shocked *hs-svp* retinas (Figs 2, 3).

The posteriormost region of the affected stripe in *hs-svp1* and *hs-svp2* retinas contained ommatidia that lacked a photoreceptor with normal R7 cell morphology in the apical sections at the level that R7 cell is present in wild-type ommatidia. Even in such abnormal ommatidia, the cell corresponding to R7 can be identified by comparison with flanking normally constructed ommatidia. The rhabdomere of the affected R7 cells (R7T) are larger than those of the normal R7 cells, resembling those of the outer photoreceptor cells. Moreover the R7T rhabdomere was not in its normal central position, but is located between those of R1 and R6. Serial sections revealed that the R7T rhabdomere indeed extended throughout the depth of the retina, as those of normal R1-R6 subtypes (Fig. 2B,C,I,L). These phenotypes are indistinguishable from those observed in flies that express the rough protein in the R7 cell, transforming R7 cell into R1-R6 subtype (Basler et al., 1990; Kimmel et al. 1990). We conclude that ectopic expression of *svp*, like that of rough, causes transformation of the R7 cell to outer pho-

totoreceptor cells. Since the width of the ommatidial rows with this specific phenotype was approximately two rows, the R7 cell must be sensitive to ectopic expression of *svp* for at least 4 hours.

Perturbation of R8 development causes loss of R7

In addition to the transformation of the R7 cell to an outer photoreceptor cell described above, heat-shocked *hs-svp1* and *hs-svp2* retinas contained another substripe that lacked the R7 cell. This substripe was found two to five rows posterior to the anterior margin of the affected stripe (Figs 2, 3). Examination of basal sections revealed that most ommatidia that lacked R7 also were missing R8 (Figs 2,3). Of a total of 13 ommatidia arranged in a row in a heat-shocked *hs-svp1* retina, we found 7 ommatidia that lacked a central photoreceptor in apical sections, all of which also lacked the R8 in basal sections (Fig. 2). Two possibilities exist as to how the R7 cell was lost. First, expression of *svp* in the R7 precursor might have prevented its differentiation as a R7 neuron in a cell autonomous manner. Alternatively, since R8 is known to induce R7 differentiation by expressing boss on its surface (Kramer et al., 1991; Van Vactor et




Fig. 4. Expression of R8-specific markers upon gain and loss of *svp* expression. (A-C) Anti-boss staining. (A) A wild-type disc. The morphogenetic furrow is indicated by an arrow. The boss staining can be first detected in row three and extends towards the posterior end of the disc. (B) *hs-svp1*; (C) *hs-svp2*. Discs from animals that had been heat shocked for 45 minutes at 37°C and chased at 22°C for 19 hours. The open arrow marks the position of the boss-negative stripe. (D-F) Histochemical detection of β -galactosidase activity of BB02 enhancer trap marker in 40 hour pupal eyes. Third instar larvae were heat shocked for 45 minutes and then reared at 22°C. (D) Wild type, (E) *hs-svp1* and (F) *hs-svp2*. BB02 expresses β -galactosidase in R8 (arrow) and R7 (arrowhead), with R8 having higher levels of expression. In heat-shocked *hs-svp* eyes, the regular ommatidial array is disrupted by a stripe of ommatidia that lack β -galactosidase expression. Most ommatidia within such a stripe (indicated by an open arrow) lack β -galactosidase expression in both R7 and R8. Further posterior (towards bottom of the figure) is a region that lacks staining of the R7 cell. An example of such ommatidia is indicated by an arrow. (G) A mosaic disc containing *svp^{e22}, M⁺* clone stained with an anti-BarH1 antibody. In the *svp⁺* region, BarH1 is expressed strongly in R1/R6, and weakly in all cells located basally in the disc (out of focus). In the *svp⁻* region, R1/R6 do not express BarH1, whereas expression in the basal cells within the same clone is unaffected. At the boundary of *svp⁺* and *svp⁻* regions, ommatidia with a single BarH1-expressing cell are present. This suggests that the effect of loss of *svp* on BarH1 expression is cell autonomous. (H) A disc containing *svp^{e22}, M⁺* clone stained with a mixture of anti-BarH1 and anti-boss antibodies. *svp⁻* ommatidia can be identified as a region that does not express BarH1 in R1/R6 (slightly out of focus). boss expression is confined to a single cell in both *svp⁺* and *svp⁻* ommatidia.

al., 1991), the loss of R7 could be a consequence of the loss of R8. We explored the latter possibility first by examining whether or not boss expression was affected by ectopic expression of *svp*. Following a 1 hour heat shock, third instar larvae were reared at 22°C for various periods and then stained with an anti-boss monoclonal antibody (Kramer et al., 1991). In wild-type larvae that were heat shocked, as well as in non-heat-shocked *hs-svp* animals, boss protein could first be seen in R8 three rows posterior

to the morphogenetic furrow and remained detectable towards the posterior end of the eye disc (Kramer et al., 1991). When eye discs from *hs-svp* larvae were fixed 2 hours after heat shock, no abnormalities in boss expression were detected. However, a chase of 9 hours or longer resulted in appearance of a stripe of ommatidia where boss expression was absent or greatly reduced (Figs 4, 5). Since boss has an essential role in the induction of R7, we expect failure of R7 differentiation to occur in such ommatidia.




Fig. 5. Position of the boss-negative stripe in heat-shocked *hs-svp* animals after various recovery periods. The number of boss⁺ ommatidia anterior to the boss-negative stripe was plotted against the chase period after the heat-shock treatment. Since anti-boss staining starts in row 3 in heat-shocked wild-type disc, the position of the morphogenetic furrow is marked at row -3.

The width of the boss-negative stripe was much narrower than the stripe of abnormally constructed ommatidia seen in retinal sections but correlated with the width of the substripe that lacks central photoreceptor cells. This means that there is a narrow developmental window in which ubiquitous expression of *svp* results in the loss of boss expression.

The induction of R7 through a boss-sev interaction takes place eight to ten rows behind the furrow (Basler et al., 1989; Mullins and Rubin, 1991; Hart et al., 1991). We wished to determine whether ectopic expression of *svp* causes an early defect in the initiation of boss expression and/or R8 differentiation, or if it causes a loss of boss expression at the time that the boss-sev interaction occurs. The distance between the morphogenetic furrow and the boss-negative stripe increased with longer chase periods; chases of 12 and 20 hours produced boss-negative stripes 7 and 11 rows posterior to the furrow respectively (Fig. 5). This meant that a new row of ommatidia was produced every 2 hours, indicating that there was no significant effect in advancement of the morphogenetic furrow after heat-shock treatment, at least between these two time points. Extrapolation of such data indicated that the time that *svp* expression affected boss expression is one to two rows behind the furrow, anterior to the position at which boss protein could first be detected (Fig. 5). We conclude that ectopic expression of *svp* results in a failure to initiate boss expression, and that proper expression of boss in such cells is never recovered following heat shock.

In order to determine whether ectopic expression of *svp* resulted in specific repression of boss expression or a general defect in R8 identity, we examined the behavior of enhancer trap marker lines that express β -galactosidase in the R8 cell. Two enhancer trap lines BB02 (Hart et al., 1990) and rO156 (Ulrike Gaul, personal communication) express β -galactosidase in R8 from row 10 on throughout imaginal disc development, independent of *boss* function. Third instar larvae carrying these marker genes, as well as *hs-svp1* and *hs-svp2* transgenes were heat shocked for 1 hour and stained for β -galactosidase activity in 40-hour-old pupae. With both

hs-svp1 and *hs-svp2*, there were one to two rows in which most ommatidia failed to express these R8 markers (Fig. 4D-F; and data not shown). With BB02, which expresses β -galactosidase also in R7 albeit at a lower level than R8, simultaneous loss of R8 and R7 was observed (Fig. 4E,F). Since multiple independent R8 markers showed loss of expression, it is likely that ubiquitous expression of *svp* isoforms resulted in a change in R8 identity.

To follow the fate of the affected R8 cell, we used the D120 line, which is an enhancer trap insertion in the *scabrous* gene (Mlodzik et al., 1990a). In this line, β -galactosidase expression starts in the R8 cell in the morphogenetic furrow, reflecting R8 specification (Mlodzik et al., 1990a; Baker et al., 1990). Although *scabrous* transcript and protein are expressed only in the morphogenetic furrow region, due to the stability of *lacZ* transcript and/or protein, β -galactosidase is detectable in R8 towards the posterior edge of the eye disc, thus serving as an excellent marker to follow R8 development. Larvae carrying the D120 insertion and either *hs-svp1* or *hs-svp2* genes were subjected to a 1 hour heat shock, and stained with anti-boss and anti- β -galactosidase antibodies after various chase periods. With 2 and 4 hours of chase, no abnormality in boss or β -galactosidase expression could be detected (data not shown). At 6 hours after heat shock, ommatidia that should have initiated boss expression failed to stain with boss antibody, yet all such ommatidia still contained R8 that expressed β -galactosidase (Fig. 6B,D). This means that the specification of the R8 precursor was normal and that the R8 cell was present despite its failure to initiate boss expression. With 11 hours of chase, the level of β -galactosidase in R8 in many of boss-negative ommatidia was either low or undetectable, although ommatidia further posterior still expressed β -galactosidase (Fig. 6D,E). Upon a 13 hour chase, all boss-negative ommatidia also failed to express β -galactosidase (data not shown). These results suggest that the R8 precursor undergoes cell death between 6 and 13 hours following the ectopic expression of *svp*.

Loss of *svp* function does not cause extra R8 formation

The finding that ectopic expression of *svp* interferes with R8 differentiation prompted us to examine whether *svp* suppresses R8 differentiation in normal development. We asked whether the loss of *svp* function results in expression of R8 traits in photoreceptor cells other than R8 itself. Due to embryonic lethality of *svp* mutants, clones that were genotypically mutant for *svp* were generated in the eye imaginal discs. We found that in *svp* clones the BarH1/BarH2 proteins, which are normally expressed in R1 and R6 (Higashijima et al., 1992), are not expressed, consistent with transformation of their fates (Fig. 4G). Within such *svp* mutant clones, expression of boss was still restricted to a single cell within each ommatidium (Fig. 4H). We conclude that although *svp* can suppress R8 differentiation, it is not utilized or not essential for restriction of R8 differentiation during normal ommatidial development.

Effects of ectopic expression of *svp* in R7 and in cone cells

Experiments using *hs-svp* animals show that ubiquitous




Fig. 6. The fate of the R8 cell in heat-shocked *hs-svp1* animals. Wild-type (A,D) or *hs-svp1* (B,C,E,F) third instar larvae were heat shocked for 1 hour and stained with a mixture of anti-boss and anti- β -galactosidase antibodies, 6 hours (A,B,D,E) or 11 hours (C,F) after the heat shock. The morphogenetic furrow is located at the top edge of the picture in all panels. (A-C) Focused apically to show boss expression, which is seen as small dots; (D-F) a more basal focal plane at the level of the R8 nuclei that express β -galactosidase. The faint round stainings seen in the apical focal plane are the out-of-focus images of the β -galactosidase-positive nuclei. The position of the boss-negative stripe is indicated by open arrows in B, C. Arrows indicate

examples of ommatidia that fail to express boss, yet contain β -galactosidase-positive R8 nuclei. Arrowheads in C, F show ommatidia that express neither boss nor β -galactosidase, suggesting that the R8 cell has undergone cell death.

expression of *svp* causes transformation of R7 cell to outer photoreceptor cell fate at a specific stage during ommatidial development. To test if a similar phenotype is produced by specifically expressing *svp* in the R7 cell, transgenic animals that carry *svp* cDNAs under the control of *sev* regulatory elements were generated (Fig. 1). Sequences from the *sev* promoter and enhancer used in these fusion genes direct high levels of expression in R3/R4/R7 and in the cone cells and lower levels in R1/R6 (Bowtell et al., 1989b; Basler et al., 1990; Kimmel et al., 1990), thus achieving ectopic expression of *svp* in R7 and the cone cells. Seven lines carrying the *sev-svp1* gene and nine lines carrying the *sev-svp2* gene were examined for abnormality in ommatidial assembly. Six of the nine *sev-svp2* lines showed mild roughening of the eye with one copy of the transgene. Surprisingly, sections of such retinæ revealed that the roughening is not caused by the loss or transformation of R7 cells. On the contrary, some ommatidia had an extra photoreceptor with R7-like morphology, i.e. rhabdomere of small diameter located apically (Fig. 7A). When the copy number of the construct was increased, we did observe a small number of ommatidia with extra outer photoreceptor cells and others that did not have an R7 cell with R8 present basally. The majority of abnormal ommatidia, however, had extra R7-like cells, the frequency of such ommatidia and the number of extra R7 cells increasing with the copy number of the *sev-svp* gene (Fig. 7A,B). To establish the identity of the extra photoreceptor cells, we used a reporter strain where the promoter of the *Rh4* rhodopsin gene, which is expressed exclusively in the R7 cells, has been fused to

the *lacZ* gene (Fortini and Rubin, 1990). In *sev-svp2* retinæ, extra photoreceptors expressed this R7 specific trait indicating that the extra cells have an R7 identity (Fig. 8A,B).

There are two possible mechanisms that could generate this phenotype. First, expression of *svp* in R7 may have affected its differentiation, but such an effect is compensated by R7 differentiation from other cells. Alternatively, R7 development may not have been affected in these transgenic animals and the phenotype is caused solely by the formation of extra R7 cells. The fate of the R7 precursor and the origin of the R7-like cells were examined using an enhancer trap line H214 that expresses β -galactosidase at high levels in the R7 cell in the eye imaginal disc (Mlodzik et al., 1992). In *sev-svp2* larval eye discs, not only the R7 cell but also the cone cells showed strong expression of β -galactosidase (Fig. 8D). We also observed expression of the neuron specific antigens *elav* (Robinow and White, 1991) and the 22C10 antigen (Zipursky et al., 1984) in the cone cells (Fig. 8G, and data not shown), establishing that cone cells were transformed to R7 cells in this genotype. In discs containing four copies of the *sev-svp2* transgene, some nuclei located basally in the disc also expressed *elav* (Fig. 8I). These may be uncommitted cells that have not yet been recruited to the ommatidial cluster. We conclude that *svp* type 2 protein expressed under the *sev* enhancer/promoter does not interfere with R7 differentiation, but rather transforms cone cells towards R7 cells. Despite this transformation, most *sev-svp2* ommatidia had four or more cone cells, as visualized with cobalt sulfide staining of pupal discs (data not




Fig. 7. Retinal phenotype of *sev-svp* animals. (A) *sev-svp2* one copy, (B) *sev-svp2* two copies, (C) *sev-svp1* three copies, (D) *sev^{d2}*; *sev-svp2* two copies, (E) *sina³/sina¹*; *sev-svp2* two copies, (F) *sev-svp Mlu* four copies. Sections are at the apical level where R7 is visible. Refer to Fig. 2A for a wild-type control. Animals shown in D and E carry *white* and *scarlet* mutations, respectively, and do not contain normal pigment granules.

shown). Thus, additional cells appear to be recruited to become cone cells when cone cell precursors enter the neuronal pathway. This is in agreement with other reports regarding the transformation of cone cell precursors (Basler et al., 1991; Gaul et al., 1992).

In contrast to *sev-svp2* lines, none of the *sev-svp1* lines showed any abnormality in ommatidial structure in single copy. When the copy number was increased to two or three, a phenotype similar to that seen in *sev-svp2* lines was observed (Fig. 7C). Thus the effect of *sev-svp1* appears to be weaker than that of *sev-svp2*.

***sina*, but not *sev* or *boss*, is required for *svp*-mediated cone cell transformation**

Three genes, *sev*, *boss* and *sina*, are required for specification of the R7 precursor and in respective retinæ all ommatidia lack the R7 cell. We examined whether these three genes are also required for the formation of R7 cells developing from the cone cells in *sev-svp2* lines. In the presence of the *sev-svp2* gene, many ommatidia contained one or two R7 cells even when *sev* or *boss* activity was removed (Fig. 7D, and data not shown). Such retinæ contained ommatidia that expressed the *Rh4/lacZ* fusion gene, while in control eyes (*sev* or *boss* mutant alone) lacking R7, this R7 specific marker was never expressed (data not shown). On the contrary, in *sina*; *sev-svp2* animals no R7 cells formed, indicating that *sina* function is required for R7 differentiation in transformed cone cells (Fig. 7E).

To analyze the fate of the R7 precursor and the cone cells in *sev*; *sev-svp2* ommatidia, we examined expression of the H214 *lacZ* marker in larval eye discs. The cone cells expressed high levels of β -galactosidase as does the R7 cell

in wild type, whereas the β -galactosidase level in the R7 precursor was low, if not undetectable, as in *sev* discs (Fig. 8E). This suggests that the R7 precursor failed to develop as the R7 cell in the absence of *sev*, whereas R7-like cells differentiating from the cone cells do not require *sev* activity. Indeed, in clusters that contained elav-positive cone cells, a majority of R7 precursors failed to express elav (Fig. 8H). To test whether the R7-like cells differentiating from cone cell precursors are functional R7 neurons, we tested the UV phototactic behavior of *sev*; *sev-svp2* animals. Animals in which the R7 precursor and the cone cells have switched their fates showed normal phototactic behavior, indicating that transformed cone cells have normal functional properties of the R7 neuron (Fig. 9).

Three genetic situations are known that cause transformation of cone cells into R7: ubiquitous expression of *boss* that results in activation of the *sev* pathway (Van Vactor et al., 1991) expression of a *boss*-independent form of *sev* (Basler et al., 1991) and reduction of *Gap1* activity (Gaul et al., 1992; Rogge et al., 1992; Buckles et al., 1992). Since in *sev-svp2* discs, cone cells differentiate into R7 in the absence of *boss* or *sev* function, it is unlikely that *svp* causes its effect by regulation of *boss* or *sev* expression and/or function. A possible target for *svp* is *Gap1*, which shows highly regulated expression that is confined to photoreceptor cells and cone cells posterior to the morphogenetic furrow (Gaul et al., 1992). Using an enhancer trap insert in *Gap1*, the r1533 line, we tested whether *sev-svp2* down-regulated *Gap1* expression in the cone cells. In *sev-svp2* discs, cone cells still expressed β -galactosidase (Fig. 8K), indicating that the effect of *sev-svp2* was not mediated by repression of *Gap1* transcription.




Fig. 8. Expression of cell specific markers in *sev-svp2* eyes. (A, B) Anti- β -galactosidase staining of retina carrying the *Rh4/lacZ* fusion gene, which is expressed in a subset of the R7 cells. (A) Wild type; (B) *sev-svp2* two copies; (C-E) anti- β -galactosidase staining of eye imaginal disc carrying the H214 enhancer trap marker; (C) wild type; (D) *sev-svp2* two copies and (E) *sev^{LY3}; sev-svp2* two copies. The arrowhead shows the R7 precursor, arrows point to β -galactosidase-positive cone cells. (F-I) Anti-elav staining. (F) Wild type; photoreceptors are numbered in an ommatidium; R2/R5/R8 are not in this focal plane. (G) *sev-svp2* two copies. The arrowhead and the arrows show the R7 precursor and anti-elav-positive cone cells, respectively. (H) *sev^{LY3}; sev-svp2* two copies. Open triangle show the position of the R7 precursor, which fail to express elav. (I) *sev-svp2* four copies, stained with anti-elav. This is a composite photograph spliced approximately at the dorsoventral midline of the eye disc, with the left half focused on the photoreceptor precursor nuclei, and the right half focused on the *elav*⁺ cells located basally. (J, K) Anti- β -galactosidase staining of eye discs carrying an enhancer trap insertion in the *Gap1* gene (r1533 line). (J) Wild type and (K) *sev-svp2* two copies. Some of the cone cells expressing β -galactosidase are indicated with arrows.

The putative ligand-binding domain is required for expression of the dominant phenotypes

The observation that both isoforms of *svp* cause similar phenotypes when ectopically expressed raised the possibility that ectopically expressed *svp* may not require the putative ligand-binding domain for its function. Since many of the steroid receptors function as ligand-independent transcriptional activators when their ligand-binding domain is deleted (reviewed in Evans, 1988), we tested whether truncated *svp* proteins lacking the ligand-binding domain could produce the same phenotype as that caused by *svp* type 1 and type 2 isoforms. Three constructs were made: *sev-svp Mlu*, in which *sev* regulatory elements direct

expression of a *svp* protein truncated shortly after the DNA-binding domain, *sev-svp Sal*, where *svp* protein is truncated immediately before the point where type 1 and type 2 isoforms diverge, and *hs-svp Mlu* where the same protein as *sev-svp Mlu* would be produced under the heat inducible promoter (Fig. 1). Animals that carry up to four copies of either the *sev-svp Mlu* or *sev-svp Sal* genes were generated, but their ommatidia showed normal morphology in retinal sections. We also failed to detect any phenotype after heat shocking animals carrying the *hs-svp Mlu* gene (Fig. 7F and data not shown). Thus C-terminal portions of both type 1 and type 2 isoforms are necessary for production of dominant phenotypes in these misexpression experiments. This requirement for the C-terminal portion could




Fig. 9. Color choice preference for wild-type (Canton-S), *sev*^{d2} and *sev*^{d2}; *sev-svp2* flies. Flies were tested for the light color choice preference at 350 nm UV light and 550 nm green light. The phototactic value is calculated as follows:

$$= \frac{N(\text{green}) - N(\text{UV})}{N(\text{green}) + N(\text{UV})}$$

where N(green) and N(UV) are the numbers of flies attracted by green light and the UV light, respectively. Flies that lack functional R7 cells, unlike wild type, are attracted by the green light. Three independent lines carrying a single copy of the *sev-svp2* transgene were tested in *sev*^{d2} background. All show strong response to the UV light, indicating that the R7 cells that develop from cone cell precursors are functional and make proper connections in the medulla.

be in the activity of the *svp* protein, the production of the stable product, or both.

DISCUSSION

Within an ommatidium, three cell types are known to have the potential to develop as the R7 neuron: the R7 precursor, the cone cells and the precursors of R1 through R6. During wild-type development, however, the R7 precursor is the only cell that adopts the R7 fate. Cone cells are prevented from becoming R7 first because they are not in contact with R8 and are thus unable to activate the *sev* pathway through *sev*-boss interaction and, second, because expression of *Gap1* in cone cells reduces their intrinsic ras activity. R3/R4/R1/R6 are prevented from becoming R7 by the expression of the *svp* gene in these cells (Mlodzik et al., 1990b). Although *svp*⁺ function is normally not required in R2/R5, in *rough* mutant ommatidia *svp* is ectopically expressed in R2/R5 and is required to prevent them from becoming R7 (Heberlein et al., 1991). Our results of ectopic expression of the *svp* isoforms show that proper transcriptional regulation of *svp* expression is essential for the proper development of two other cell types that have potential to become the R7 neuron, the R7 precursor and the cone cells.

Since expression of *svp* is restricted to R3/R4/R1/R6, *svp* could act as a genetic switch between the R3/R4/R1/R6 neuronal type and the R7 type, in a manner similar to the homeotic selector genes. In such a case, one would expect ectopic expression of *svp* in the R7 cell to interfere with its differentiation, and possibly transform it towards the R3/R4/R1/R6 fate. Indeed, in both *hs-svp1* and *hs-svp2* animals, we found ommatidia in which the R7 cell was transformed to an outer photoreceptor fate. Curiously, we failed to observe similar transformation in animals carrying *sev-svp* transgenes, which express *svp* isoforms in the R7 cell under the control of the *sev* enhancer/promoter. There

are a few possibilities that could explain this discrepancy. First, the R7 to outer photoreceptor transformation in *hs-svp* retinæ may not be caused by ectopic expression in the R7 cell, but may be due to expression in other cells that influence the differentiation of the R7 precursor. Alternatively, under the *sev* enhancer/promoter control, the *svp* protein may not have been expressed in the R7 cell in the appropriate temporal pattern to effect this transformation. We favor the latter possibility because the stage that is susceptible to R7-to-outer photoreceptor transformation maps late in ommatidial assembly. In *hs-svp1* retinæ, the substripe containing transformed R7 was located posterior to the substripe containing extra R7 cells, which is likely to correspond to the cone cell-to-R7 transformation seen in *sev-svp* retinæ. Since expression in R7 directed by *sev* enhancer/promoter sequences ceases before expression in the cone cells does (Bowtell et al., 1989b), expression of *svp* in R7 may not have persisted long enough to cause transformation of the R7 cell. In addition, in the case of the *rough* gene that specifies the R2/R5 cell fate, its ectopic expression in R7 alone is sufficient to transform R7 to the outer photoreceptor fate (Basler et al., 1990; Kimmel et al., 1990) and no data thus far available suggest a requirement for an inductive signal controlling the photoreceptor subtype decision of the R7 cell.

Another phenotype caused by ectopic expression of *svp* that interferes with R7 differentiation is the loss of R7 seen in *hs-svp* retinæ. It is unlikely that this phenotype is caused by ectopic expression in the R7 cell itself, for the following reasons. First, the loss of the R7 cell is usually accompanied by the loss of R8. Concomitant loss of R7 and R8 suggests that these two events are not caused by independent effects on R7 and R8. Second, ubiquitous expression of *svp* affects expression of the boss protein which serves as a ligand for the *sev* receptor tyrosine kinase. Since activation of *sev* is essential for the R7 precursor to assume a neuronal fate, the effect on boss expression can alone account for failure of R7 differentiation. Third, the ectopic expression of *svp* in R7 directed by the *sev* enhancer/promoter does not affect R7 differentiation. Taken together, these data strongly suggest that the loss of R7 caused by the ubiquitous expression of *svp* is due to the loss of the R8 cell. Whether the loss of R8 is caused by expression of *svp* in the R8 cell, or due to a nonautonomous mechanism, is not known.

Since R8 is believed to play a central role in initiating a series of induction steps (Tomlinson and Ready, 1987a), it is rather surprising that, in an ommatidium lacking R8, assembly proceeds with minor effects on the induction of outer photoreceptor cells. The sensitive period for the effect on R8 is quite early, one or two rows posterior to the morphogenetic furrow (Fig. 5). We show, however, that *hs-svp* does not affect specification of the R8 cell, as visualized by the expression of an enhancer trap insertion in the *scabrous* gene, and that the loss of R8 is likely to be caused by cell death that occurs between 6 and 13 hours after heat shock. Anterior to the stripe of ommatidia that have lost R8, there is another substripe consisting of ommatidia with different phenotypes (Figs 2, 3). This implies that there is yet another step in ommatidial assembly that *svp* can interfere with, which takes place prior to the effect on R8 differentiation and boss expression. Therefore, it is likely that the R8 cell

had already performed its function to induce other cells (e.g. R2/5) prior to the step that is affected by the ectopic expression of *svp*. Such a process must take place at or near the morphogenetic furrow, possibly prior to the formation of the 5-cell precluster (Banerjee and Zipursky, 1990).

Both the loss-of-function phenotype and the R7-to-outer photoreceptor transformation phenotype caused by *hs-svp* transgenes is consistent with the idea that *svp* acts by preventing R7 differentiation. It was thus unexpected that the ectopic expression of *svp* would cause the transformation of the cone cells towards R7 neurons. R7 differentiation from the cone cells caused by *sev-svp* transgene expression does not require the function of *boss* or *sev*, whereas it is completely suppressed by mutations in *sina*. Thus, ectopically expressed *svp* acts downstream of the *sev* receptor tyrosine kinase, but acts either upstream or in parallel to the *sina* gene. These epistatic relationships are similar to other conditions that cause the same cellular transformation, i.e., the expression of activated *Ras1* (Fortini et al., 1991) and the reduction in the *Gap1* activity (Gaul et al., 1992; Rogge et al., 1992; Buckles et al., 1992). Similarities in phenotypes and genetic relationships of *svp* and the activated ras pathway suggest that *svp* might act through ras to provide the potential to become a neuron. In support of this view, we have identified alleles of the *Ras1* gene among dominant suppressors of rough eye phenotype caused by the *sev-svp2* transgene (S. Kramer, F. Birkmeyer, M. M. and Y. H. unpublished). Although phenotypes caused by the loss of *svp* function indicate that *svp* is involved in a decision between two neuronal cell types, there is some evidence suggesting that *svp* plays a role similar to that of *sev* in providing neuronal fate per se. In ommatidia that are doubly mutant for *svp* and *sev*, not only R7 but also some of the outer photoreceptors fail to adopt a photoreceptor cell fate (Mlodzik et al., 1990b). Since *sev* is not required in R3/R4/R1/R6 in *svp*⁺ ommatidia, it appears that the role that *svp* plays in R3/R4/R1/R6 is not simply to control their photoreceptor subtype, but also to decide between neuronal versus non-neuronal fate. Thus the role of *svp* appears different from that of the *rough* gene, which specifies a subtype of outer photoreceptors, but does not have the potential to induce neuronal development of cone cells (Kimmel et al., 1990; Basler et al., 1990). It should be noted, however, that many of the dominant phenotypes can be caused by both type 1 and type 2 isoforms, which differ in the putative ligand-binding domain. It is therefore possible that the phenotypes observed are generated in a ligand-independent way, due to a function that is at least in part different from the one that *svp* performs in R3/R4/R1/R6.

We have shown that spatially restricted expression of the *svp* gene is essential for execution of its normal function. This result is similar to those obtained with another *Drosophila* member of the steroid receptor gene family, the *tailless* gene (Steingrímsson et al., 1991) which, like *svp*, is expressed in the region of the embryo requiring its function. Ectopically expressed *tailless* appears to have the same function as the endogenous gene product, suggesting that either the *tailless* ligand is uniformly distributed, or the *tailless* function is ligand-independent. On the contrary, ubiquitous expression of the *ultraspiracle* gene, which is the *Drosophila* homolog of the retinoid X receptor and is likely

to function as a heterodimer with the ecdysone receptor (Yao et al., 1992), do not interfere with normal development (Oro et al., 1992). These differences may reflect differences in strategies that steroid receptors utilize to regulate their functions, such as the distribution of the receptor itself, distribution of the ligand, or the distribution of the receptor's heterodimeric partner.

We thank K. Golic, P. Lawrence, U. Gaul, M. Fortini and L. Zipursky for providing fly strains, K. Saigo and L. Zipursky for gifts of antibodies, D. Bowtell and C. Thummel for their expression vectors, D. Brunner and E. Hafen for help with phototactic behavior experiments, S. Kramer for help in *sev-svp2* experiments, and A. Ephrussi, A. Bejsovec and S. Roth for valuable comments on the manuscript. This work was supported by a National Institutes of Health Training Grant (S. R. W.) and grants from National Institutes of Health (R01 NS29662) and Pew Scholars Program (Y. H.), and Swiss National Science Foundation (M. M.). C. S. G. and G. M. R. are investigators of the Howard Hughes Medical Institute.

REFERENCES

- Baker, N. E., Mlodzik, M. and Rubin, G. M. (1990). Spacing differentiation in the developing *Drosophila* eye: a fibrinogen-related lateral inhibitor encoded by *scabrous*. *Science* **250**, 1370-1377.
- Banerjee, U., Renfranz, P. J., Hinton, D. R., Rabin, B. A. and Benzer, S. (1987). The *sevenless* protein is expressed apically in cell membranes of developing *Drosophila* retina: it is not restricted to cell R7. *Cell* **51**, 151-158.
- Banerjee, U. and Zipursky, S. L. (1990). The role of cell-cell interaction in the development of the *Drosophila* visual system. *Neuron* **4**, 177-187.
- Basler, K. and Hafen, E. (1988). Control of photoreceptor cell fate by the *sevenless* protein requires a functional tyrosine kinase domain. *Cell* **54**, 299-311.
- Basler, K. and Hafen, E. (1989a). Ubiquitous expression of *sevenless*: position-dependent specification of cell fate. *Science* **243**, 931-934.
- Basler, K. and Hafen, E. (1989b). Dynamics of *Drosophila* eye development and temporal requirements of *sevenless* expression. *Development* **107**, 723-731.
- Basler, K., Siegrist, P. and Hafen, E. (1989). The spatial and temporal expression pattern of *sevenless* is exclusively controlled by gene-internal elements. *EMBO J.* **8**, 2381-2386.
- Basler, K., Christian, B. and Hafen, E. (1991). Ligand-independent activation of the *sevenless* tyrosine kinase changes the fate of cells in the developing *Drosophila* eye. *Cell* **64**, 1069-1081.
- Basler, K., Yen, D., Tomlinson, A. and Hafen, E. (1990). Reprogramming cell fate in the developing *Drosophila* retina: transformation of R7 cells by ectopic expression of *rough*. *Gene Dev.* **4**, 728-739.
- Bonfanti, L., Karlovich, C. A., Dasgupta, C. and Banerjee, U. (1992). The *Son of sevenless* gene product: a putative activator of ras. *Science* **255**, 603-606.
- Bowtell, D. D. L., Simon, M. A. and Rubin, G. M. (1988). Nucleotide sequence and structure of the *sevenless* gene of *Drosophila melanogaster*. *Genes Dev.* **2**, 620-634.
- Bowtell, D. D. L., Simon, M. A. and Rubin, G. M. (1989a). Ommatidia in the developing *Drosophila* eye require and can respond to *sevenless* for only a restricted period. *Cell* **56**, 931-936.
- Bowtell, D. D. L., Kimmel, B. E., Simon, M. A. and Rubin, G. M. (1989b). Regulation of the complex pattern of *sevenless* expression in the developing *Drosophila* eye. *Proc. Natl. Acad. Sci. USA* **86**, 6245-6249.
- Buckles, G. R., Smith, Z. D. J. and Katz, F. N. (1992). *mip* causes hyperinnervation of a retinotopic map in *Drosophila* by excessive recruitment of R7 photoreceptor cells. *Neuron* **8**, 1015-1029.
- Cagan, R. L., Kramer, H., Hart, A. C. and Zipursky, S. L. (1992). The bride of *sevenless* and *sevenless* interaction: internalization of a transmembrane ligand. *Cell* **69**, 393-399.
- Carthew, R. W. and Rubin, G. M. (1990). *seven in absentia*, a gene required for specification of R7 cell fate in the *Drosophila* eye. *Cell* **63**, 561-577.

- Evans, R. M. (1988). The steroid and thyroid hormone receptor superfamily. *Science* **240**, 889-895.
- Fortini, M. E. and Rubin, G. M. (1990). Analysis of cis-acting requirements of the Rh3 and Rh4 genes reveals a bipartite organization to rhodopsin promoters in *Drosophila melanogaster*. *Genes Dev.* **4**, 444-463.
- Fortini, M. E., Simon, M. A. and Rubin, G. M. (1991). Signalling by the *sevenless* protein tyrosine kinase is mimicked by Ras1 activation. *Nature* **355**, 559-561.
- Gaul, U., Mardon, G. and Rubin, G. M. (1992). A putative Ras GTPase activating protein acts as a negative regulator of signalling by the *Sevenless* receptor tyrosine kinase. *Cell* **68**, 1007-1019.
- Golic, K. G. and Lindquist, S. (1989). The FLP recombination of yeast catalyzes site-specific recombination in the *Drosophila* germline. *Cell* **59**, 499-509.
- Golic, K. G. and Lindquist, S. (1991). Site-specific recombination between homologous chromosomes in *Drosophila*. *Science* **252**, 958-961.
- Green, S. and Chambon, P. (1988). Nuclear receptors enhance our understanding of transcription regulation. *Trends Genet.* **4**, 309-314.
- Hafen, E. (1991). Patterning by cell recruitment in the *Drosophila* eye. *Current Opinion in Genetics and Development* **1**, 268-274.
- Hafen, E., Basler, K., Edstroem, J.-E. and Rubin, G. M. (1987). *Sevenless*, a cell-specific homeotic gene of *Drosophila*, encodes a putative transmembrane receptor with a tyrosine kinase domain. *Science* **236**, 55-63.
- Hardie, R. C. (1986). The photoreceptor array of the dipteran retina. *Trends in Neurosci.* **4**, 419-423.
- Hart, A. C., Krämer, H., Van Vactor, Jr., D. L., Paidhungat, M. and Zipursky, S. L. (1990). Induction of cell fate in the *Drosophila* retina: the bride of *sevenless* protein is predicted to contain a large extracellular domain and seven transmembrane segments. *Genes Dev.* **4**, 1835-1847.
- Heberlein, U., Mlodzik, M. and Rubin, G. M. (1991). Cell-fate determination in the developing *Drosophila* eye: role of the *rough* gene. *Development* **112**, 703-712.
- Higashijima, S., Kojima, T., Michiue, T., Ishimaru, S., Emori, Y. and Saigo, K. (1992). Dual *Bar* homeo box genes of *Drosophila* required in two photoreceptor cells, R1 and R6, and primary pigment cells for normal eye development. *Genes Dev.* **6**, 50-60.
- Kimmel, B. E., Heberlein, U. and Rubin, G. M. (1990). The homeo domain protein *rough* is expressed in a subset of cells in the developing *Drosophila* eye where it can specify photoreceptor cell subtype. *Genes Dev.* **4**, 712-727.
- Krämer, H., Cagan, R. L. and Zipursky, S. L. (1991). Interaction of *bride of sevenless* membrane-bound ligand and the *sevenless* tyrosine-kinase receptor. *Nature* **352**, 207-212.
- Ladiaz, J. A. A. and Karathanasis, S. K. (1991). Regulation of the Apolipoprotein AI gene by ARP-1, a novel member of the steroid receptor superfamily. *Science* **251**, 561-565.
- Lawrence, P. A. and Green, S. M. (1979). Cell lineage in the developing retina of *Drosophila*. *Dev. Biol.* **71**, 142-152.
- Miyajima, N., Kadowaki, Y., Fukushige, S., Shimizu, S., Semba, K., Yamanashi, Y., Matsubara, K., Toyoshima, K. and Yamamoto, T. (1988). Identification of two novel members of *erbA* superfamily by molecular cloning: the gene products of the two are highly related to each other. *Nucl. Acids Res.* **16**, 11057-11074.
- Mlodzik, M., Baker, N. E. and Rubin, G. M. (1990a). Isolation and expression of *scabrous*, a gene regulating neurogenesis in *Drosophila*. *Genes Dev.* **4**, 1848-1861.
- Mlodzik, M., Hiromi, Y., Weber, U., Goodman, C. S. and Rubin, G. M. (1990b). The *Drosophila seven-up* gene, a member of the steroid receptor gene superfamily, controls photoreceptor cell fates. *Cell* **60**, 211-224.
- Mlodzik, M., Hiromi, Y., Goodman, C. S. and Rubin, G. M. (1992). The presumptive R7 cell of the developing *Drosophila* eye receives positional information independent of *sevenless*, *boss* and *sina*. *Mech. Dev.* **37**, 37-42.
- Mullins, M. C. and Rubin, G. M. (1991). Isolation of temperature-sensitive mutations of the tyrosine kinase receptor *sevenless* (*sev*) in *Drosophila* and their use in determining its time of action. *Proc. Nat. Acad. Sci. USA* **88**, 9387-9391.
- Oro, A. E., McKeown, M. and Evans, R. M. (1992). The *Drosophila* retinoid X receptor homolog *ultraspiracle* functions in both female reproduction and eye morphogenesis. *Development* **115**, 449-462.
- Ready, D. F. (1989). A multifaceted approach to neural development. *Trends in Neurosci.* **12**, 102-110.
- Ready, D. F., Hanson, T. E. and Benzer, S. (1976). Development of the *Drosophila* retina, a neurocrystalline lattice. *Dev. Biol.* **53**, 217-240.
- Reinke, R. and Zipursky, S. L. (1988). Cell-cell interaction in the *Drosophila* retina: the *bride of sevenless* gene is required in photoreceptor cell R8 for R7 cell development. *Cell* **55**, 321-330.
- Robertson, H. M., Preston, C. R., Phillis, R. W., Johnson-Schlitz, D. M., Denz, W. K. and Engels, W. R. (1988). A stable genomic source of P element transposase in *Drosophila melanogaster*. *Genetics* **118**, 461-470.
- Robinow, S. and White, K. (1991). Characterization and spatial distribution of the ELAV protein during *Drosophila melanogaster* development. *J. Neurobiol.* **22**, 443-461.
- Rogge, R. D., Karlovich, C. A. and Banerjee, U. (1991). Genetic dissection of a neurodevelopmental pathway: *son of sevenless* functions downstream of the *sevenless* and EGF receptor tyrosine kinases. *Cell* **64**, 39-48.
- Rogge, R. D., Cagan, R., Majumdar, A., Dulaney, T. and Banerjee, U. (1992). Neuronal development in the *Drosophila* retina: the *sextra* gene defines an inhibitory component in the developmental pathway of R7 photoreceptor cells. *Proc. Nat. Acad. Sci. USA* **89**, 5271-5275.
- Rubin, G. M. (1991). Signal transduction and the fate of the R7 photoreceptor in *Drosophila*. *Trends Genet.* **7**, 372-377.
- Simon, M. A., Bowtell, D. D. L. and Rubin, G. M. (1989). Structure and activity of the *sevenless* protein: A protein tyrosine kinase receptor required for photoreceptor development in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **86**, 8333-8337.
- Simon, M. A., Bowtell, D. D. L., Dodson, G. S., Laverty, T. R. and Rubin, G. M. (1991). Ras1 and a putative guanine nucleotide exchange factor perform crucial steps in signalling by the *sevenless* protein tyrosine kinase. *Cell* **67**, 701-716.
- Steingrímsson, E., Pignoni, F., Liaw, G.-J. and Lengyel, J. A. (1991). Dual role of the *Drosophila* pattern gene *tailless* in embryonic termini. *Science* **254**, 418-421.
- Thummel, C. S. and Pirrotta, V. (1991). New CaSpeR P-element vectors. *Drosophila Information News Letter* **2**.
- Tomlinson, A. (1988). Cellular interactions in the developing *Drosophila* eye. *Development* **104**, 183-189.
- Tomlinson, A. and Ready, D. F. (1986). *Sevenless*: a cell-specific homeotic mutation of the *Drosophila* eye. *Science* **231**, 400-402.
- Tomlinson, A. and Ready, D. F. (1987a). Neuronal differentiation in the *Drosophila* ommatidium. *Dev. Biol.* **120**, 366-376.
- Tomlinson, A. and Ready, D. F. (1987b). Cell fate in the *Drosophila* ommatidium. *Dev. Biol.* **123**, 264-275.
- Tomlinson, A., Bowtell, D. D. L., Hafen, E. and Rubin, G. M. (1987). Localization of the *sevenless* protein, a putative receptor for positional information, in the eye imaginal disc of *Drosophila*. *Cell* **51**, 143-150.
- Tomlinson, A., Kimmel, B. E. and Rubin, G. M. (1988). *rough*, a *Drosophila* homeobox gene required in photoreceptors R2 and R5 for inductive interactions in the developing eye. *Cell* **55**, 771-784.
- Van Vactor, D. L. Jr., Cagan, R. L., Kramer, H. and Zipursky, S. L. (1991). Induction in the developing compound eye of *Drosophila*: multiple mechanisms restrict R7 induction to a single retinal precursor cell. *Cell* **67**, 1145-1155.
- Wang, L.-H., Tsai, S.-Y., Cook, R. G., Beattie, W. G., Tsai, M.-J. and O'Malley, B. W. (1989). COUP transcription factor is a member of the steroid receptor superfamily. *Nature* **340**, 163-166.
- Wolf, T. and Ready, D. F. (1991). The beginning of pattern formation in the *Drosophila* compound eye: the morphogenetic furrow and the second mitotic wave. *Development* **113**, 841-850.
- Yao, T.-P., Segraves, W. A., Oro, A. E., McKeown, M. and Evans, R. M. (1992). *Drosophila* ultraspiracle modulates ecdysone receptor function via heterodimer formation. *Cell* **71**, 63-72.
- Zipursky, S. L., Venkatesh, T. R., Teplow, D. B. and Benzer, S. (1984). Neuronal development in the *Drosophila* retina: monoclonal antibodies as molecular probes. *Cell* **36**, 15-26.