

Novel genes expressed differentially in ascidians with alternate modes of development

Billie J. Swalla^{1,2,*}, Kazuhiro W. Makabe^{1,3,†}, Noriyuki Satoh³ and William R. Jeffery^{1,2}

¹Department of Zoology and Bodega Marine Laboratory, University of California, Davis, PO Box 247, Bodega Bay, CA 94923, USA

²Station Biologique, Place Georges Teissier, 29680 Roscoff, France

³Department of Zoology, Kyoto University, Kyoto, 606, Japan

*Author for correspondence at address 1

†Present address: Division of Biology, California Institute of Technology, Pasadena, CA 91125, USA

SUMMARY

We have used a subtractive procedure to isolate cDNA clones encoding genes expressed differentially in ascidian species with alternate modes of development. The ascidians used in this study were *Molgula oculata*, which develops a tailed (urodele) larva, and *Molgula occulta*, which develops a tailless (anural) larva. Two of the identified clones, *Uro-2* and *Uro-11*, are described. Southern blots show that the *Uro-2* and *Uro-11* genes are present in both species, but the corresponding mRNAs are expressed preferentially in the urodele species. In situ hybridization showed that *Uro-2* and *Uro-11* transcripts accumulate in small oocytes during oogenesis. The maternal *Uro-2* and *Uro-11* transcripts were distributed throughout the oocyte cytoplasm. Transcript concentrations declined during vitellogenesis, but mature eggs still contain detectable levels of *Uro-2* and *Uro-11* mRNA. After fertilization, the maternal *Uro-2* and *Uro-11* transcripts were localized in the ectoplasm of uncleaved zygotes and mostly entered the ectoderm cells during cleavage. The *Uro-2* gene appears to produce only maternal transcripts. In contrast, the *Uro-11* gene may also produce zygotic transcripts, which accumulate

between gastrulation and neurulation in posterior epidermis, neural and tail muscle cells. Zygotic expression of the *Uro-11* gene was not detected in embryos of the anural species. The deduced amino acid sequences of the *Uro-2* and *Uro-11* cDNAs suggest that they encode novel basic proteins with distinctive structural features. The predicted *Uro-2* protein contains a leucine zipper motif, suggesting that it may dimerize with another protein. The predicted *Uro-11* protein contains a nuclear localization signal, a region with similarity to part of the DNA-binding motif in the bacterial histone-like HU and IHF proteins, 12 repeats of the proposed DNA-binding motif S(T)PXX, and a potential zinc finger of the C₆ or C₆H₂ class, suggesting that it may be a DNA-binding protein. The *Uro-2* and *Uro-11* proteins are candidates for regulatory factors involved in the evolutionary transition from urodele to anural development.

Key words: alternate modes of development, evolution, subtractive cloning, *Uro* genes, ascidians

INTRODUCTION

Little is known about the molecular mechanisms underlying evolutionary changes in development. Closely related species with different modes of development are attractive systems to investigate this problem (Raff, 1987). Ascidians are urochordates that have evolved alternate modes of development (reviewed in Jeffery and Swalla, 1992a). Most ascidian species form a tailed (urodele) larva, which is the dispersal phase in the life cycle. The urodele larva consists of a head, containing a brain with a pigmented sensory organ(s), and a tail, containing a notochord flanked by bands of striated muscle cells. The larval tail is formed by concerted movements of the prospective notochord (Cloney, 1964; Miyamoto and Crowther, 1985), muscle and

epidermal cells in the posterior region of the embryo. As the tail elongates, myoblasts differentiate into muscle cells and a swimming tadpole larva is formed. The tail muscle cells develop autonomously, mediated by determinants segregated from the egg into the myoblast lineages (reviewed in Satoh et al., 1990; Swalla, 1992). In contrast, the brain sensory cells are specified by inductive signals (Nishida and Satoh, 1989), probably arising from presumptive notochord cells during gastrulation (reviewed in Venuti and Jeffery, 1989). The notochord cells themselves may be specified by an earlier induction during cleavage (Nishida, 1992).

Anural development is an alternate mode of development in ascidians, which results in the formation of a tailless (anural) embryo (reviewed in Jeffery and Swalla, 1990). The anural embryo lacks many of the typical urodele features

and forms an immotile larva, which develops directly into an adult. Fewer than 20 ascidian species have been shown to exhibit anural development and almost all of these species are members of the family Molgulidae (reviewed in Jeffery and Swalla, 1990). Urodele development is thought to be the ancestral mode of development in ascidians (Berrill, 1931). This viewpoint is supported by the retention of vestigial urodele features in embryos of some anural species (Whittaker, 1979a; Swalla and Jeffery, 1990, 1991). Anural ascidians appear to have evolved in habitats such as subtidal sand and mud flats in which there is less selective advantage for a larval dispersal phase.

The mechanisms underlying the transition from urodele to anural development have been investigated in two closely related species of ascidians with different modes of development (Swalla and Jeffery, 1990; Jeffery and Swalla, 1991). *Molgula oculata* shows conventional urodele development, whereas *Molgula occulta* shows anural development. Fertilization, cleavage, gastrulation and neurulation are similar in both species, but induction of brain sensory cells and tail morphogenesis does not occur in embryos of the anural species (Swalla and Jeffery, 1990). Remarkably, however, when eggs of the anural species are fertilized with sperm of the urodele species, hybrids develop some of the ancestral urodele features, including a brain sensory cell and a short tail containing a notochord (Swalla and Jeffery, 1990). Expression of the urodele genome in the anural zygote is responsible for restoring these urodele features (Jeffery and Swalla, 1992b). In contrast, the reciprocal cross, in which eggs of the urodele species are fertilized with sperm of the anural species results in hybrids with urodele development (Jeffery and Swalla, 1992a). These results suggest that loss-of-function mutations in zygotic genes are responsible for some of the developmental changes in the anural species. However, muscle cells did not differentiate in hybrids with restored urodele features, suggesting that maternal changes are also important in the transition to anural development. Consistent with this possibility, the myoplasm, an egg cytoplasmic region that is segregated to muscle lineages in ascidian embryos (see Swalla, 1992), is modified in eggs of anural species (Swalla et al., 1991; Jeffery and Swalla, 1992b).

The identification of regulatory genes expressed differentially in urodele and anural ascidians is needed to define the molecular mechanisms underlying the evolutionary transition to anural development. Here we identify and characterize two cDNAs cloned by subtractive procedures encoding *Uro* (urodele-specific) genes that are expressed during development in the urodele ascidian *M. oculata*, but are inactive or down-regulated in the anural ascidian *M. occulta*. The deduced *Uro* proteins contain potential leucine zipper and DNA-binding motifs and are candidates for regulatory factors involved in changing the mode of larval development.

MATERIALS AND METHODS

Biological materials

The urodele ascidian *Molgula oculata* and the anural ascidian *Molgula occulta* were obtained by dredging sand flats near Point

de Blosson, Roscoff, France. Animals were maintained in tanks of running sea water and gametes were obtained from dissected gonads. Cultures of mature eggs obtained in this way also contain large numbers of immature oocytes. Insemination of eggs, culture of embryos and developmental times have been described previously (Swalla and Jeffery, 1990).

Isolation of DNA and RNA

The method of Davis et al. (1986) was used to prepare genomic DNA for southern analysis. Total RNA was obtained by homogenization of dissected gonads or embryos using the guanidine isothiocyanate method (March et al., 1985). Poly(A)⁺ RNA was isolated by purification with Oligotex-dT30 beads (Roche, Tokyo, Japan).

Preparation and PCR amplification of cDNA libraries

cDNA was synthesized from *M. oculata* and *M. occulta* gonad poly(A)⁺ RNA using a Riboclone cDNA kit (Promega, Madison, WI). Double-stranded cDNA was size fractionated on a Sepharose CL-4B (Sigma Chemical Company, St. Louis, MO) column and fractions greater than 300 bp were collected. cDNA prepared from *M. oculata* gonad mRNA was directionally cloned into the *EcoRI-XhoI* site of a lambda Uni-ZAP vector (Stratagene, La Jolla, CA). *M. occulta* cDNA was ligated into the *EcoRI* site of a lambda ZAP-II vector (Stratagene) in either direction.

PCR reactions were performed in 100 µl volumes using Taq polymerase (Promega) and a Perkin-Elmer Thermal Cycler (Perkin-Elmer Corporation; Norwalk, Conn.). Final concentrations of the reaction mixture were 10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl₂ (pH 9 at 25°C), 0.1% Triton X-100 (Promega). A 68.5 µl aliquot of the *M. oculata* or *M. occulta* cDNA library was boiled in water and added to the PCR reaction at a final concentration of 10⁵ pfu/µl. Each PCR reaction included 100 pmol of T3 primer (5'-ATTAACCTCACTAAAGGGA-3') and T7 primer (5'-GCGTAATACGACTACTATA-3') for the *M. oculata* library or SK20 primer (5'-CCGCTCTAGAACTAGTGGAT-3') and T7 primer for the *M. occulta* library. The primers were synthesized using an ABI 380B-02 DNA synthesizer (Applied Biosystems Inc., Foster City, CA). Amplification used 15 cycles of 1 minute at 94°C, 2 minutes at 55°C, and 3 minutes at 72°C. One fifth aliquot of the amplified inserts was then reamplified in 100 µl of a fresh reaction mixture for 4 cycles and chased by 2 additional cycles that omitted the 94°C incubation (Timblin et al., 1990). After the final cycle, the sample was incubated at 72°C for 15 minutes, then chilled to 4°C.

Subtractive hybridization and library screening

To synthesize RNA from the amplified libraries, template DNA from each library was amplified as described above, extracted with phenol and precipitated with ethanol, and a quarter of the DNA was dissolved in 20 µl of RNA synthesis mixture according to Van Gelder et al. (1990). 80 units of T7 RNA polymerase (Bethesda Research Laboratories, Bethesda, MD) was used to synthesize antisense RNA from the *M. oculata* cDNA library. After incubation for 2 hours at 37°C, 1 unit of RQ1 DNase (Promega; Madison, WI) was added to digest the DNA template. The *M. oculata* library RNA was converted into cDNA using the T3 primer, and RNA was subsequently hydrolyzed in 1 N NaOH, 0.5% SDS. Because the *M. occulta* cDNA library was not directionally cloned, RNA prepared using the T7 promoter consisted of both sense and antisense messages. The *M. occulta* library RNA was labeled with photoactivatable biotin (Clontech Laboratories, Inc., Palo Alto, CA) by use of a PAB sunlamp (Clontech). Subtractive hybridization, in which 1.46 µg of *M. oculata* single-stranded cDNA and 20 µg of *M. occulta* library RNA were mixed, was performed according to the procedure of Sive and St. John (1988), except that 3 rounds of hybridization were done. After hybridization, 5 µg of streptavidin was added and the mixture was phenol extracted twice to remove the biotin-labeled transcripts. The subtracted cDNA (92 ng) was

then labelled with [α - 32 P]dCTP (3000 Ci/mmol, Amersham, Arlington Heights, IL) by random primer labelling (BMB; Boehringer Mannheim Biochemicals, Indianapolis, IN) and used as a probe to screen the *M. oculata* cDNA library.

Screening was performed in 60 ml of hybridization solution (50% formamide, 6 \times SSPE, 5 \times Denhardt's, 0.5% SDS, 100 μ g/ml salmon sperm DNA) with 2 \times 10⁶ cts/minute/ml probe. A total of 5 \times 10⁴ phage were screened at high stringency with the subtracted probe at low plaque density (5 \times 10³ per 10 \times 14 cm plate). The 24 positive *M. oculata* clones obtained in the first screen were then hybridized with a cDNA probe prepared from RNA synthesized in vitro from the *M. occulta* cDNA library, according to the method described above, except that washes were at low stringency (5 washes at 2 \times SSC, 0.1% SDS for 30 minutes at 55°C; 4 washes at 0.1 \times SSC, 0.1% SDS for 25 minutes at 50°C), and 8 negative clones were selected. One of the 16 positive clones was also retained as a control for RNA slot blot hybridization.

The clones that screened positive with *M. oculata* cDNA but negative with *M. occulta* cDNA (*M. oculata* positive/*M. occulta* negative clones) were further characterized by RNA slot blot hybridization. RNA synthesized in vitro from the *M. oculata* and *M. occulta* cDNA libraries as described earlier was blotted onto Hybond N⁺ filter paper (Amersham) using the formamide/formaldehyde method (Sambrook et al., 1989). The filters were hybridized separately in Rapid Hybridization Buffer (Amersham) with 1 \times 10⁶ cts/minute of random primed (BMB) 32 P-labelled (3000 Ci/mmol, Amersham) cDNA. Probes were prepared from templates obtained by PCR amplification of 7 *M. oculata* positive/*M. occulta* negative clones (1 insert did not amplify) and a control clone positive for both species. Three cDNA clones, *Uro 1*, 2 and 11 (see Results for terminology of *Uro* clones), hybridized preferentially to *M. oculata* RNA; the remaining clones hybridized to the same extent as the control clone to RNA of both species.

Clones encoding full-length cytoskeletal actin genes from *M. oculata* (*MocuCA-4*) and *M. occulta* (*MoccCA-4*) were used as controls in studies of the expression patterns of the mUro genes. The derived amino acid sequences of *MocuCA-4* and *MoccCA-4* correspond to previously described ascidian cytoskeletal actin genes (Beach and Jeffery, 1990).

Sequencing and computer analysis

Selected cDNA clones were sequenced on both strands using [35 S]-dATP (800 Ci/mmol; NEN; New England Nuclear, Boston, MA) by the dideoxy chain termination method (Sanger et al. 1977) with Sequenase (USB; United States Biochemical Corp., Cleveland, OH). Oligonucleotide primers were made on a Pharmacia LKB Gene Assembler Plus (Pharmacia Biosystems, Inc.; Piscataway, NJ) to create overlapping sequence information. Synthesized oligonucleotides were deprotected with NH₄OH and purified with Oligo-clean (USB), then diluted to 25 ng/ml with distilled water. The sequences were read and compared with the MacVector Program (IBI-A Kodak Company; New Haven, CT) on a MacIntosh IICI computer, and translated protein sequences were compared to sequences present in the National Biomedical Research Foundation protein database (IBI) by use of MacVector (IBI). Sequences were also compared by BLAST to the on-line GenBank protein sequences available through the National Library of Medicine. Sequences are available through GenBank by Accession no. L19340 (*Uro-2*) and Accession no. L19339 (*Uro-11*).

Filter hybridizations

Genomic DNA was exhaustively digested with *Eco*RI, *Hind*III, and *Bam*HI and blotted onto nylon membrane (MSI; Fisher Scientific; San Francisco, CA). After baking under vacuum for 2 hours, blots were hybridized at 42°C for 18–24 hours in a rotating hybridization chamber (Robbins Scientific Corporation, Sunnyvale, CA).

Washes were done in decreasing concentrations of SET (1 \times SET=30 mM Tris, 2 mM EDTA, 150 mM NaCl) at increasing temperatures to obtain an appropriate signal-to-noise ratio. DNA probes for blot hybridizations were labelled with [32 P]dCTP (800–3000 Ci/mmol; NEN) using random primed labelling (USB) as described above. For some hybridizations, antisense RNA probes were synthesized in vitro with either T3 or T7 polymerase (Ambion; Austin, TX) and [32 P]UTP (3,000 Ci/mole; NEN) after linearizing the clone with the appropriate restriction enzyme.

In situ hybridization

Gonads and embryos were embedded in paraplast, sectioned at 8 μ m, attached to subbed microscope slides and subjected to in situ hybridization (Swalla and Jeffery, 1990). Single-stranded RNA was synthesized from full-length *Uro* cDNA inserts using [35 S]ATP (1000–1500 Ci/mmol; NEN) and T7 RNA polymerase (MAXIscript Kit; Ambion). Approximately 1–5 \times 10⁶ cts/minute of labelled RNA probe was added to each pretreated slide in 50 μ l of hybridization buffer (Swalla and Jeffery, 1990) containing 100 mM dithiothreitol (DTT). Coverslips were removed in 4 \times SSC with 10 mM DTT, and the slides were treated with 20 μ g/ml pancreatic RNase A for 30 minutes at 37°C. Following RNase treatment, slides were washed at room temperature for 30 minutes with 2 \times SSC and then for 45 minutes with 1 \times SSC. Autoradiography was performed with NTB-2 emulsion (IBI-Kodak) and sections were stained through the emulsion with Harris hematoxylin and eosin (Humason, 1972). Background radioactivity was assessed from hybridizations carried out with sense RNA probes.

RESULTS

Isolation of *Uro* cDNA clones

Fig. 1 shows larvae of the urodele ascidian *M. oculata* and the anural ascidian *M. occulta*. The larva of the urodele species has a pigmented brain sensory cell and a tail, whereas the larva of the anural species lacks these features. A subtractive hybridization procedure was used to identify genes that are expressed preferentially in *M. oculata* oocytes, and thus are candidates for regulatory factors that initiate the changes leading to anural development. An *M. oculata* (urodele) gonad cDNA library was screened with a *M. oculata* subtracted cDNA probe and 24 positive clones were identified. Subsequent screening of the positive clones with an *M. occulta* (anural) gonad cDNA probe yielded 8 clones that were candidates for encoding differentially expressed genes. The PCR-amplified inserts of 7 of these clones were used to probe slot blots containing RNA synthesized from the urodele and anural gonad cDNA libraries, and 3 clones were identified that hybridized preferentially with *M. oculata* RNA (data not shown). These were designated *Uro* (urodele specific) clones because of their preferential expression in *M. oculata*, the urodele species. Here we describe two of these cDNA clones: *Uro-2* and *Uro-11*.

Identification of *Uro* transcripts

The *Uro-2* and *Uro-11* transcripts were identified by hybridizing northern blots containing RNA of the urodele and anural species with probes prepared from each insert. The *Uro-2* probe detected a single 1.8 kb mRNA (Fig. 2A; lane c) and the *Uro-11* probe a major 1.8 kb transcript and a minor 2.3 kb transcript (Fig. 2B; lane c) in gonad RNA from the urodele species. Neither the *Uro-2* nor *Uro-11*



Fig. 1. A photomicrograph showing larvae of the anural ascidian *M. occulta* (top) and the urodele ascidian *M. oculata* (bottom). The anural larva lacks the neural sensory cell (melanin pigment spot) and the tail of the urodele larva. Scale bar is 50 μ m.

transcripts was detected in gonads or 7.5 hour embryos of the anural species (Fig. 2A, B; lanes a-b). Reducing the hybridization stringency to much lower levels did not result in detection of *Uro* mRNAs in gonads or zygotes of the anural species (data not shown). In addition, longer exposures of the blots shown in Fig. 2 failed to detect *Uro-2* transcripts in the anural species, although very weak 1.8 kb and 2.3 kb bands were sometimes observed in blots hybridized with the *Uro-11* probe. As a control, an identical northern blot was hybridized with a probe containing the coding region of an *M. oculata* cytoskeletal actin gene. The 1.4 kb cytoskeletal actin transcript was expressed in gonads of both species (Fig. 2C, lanes a, c), as well as in embryos of the anural species (Fig. 2C, lane b). Unfortunately, filter hybridization procedures could not be used to test for accumulation of zygotic *Uro* mRNAs in the urodele species because the embryo cultures also contain immature oocytes (see Materials and Methods; Fig. 6), which accumulate high levels of both maternal *Uro* transcripts. We conclude that the *Uro-2* and *Uro-11* clones encode transcripts that are preferentially expressed in gonads of the urodele species.

***Uro* genes are present in the urodele and anural species**

The preferential accumulation of *Uro-2* and *Uro-11* transcripts in gonads of the urodele species could be caused by differential gene expression or by modification or deletion




Fig. 2. Identification of *Uro* transcripts by northern blot hybridization. Each lane was loaded with 10 μ g of total RNA isolated from *M. occulta* gonads (a), *M. occulta* 7.5 hour embryos (b), or *M. oculata* gonads (c) and hybridized with labeled RNA probes prepared from either *Uro-2* (A), *Uro-11* (B), or *MoccCA-4* (C) plasmids. The approximate size of the transcripts (see text) was determined by comparing the bands to RNA standards. Northern blots were washed first in 2 \times SET at 42 $^{\circ}$ C and then at increasingly higher stringencies. The blots in this photograph were washed in 1 \times SET at 60 $^{\circ}$ C. Exposure was 4 days in A,B and 1 day in C.

of the corresponding genes in the anural species. The presence of the *Uro-2* and *Uro-11* genes in the urodele and anural species was examined by Southern blot hybridization (Fig. 3). Only one or two labeled *M. oculata* DNA fragments were detected after hybridization with *Uro-2* (Fig. 3B) or *Uro-11* (Fig. 3D) inserts, suggesting that the corresponding genes are single copy in the urodele species. Hybridization was also observed in Southern blots containing genomic DNA fragments from the anural species (Fig. 3A,C). One or two labeled DNA fragments were observed in *M. occulta* DNA after hybridization with the *Uro-2* insert (Fig. 3A), suggesting that the *Uro-2* gene is also single copy in the anural species. However, two or more bands were detected in every lane in blots probed with the *Uro-11* insert (Fig. 3C), suggesting that multiple copies of *Uro-11*-related genes may be present in the anural species. The results suggest that homologous *Uro-2* and *Uro-11* genes may be present in the genome of the anural species.

***Uro* gene expression during development**

Further information on the developmental expression of the *Uro* genes was obtained by in situ hybridization. Figs 4 and 5 show sectioned gonads of the urodele and anural species hybridized with *Uro-2* (Fig. 5A,B), *Uro-11* (Figs 4A,B, 5C, D) and actin (Figs 4C,D, 5E,F) RNA probes. The hermaphroditic gonads of each species contain sperm and oocytes at various stages of development. Consistent with the northern blot results (Fig. 2), *Uro-2* and *Uro-11* transcripts were present in gonads of the urodele species (Fig. 5A,C), but were undetectable (Fig. 5B) or present at very low levels (Fig. 5D) respectively in the anural species. In contrast, actin mRNA was observed in gonads of both species (Figs 4C,D, 5E,F). The *Uro-2* and *Uro-11* transcripts showed the same temporal and spatial expression patterns in gonads. Transcripts were observed in oocytes but not in accessory cells




Fig. 3. Southern blots of *M. occulta* (A,C) and *M. oculata* (B,D) genomic DNA digested with *Bam*HI (a), *Eco*RI (b), or *Hind*III (c). 10 µg of digested DNA from each species was loaded in each lane of the gels and blots were hybridized with labelled *Uro-2* (A,B) or *Uro-11* (C,D) probes and washed in 2× SSC at 50°C. Exposure was 2 days for A,C and D and 1 day for B.

surrounding the oocytes (data not shown) or in developing spermatocytes within *M. oculata* gonads (Fig. 4A). The highest concentration of *Uro-2* and *Uro-11* transcripts was seen in small previtellogenic oocytes (Fig. 5A,C). The concentration of *Uro-2* and *Uro-11* mRNA gradually declined during vitellogenesis (Fig. 5A,C). A similar decline in actin transcripts was observed in vitellogenic oocytes of both species (Fig. 5E,F). The *Uro-2* and *Uro-11* transcripts were distributed uniformly in the cytoplasm of both previtellogenic and vitellogenic oocytes (Fig. 5A,C). The results are consistent with the northern analysis described above and confirm that maternal *Uro* mRNAs accumulate preferentially in oocytes of the urodele species.

Although the concentration of *Uro* transcripts declined during oogenesis, signal above background was still present in mature oocytes (Fig. 5A,C), implying that maternal mRNA may persist in eggs and embryos. The expression and distribution of *Uro* mRNA in eggs and embryos of the urodele species was examined by in situ hybridization. The results are shown in Fig. 6. In this figure, developing embryos are shown mixed with immature oocytes (see Materials and Methods), which served as internal controls. Maternal transcripts of both *Uro* genes were present in unfertilized eggs. After fertilization and ooplasmic segregation, the maternal *Uro-2* and *Uro-11* mRNAs were localized in the ectoplasm (Conklin, 1905) of uncleaved zygotes and distributed to ectodermal blastomeres in the animal hemisphere during cleavage (Fig. 6A-D). The maternal *Uro-2*




Fig. 4. Expression of *Uro-11* and actin transcripts in *M. occulta* and *M. oculata* gonads. Sections of gonads from each species were hybridized in situ with radioactive RNA probes and are shown here at low magnification. Sections of entire *M. oculata* (A,C) and *M. occulta* (B,D) gonads hybridized with *Uro-11* (A,B) or *MocuCA-4* (C,D) RNA. In A-B, *Uro-11* transcripts are expressed in small oocytes of the urodele (A) but not the anural (B) species (arrowheads) and are absent from the testes (T) of both species. In C-D, actin transcripts are present in oocytes (arrowheads) of both species. Scale bar in A represents 500 µm; magnification is the same in A-D.

transcripts could no longer be detected by the late cleavage stages, and did not accumulate to levels above background during the remainder of embryogenesis (Fig. 6E,G,I). Thus, expression of the *Uro-2* gene appears to be strictly maternal. In contrast, *Uro-11* transcripts were detectable through the neurula stage (Fig. 6F,H), but could not be observed at later stages of embryogenesis (Fig. 6J). In other ascidian species, maternal transcripts disappear during cleavage and embryonic transcription begins by the early gastrula stage (Beach and Jeffery, 1990, 1992; Kusakabe et al., 1991). Therefore, it is likely that the *Uro-11* gene shows both maternal and zygotic expression. The putative zygotic *Uro-11* transcripts accumulated primarily in the posterior epidermis, the developing nervous system (neural folds and tube) and the presumptive myoblasts (Fig. 6D,F,H).

However, zygotic *Uro-2* and *Uro-11* transcripts were not detected in embryos of the anural species (data not shown; Fig. 2B, lane b). The results suggest that maternal *Uro-2* and *Uro-11* transcripts are localized in the zygote and segregated to ectodermal cells during cleavage. Afterwards, expression of the *Uro* genes appears to diverge: the *Uro-2* gene appears to be zygotically inactive, whereas the *Uro-11* gene may produce zygotic transcripts in posterior tissues of gastrulae and neurulae.

The predicted *Uro-2* protein contains a leucine zipper motif

The nucleotide and predicted amino acid sequences of the *Uro-2* clone are shown in Fig. 7A. The cDNA clone contains a single opening reading frame of 1092 nucleotides which




Fig. 5. Expression of *Uro-2*, *Uro-11* and actin transcripts in developing oocytes in *M. oculata* (A,C,E) and *M. occulta* (B,D,F) gonads hybridized in situ with *Uro-2* (A,B), *Uro-11* (C,D) or *MocuCA-4* (E,F) RNA probes. The *Uro-2* and *Uro-11* transcripts are distributed throughout the cytoplasm of small previtellogenic oocytes in the urodele (A,C) but not the anural (B,D) species, and gradually decrease in concentration in larger vitellogenic oocytes. The actin transcripts are present in previtellogenic oocytes of both species (E,F). Scale bar in E represents 50 μ m; magnification is the same in A-F.

is flanked by a 5' leader of 61 nucleotides and a 3' untranslated region of 684 nucleotides. The first ATG triplet is in the context most closely resembling the consensus eukaryotic translational initiation site (Kozak, 1991), and the 3' non-coding region contains a putative polyadenylation signal. The size of the *Uro-2* cDNA clone (1836 bp) is consistent with the 1.8 kb *Uro-2* transcript identified in northern blots (Fig. 2A, lane c).

The *Uro-2* nucleotide sequence predicts a basic protein

(pI 10.0) of 364 amino acids with a calculated molecular mass of 42 kD (Genbank Accession no. L19340). No significant matches were revealed between the *Uro-2* amino acid sequence and proteins represented in the protein databases. The *Uro-2* protein contains a tyrosine kinase phosphorylation motif (KKIDPHSY) and a region containing mostly hydrophobic amino acids (Fig. 7A). The hydrophobic region lacks charged amino acids and is extensive enough to be a transmembrane domain, although a conventional signal sequence (von Heijne, 1986) is not present at the N terminus (Fig. 7B). The most notable feature of the *Uro-2* protein is a heptad repeat of four leucine residues in the C-terminal region (Fig. 7A). This region may be a leucine zipper, the dimerization motif present in certain transcription factors (Landschultz et al., 1988) and integral membrane proteins (Buckland and Wild, 1989; McCormack et al., 1989; White and Weber, 1989; Inoue et al., 1991). Although a short basic region precedes the putative leucine zipper in the *Uro-2* protein, the sequence lacks the conserved basic domain typical of the bZIP family of transcription factors. Thus, the *Uro-2* gene may encode a novel leucine zipper protein.

The predicted *Uro-11* protein contains DNA-binding motifs

The *Uro-11* nucleotide and predicted amino acid sequences are shown in Fig. 8. The cDNA clone contains an open reading frame of 1674 nucleotides, a 5' leader of 428 nucleotides, and a 3' untranslated region of 192 nucleotides (GenBank Accession no. L19339). The first ATG is in the most reasonable context for translational initiation (Kozak, 1991) and a polyadenylation signal is present in the 3' untranslated region. The inferred *Uro-11* mRNA (2294 nucleotides) is sufficient to encode either the 1.8 or 2.3 kb transcripts identified by northern analysis (Fig. 2B, lane c).

The *Uro-11* sequence predicts a basic protein (pI 10.0) of 558 amino acids with a calculated molecular mass of 62 kD.




Fig. 6. Expression of *Uro-2* and *Uro-11* transcripts in *M. oculata* embryos hybridized in situ with *Uro-2* (A,C,E,G,I) and *Uro-11* (B,D,F,H,J) RNA probes. (A,B) Sections of 2-cell embryos showing *Uro-2* and *Uro-11* transcripts localized in the ectoplasm. The animal pole is at the top in both figures. In B, the 2-cell embryo is flanked by a labeled immature oocyte (lower right). (C,D) Sections through the animal hemisphere and parallel to the anteroposterior axis (top to bottom) of 32-64 cell embryos showing *Uro-2* and *Uro-11* transcripts in ectoderm cells. In C, the concentration of *Uro-2* mRNA has decreased relative to A and the adjacent (right) labeled immature oocyte. In D, the concentration of *Uro-11* mRNA has increased relative to B, suggesting that zygotic *Uro-11* transcripts have started to accumulate. (E,F) Parasagittal sections through the anteroposterior axis (right to left) of mid-gastrulae. In E, *Uro-2* transcripts are undetectable at the gastrula stage. In contrast, in F, *Uro-11* transcripts are located in the posterior epidermis (E) and presumptive muscle (M) cells of a gastrula. (G,H) Cross-sections through the dorsoventral axis (top to bottom) of neurulae showing lack of *Uro-2* transcripts (G) and *Uro-11* transcript accumulation in the neural folds (N) and neural tube in two neurulae (H). In G, there is a labeled immature oocyte at the upper right side of the neurula. (I,J) Parasagittal sections through early tailbud stage embryos showing lack of *Uro-2* (I) and *Uro-11* (J) transcripts. Scale bar in A represents 20 μ m; magnification is the same in each frame.

A

```

661 668 675 682 689 696 703 710 717 724 731 738 745 752 759 766 773 780 787 794 801 808 815 822 829 836 843 850 857 864 871 878 885 892 900 907 914 921 928 935 942 949 956 963 970 977 984 991 998 1005 1012 1019 1026 1033 1040 1047 1054 1061 1068 1075 1082 1089 1096 1103 1110 1117 1124 1131 1138 1145 1152 1159 1166 1173 1180 1187 1194 1201 1208 1215 1222 1229 1236 1243 1250 1257 1264 1271 1278 1285 1292 1299 1306 1313 1320 1327 1334 1341 1348 1355 1362 1369 1376 1383 1390 1397 1404 1411 1418 1425 1432 1439 1446 1453 1460 1467 1474 1481 1488 1495 1502 1509 1516 1523 1530 1537 1544 1551 1558 1565 1572 1579 1586 1593 1600 1607 1614 1621 1628 1635 1642 1649 1656 1663 1670 1677 1684 1691 1698 1705 1712 1719 1726 1733 1740 1747 1754 1761 1768 1775 1782 1789 1796 1803 1810 1817 1824 1831 1838 1845 1852 1859 1866 1873 1880 1887 1894 1901 1908 1915 1922 1929 1936 1943 1950 1957 1964 1971 1978 1985 1992 2000 2007 2014 2021 2028 2035 2042 2049 2056 2063 2070 2077 2084 2091 2098 2105 2112 2119 2126 2133 2140 2147 2154 2161 2168 2175 2182 2189 2196 2203 2210 2217 2224 2231 2238 2245 2252 2259 2266 2273 2280 2287 2294 2301 2308 2315 2322 2329 2336 2343 2350 2357 2364 2371 2378 2385 2392 2399 2406 2413 2420 2427 2434 2441 2448 2455 2462 2469 2476 2483 2490 2497 2504 2511 2518 2525 2532 2539 2546 2553 2560 2567 2574 2581 2588 2595 2602 2609 2616 2623 2630 2637 2644 2651 2658 2665 2672 2679 2686 2693 2700 2707 2714 2721 2728 2735 2742 2749 2756 2763 2770 2777 2784 2791 2798 2805 2812 2819 2826 2833 2840 2847 2854 2861 2868 2875 2882 2889 2896 2903 2910 2917 2924 2931 2938 2945 2952 2959 2966 2973 2980 2987 2994 3001 3008 3015 3022 3029 3036 3043 3050 3057 3064 3071 3078 3085 3092 3099 3106 3113 3120 3127 3134 3141 3148 3155 3162 3169 3176 3183 3190 3197 3204 3211 3218 3225 3232 3239 3246 3253 3260 3267 3274 3281 3288 3295 3302 3309 3316 3323 3330 3337 3344 3351 3358 3365 3372 3379 3386 3393 3400 3407 3414 3421 3428 3435 3442 3449 3456 3463 3470 3477 3484 3491 3498 3505 3512 3519 3526 3533 3540 3547 3554 3561 3568 3575 3582 3589 3596 3603 3610 3617 3624 3631 3638 3645 3652 3659 3666 3673 3680 3687 3694 3701 3708 3715 3722 3729 3736 3743 3750 3757 3764 3771 3778 3785 3792 3799 3806 3813 3820 3827 3834 3841 3848 3855 3862 3869 3876 3883 3890 3897 3904 3911 3918 3925 3932 3939 3946 3953 3960 3967 3974 3981 3988 3995 4002 4009 4016 4023 4030 4037 4044 4051 4058 4065 4072 4079 4086 4093 4100 4107 4114 4121 4128 4135 4142 4149 4156 4163 4170 4177 4184 4191 4198 4205 4212 4219 4226 4233 4240 4247 4254 4261 4268 4275 4282 4289 4296 4303 4310 4317 4324 4331 4338 4345 4352 4359 4366 4373 4380 4387 4394 4401 4408 4415 4422 4429 4436 4443 4450 4457 4464 4471 4478 4485 4492 4500 4507 4514 4521 4528 4535 4542 4549 4556 4563 4570 4577 4584 4591 4598 4605 4612 4619 4626 4633 4640 4647 4654 4661 4668 4675 4682 4689 4696 4703 4710 4717 4724 4731 4738 4745 4752 4759 4766 4773 4780 4787 4794 4801 4808 4815 4822 4829 4836 4843 4850 4857 4864 4871 4878 4885 4892 4899 4906 4913 4920 4927 4934 4941 4948 4955 4962 4969 4976 4983 4990 4997 5004 5011 5018 5025 5032 5039 5046 5053 5060 5067 5074 5081 5088 5095 5102 5109 5116 5123 5130 5137 5144 5151 5158 5165 5172 5179 5186 5193 5200 5207 5214 5221 5228 5235 5242 5249 5256 5263 5270 5277 5284 5291 5298 5305 5312 5319 5326 5333 5340 5347 5354 5361 5368 5375 5382 5389 5396 5403 5410 5417 5424 5431 5438 5445 5452 5459 5466 5473 5480 5487 5494 5501 5508 5515 5522 5529 5536 5543 5550 5557 5564 5571 5578 5585 5592 5600 5607 5614 5621 5628 5635 5642 5649 5656 5663 5670 5677 5684 5691 5698 5705 5712 5719 5726 5733 5740 5747 5754 5761 5768 5775 5782 5789 5796 5803 5810 5817 5824 5831 5838 5845 5852 5859 5866 5873 5880 5887 5894 5901 5908 5915 5922 5929 5936 5943 5950 5957 5964 5971 5978 5985 5992 6000 6007 6014 6021 6028 6035 6042 6049 6056 6063 6070 6077 6084 6091 6098 6105 6112 6119 6126 6133 6140 6147 6154 6161 6168 6175 6182 6189 6196 6203 6210 6217 6224 6231 6238 6245 6252 6259 6266 6273 6280 6287 6294 6301 6308 6315 6322 6329 6336 6343 6350 6357 6364 6371 6378 6385 6392 6400 6407 6414 6421 6428 6435 6442 6449 6456 6463 6470 6477 6484 6491 6498 6505 6512 6519 6526 6533 6540 6547 6554 6561 6568 6575 6582 6589 6596 6603 6610 6617 6624 6631 6638 6645 6652 6659 6666 6673 6680 6687 6694 6701 6708 6715 6722 6729 6736 6743 6750 6757 6764 6771 6778 6785 6792 6800 6807 6814 6821 6828 6835 6842 6849 6856 6863 6870 6877 6884 6891 6898 6905 6912 6919 6926 6933 6940 6947 6954 6961 6968 6975 6982 6989 6996 7003 7010 7017 7024 7031 7038 7045 7052 7059 7066 7073 7080 7087 7094 7101 7108 7115 7122 7129 7136 7143 7150 7157 7164 7171 7178 7185 7192 7200 7207 7214 7221 7228 7235 7242 7249 7256 7263 7270 7277 7284 7291 7298 7305 7312 7319 7326 7333 7340 7347 7354 7361 7368 7375 7382 7389 7396 7403 7410 7417 7424 7431 7438 7445 7452 7459 7466 7473 7480 7487 7494 7501 7508 7515 7522 7529 7536 7543 7550 7557 7564 7571 7578 7585 7592 7600 7607 7614 7621 7628 7635 7642 7649 7656 7663 7670 7677 7684 7691 7698 7705 7712 7719 7726 7733 7740 7747 7754 7761 7768 7775 7782 7789 7796 7803 7810 7817 7824 7831 7838 7845 7852 7859 7866 7873 7880 7887 7894 7901 7908 7915 7922 7929 7936 7943 7950 7957 7964 7971 7978 7985 7992 8000 8007 8014 8021 8028 8035 8042 8049 8056 8063 8070 8077 8084 8091 8098 8105 8112 8119 8126 8133 8140 8147 8154 8161 8168 8175 8182 8189 8196 8203 8210 8217 8224 8231 8238 8245 8252 8259 8266 8273 8280 8287 8294 8301 8308 8315 8322 8329 8336 8343 8350 8357 8364 8371 8378 8385 8392 8400 8407 8414 8421 8428 8435 8442 8449 8456 8463 8470 8477 8484 8491 8498 8505 8512 8519 8526 8533 8540 8547 8554 8561 8568 8575 8582 8589 8596 8603 8610 8617 8624 8631 8638 8645 8652 8659 8666 8673 8680 8687 8694 8701 8708 8715 8722 8729 8736 8743 8750 8757 8764 8771 8778 8785 8792 8800 8807 8814 8821 8828 8835 8842 8849 8856 8863 8870 8877 8884 8891 8898 8905 8912 8919 8926 8933 8940 8947 8954 8961 8968 8975 8982 8989 8996 9003 9010 9017 9024 9031 9038 9045 9052 9059 9066 9073 9080 9087 9094 9101 9108 9115 9122 9129 9136 9143 9150 9157 9164 9171 9178 9185 9192 9200 9207 9214 9221 9228 9235 9242 9249 9256 9263 9270 9277 9284 9291 9298 9305 9312 9319 9326 9333 9340 9347 9354 9361 9368 9375 9382 9389 9396 9403 9410 9417 9424 9431 9438 9445 9452 9459 9466 9473 9480 9487 9494 9501 9508 9515 9522 9529 9536 9543 9550 9557 9564 9571 9578 9585 9592 9600 9607 9614 9621 9628 9635 9642 9649 9656 9663 9670 9677 9684 9691 9698 9705 9712 9719 9726 9733 9740 9747 9754 9761 9768 9775 9782 9789 9796 9803 9810 9817 9824 9831 9838 9845 9852 9859 9866 9873 9880 9887 9894 9901 9908 9915 9922 9929 9936 9943 9950 9957 9964 9971 9978 9985 9992 10000
    
```

B




Fig. 7. (A) The nucleotide and predicted amino acid sequence of *Uro-2*. The putative tyrosine kinase phosphorylation site is shaded, the hydrophobic domain is boxed, the leucine zipper motif is indicated by a single underline with boxed heptad leucine residues, and the polyadenylation signal is indicated by a double underline. GenBank Accession no. L19340. (B) Hydrophobicity plot (Kyte and Doolittle, 1982) of the predicted *Uro-2* protein showing the central hydrophobic domain. The window size is 11.

No significant similarity was revealed by comparing the *Uro-11* amino acid sequence to proteins in the protein databases. The *Uro-11* protein contains a putative nuclear localization signal (KKKTK; Kalderon et al., 1984) and several potential DNA-binding motifs (Fig. 8). First, an N-terminal region of the protein matches the consensus of a predicted DNA-binding motif (Fig. 9) present in the bacterial histone-like proteins HU (Goshima et al., 1990) and IHF (Nash, 1990). HU and IHF are small proteins that bind to DNA as dimers via an extended arm structure consisting of a two-stranded antiparallel ribbon (reviewed in Dijk and Reinhardt, 1986). The turns (GFG and PXT) separating the strands of the ribbons as well as critical basic and hydrophobic residues in the N-terminal strand are conserved in the *Uro-11* protein. Second, the *Uro-11* protein contains 12 repeats of S(T)PXX, a proposed DNA-binding motif in certain transcription factors (Suzuki, 1989) and histone H2B (Lindsey and Thompson, 1992) (Fig. 8). Finally, the C-terminal region of the *Uro-11* protein contains a potential zinc finger (or cluster; Vallee et al., 1991) of the form CX₂CX₁₂CX₂CX₄CX₃C (Fig. 10). Although the number of cysteine residues would place the *Uro-11* protein

in the C₆ family of zinc finger proteins (Evans and Hollenberg, 1988), the cysteine positions are not conserved relative to the C₆ fungal transcription factors (Johnston, 1987). Histidine residues are present in the cysteine rich region of the *Uro-11* protein that could also be involved in coordination of zinc moieties. In this way, *Uro-11* resembles some of the C₆H₂ B box-containing zinc finger proteins (Reddy et al., 1992). Thus, the *Uro-11* gene may encode a DNA-binding protein.

DISCUSSION

Previous studies have shown that changes in maternal and zygotic factors are involved in the evolution of alternate modes of development in ascidians (reviewed in Jeffery and Swalla, 1992a). Here we have identified and characterized candidates for regulatory genes governing the transition from urodele to anural development. Our approach for isolating the *Uro* clones was based on the existence of two closely related ascidian species exhibiting urodele and anural development. The *Uro* clones were isolated from a




Fig. 8. The nucleotide and predicted amino acid sequence of *Uro-11*. The region with homology to part of the DNA-binding motif of the bacterial HU and IHF proteins is boxed, the cysteine-rich region containing a potential zinc finger is indicated by shading, the S(T)PXX repeats are indicated by a single (sometimes overlapping) underline and the polyadenylation signal is indicated by a double underline. The nuclear localization signal KKKTK (Kalderon et al., 1984) begins at residue 528 within the shaded region. GenBank Accession no. L19339

gonad cDNA library of the urodele species, *M. oculata*, after subtraction with a gonad cDNA library from the anural species, *M. occulta*. The *M. oculata* cDNA library was screened with a subtracted probe enriched in urodele-specific sequences to isolate clones encoding genes expressed preferentially in the urodele species. In this investigation, we have described two of these clones: *Uro-2* and *Uro-11*.

The temporal and spatial expression patterns of *Uro-2* and *Uro-11* genes are identical during oogenesis of the urodele species. Transcripts of each gene accumulate to their highest levels in previtellogenic oocytes and gradually decrease in concentration during vitellogenesis. The expression of *Uro-2* and *Uro-11* mRNA during oogenesis is similar to that of actin transcripts in the urodele and anural species and may be characteristic of messages synthesized in previtellogenic




Fig. 9. Comparison of the Uro-11 protein to part of the DNA-binding region in the bacterial HU and IHF proteins. Amino acids 90-121 of Uro-11 sequence (A) are compared to part of the DNA-binding domain of the *B. stearrowthermophilus* HBs (B), *E. coli* NS-2 (C) and *E. coli* IHF (D) proteins. (E) The consensus amino acids in part of the DNA-binding motif of the bacterial HU and IHF proteins are shown in boldface. Boxed residues indicate identical and conserved substitutions in the protein sequences. Bacterial sequences are from Dijk and Reinhardt (1986) and Miller (1984).

oocytes. A single *Uro-2* transcript was detected on northern blots, but two distinct *Uro-11* transcripts were present. The size of the *Uro-11* opening reading frame is consistent with expression of either transcript, and further experiments will be necessary to determine the relationship between the *Uro-11* transcripts and how they may be generated during RNA transcription or processing.

Ascidian eggs contain distinct cytoplasmic localizations (Conklin, 1905) and there is previous evidence that a subtype of actin mRNA is localized in the myoplasm (Jeffery et al., 1983), an egg cytoplasmic region inherited by the larval tail muscle cells. Although the maternal *Uro-2* and *Uro-11* transcripts are uniformly distributed in the oocyte cytoplasm throughout oogenesis, they become localized in the ectoplasm after fertilization and are segregated to the ectodermal cells during cleavage. This pattern of localization and segregation is similar to that of a maternal cytoskeletal actin mRNA in the ascidian *Styela clava* (Beach and Jeffery, 1990). The maternal *Uro-2* mRNA disappeared during cleavage and no further accumulation was detected during embryogenesis, indicating that this *Uro* gene shows exclusively maternal expression. In contrast, the *Uro-11* gene is present in embryos of the urodele species during gastrulation and neurulation. This suggests that there is also a zygotic expression period of the *Uro-11* gene. The accumulation of *Uro-11* transcripts in the posterior epidermis, the neural cells and the presumptive muscle cells suggests a possible function for this gene during urodele development.




Fig. 10. The proposed structure of a C₆ zinc finger in the Uro-11 protein. This structure assumes that the two histidine residues in the cysteine-rich region of the protein do not serve as zinc ligands.

After gastrulation, the larval tail is extended by concerted morphogenetic movements and differentiation of cells in the posterior region of the embryo. Interference with this process could lead to anural development. Thus, the temporal and spatial expression pattern of the *Uro-11* gene in some of the posterior embryonic cells is consistent with a role in tail formation in the urodele species.

Although *Uro* transcripts accumulate preferentially in the urodele species, the *Uro 2* and *Uro-11* genes are present in the genomes of both *M. oculata* and *M. occulta*. Therefore, the simplest explanation for lack of *Uro* transcripts in the anural species is that gene expression has been suppressed. The longer exposure times required to distinguish the *Uro-2* bands in Southern blots containing DNA from the anural species suggests that the single copy *Uro-2* gene has diverged in the anural species. An extreme possibility is that the *Uro-2* sequence detected in the genome of the anural species by Southern analysis represents hybridization with a pseudogene. In addition, the increased number of bands detected in genomic DNA of the anural species suggests that the *Uro-11* gene may have duplicated or undergone a chromosomal rearrangement. It will be interesting to investigate the sequence divergence and organization of the *Uro-2* and *Uro-11* genes in the urodele and anural species.

The *Uro-2* gene appears to encode a novel protein containing a leucine zipper motif. Leucine zippers promote protein-protein interactions in certain dimeric transcription factors (Landschultz et al., 1988) and channel-forming, integral membrane proteins (McCormack et al., 1989; Buckland and Wild, 1989; White and Weber, 1989; Inoue et al., 1991). Although a basic region precedes the leucine zipper in the *Uro-2* protein, it is not as extensive as the conserved basic domain characteristic of the bZIP family of transcription factors. The presence of a potential membrane spanning domain in the *Uro-2* protein brings up the alternate possibility that it may represent a leucine-zipper transmembrane protein. It should be pointed out, however, that the hydrophobic region and leucine zipper motif are widely separated in the *Uro-2* protein, whereas these regions are adjacent in the leucine zipper transmembrane proteins. If *Uro-2* is a transmembrane protein, its absence in eggs of the anural species could lead to structural changes in the myoplasm, a unique cytoskeletal domain linked to the egg plasma membrane (Jeffery and Meier, 1983). It has been shown previously that the organization of the myoplasm is modified in the eggs of anural ascidian species (Whittaker, 1979b; Swalla et al., 1991; Jeffery and Swalla, 1992b). Direct information on localization of the *Uro-2* protein using antibodies will be necessary to determine whether it is a nuclear protein or a component of the egg plasma membrane.

The *Uro-11* gene appears to encode a novel protein containing a potential nuclear localization signal and several different DNA-binding motifs. A region in the N-terminal portion of the predicted protein shares a part of the consensus of the DNA-binding arm in the bacterial histone-like HU and IHF proteins (Dijk and Reinhardt, 1986). Structural and biochemical studies suggest that the HU and IHF proteins interact with DNA via a two-stranded β -ribbon (Tanaka et al., 1984; Yang and Nash, 1989). Although *Uro-11* shows homology only to the N-terminal half of the HU-IHF β -ribbon, this is precisely the region known to be

important for DNA binding (Goshima et al., 1992). This is the first report of a bacterial histone-like DNA-binding motif in a eukaryotic protein, although similarities between the DNA-binding domains of the IHF protein and transcription factor TFIID have previously been recognized (Nash and Granston, 1991). Recent structural analysis of the TFIID protein indicates that its DNA-binding site forms a multi-stranded antiparallel β -sheet, which resembles a molecular 'saddle' sitting astride the DNA helix (Nicolov et al., 1992).

The Uro-11 protein also contains a potential C₆ zinc finger in its C-terminal region. Although this structure has 6 cysteine (and 2 histidine) residues, which could form a coordinated finger or cluster with 2 zinc moieties (Vallee et al., 1991), the cysteine positions are not conserved relative to known members of the C₆ zinc fingers, such as the GAL4 fungal transcription factor (Johnston et al., 1987), or the B box class of zinc fingers, some of which contain 6 cysteine and 2 histidine metal-binding ligands (Reddy et al., 1992). Therefore, the Uro-11 protein could represent a new subclass of zinc finger proteins in the C₆ or C₆H₂ family. Twelve repeats of the motif S(T)PXX are positioned between the regions of HU-IHF similarity and the potential zinc finger in the Uro-11 protein. The S(T)PXX motif has been proposed to form turn I structures that interact with the minor groove of the DNA helix (Suzuki, 1989). S(T)PXX repeats are frequently observed in transcription factors with zinc finger DNA-binding motifs, such as the *Drosophila* *krüppel* protein and the steroid hormone receptor (Suzuki, 1989), or in other DNA-binding proteins, such as histone H2B (Lindsey and Thompson, 1992), but are much less common in general proteins. Although further studies will be necessary to determine whether Uro-11 is a DNA-binding protein, it is interesting to note that this clone was identified in a library screen for mRNAs that are down regulated during development of the anural species. This result is consistent with the possibility that anural development is mediated by eliminating the expression of a transcription factor(s).

In summary, we have described potential genes that are expressed during the ontogeny of an ascidian with urodele development but not in a closely related species with anural development. Since our library screening was not exhaustive, it is possible that other genes showing this pattern of expression remain to be identified. Indeed, we have recently determined that another *Uro* clone (*Uro-1*) encodes a putative tyrosine kinase (Makabe et al., unpublished data). Interestingly, the Uro-2 and Uro-11 proteins contain potential phosphorylation sites, raising the possibility that they function in a signal transduction pathway downstream from the Uro-1 tyrosine kinase.

We thank B. Baldwin, T. Carroll, N. Ewing and K. Hadfield for technical assistance, Dr B. Blumberg for searching the Genbank database and Dr L. Meijer for use of his laboratory facilities at Roscoff. The collection staff at Station Biologique, Roscoff is thanked for their assistance in obtaining *M. occulta* and *M. oculata*. This research was supported by an NIH Postdoctoral Fellowship (HD-07493) (B. J. S.), a post-doctoral fellowship from the Japan Society for the Promotion of Science (02954052) (K. W. M.), a grant from the Monbusho International Scientific Research Program (01044077) (N. S.), and research grants from the NIH (HD-13970) and NSF (DCB-9115543) (W. R. J.).

REFERENCES

- Beach, R. L. and Jeffery, W. R. (1990). Temporal and spatial expression of a cytoskeletal actin gene in the ascidian *Styela clava*. *Dev. Genetics* **11**, 2-14.
- Beach, R. L. and Jeffery, W. R. (1992). Multiple actin genes encoding the same alpha muscle isoform are expressed differentially during ascidian development. *Dev. Biol.* **151**, 55-66.
- Berrill, N. J. (1931). Studies in tunicate development. Part II. Abbreviation of development in the Molgulidae. *Phil. Trans. R. Soc. B. Biol. Sci.* **219**, 281-346.
- Buckland, R. and Wild, F. (1989). Leucine zipper motif extends. *Nature* **338**, 547.
- Cloney, R. A. (1964). Development of the ascidian notochord. *Acta Embryol. Morphol. Exp.* **7**, 111-130.
- Conklin E. G. (1905). The organization and cell lineage of the ascidian egg. *J. Acad. Natl. Sci. Philadelphia* **13**, 1-119
- Davis, L. G., Dibner, M. D. and Batty, J. F. (1986). *Basic Methods in Molecular Biology*. New York, NY: Elsevier.
- Dijk, J. and Reinhardt, R. (1986). The structure of DNA-binding proteins from eu- and archaeobacteria. In *Bacterial Chromatin* (ed. C. O. Gualerzi and C. L. Pon), pp 185-218. Berlin: Springer-Verlag.
- Evans, R. M. and Hollenberg, S. M. (1988). Zinc fingers: gilt by association. *Cell* **52**, 1-3.
- Goshima, N., Kohno, K., Imamoto, F. and Kano, Y. (1990). HU-1 mutants of *Escherichia coli* deficient in DNA binding. *Gene* **96**, 141-145.
- Goshima, N., Inagaki, Y., Otaki, H., Tanaka, H., Hayashi, N., Imamoto, F. and Kano, Y. (1992). Chimeric HU-IHF proteins that alter DNA binding ability. *Gene* **118**, 97-102.
- Humason, G.L. (1972). *Animal Tissue Techniques*, pp 150-155. San Francisco, CA: W. H. Freeman and Company.
- Inoue, H., Takahashi, S., Fukui, K. and Miyake, Y. (1991). Genetic and molecular properties of human and rat renin-binding proteins with reference to the function of the leucine zipper motif. *J. Biochem.* **110**, 493-500.
- Jeffery, W. R. and Meier, S. (1983). A yellow crescent cytoskeletal domain in ascidian eggs and its role in embryonic development. *Dev. Biol.* **96**, 125-143.
- Jeffery, W. R. and Swalla, B. J. (1990). Anural development in ascidians: Evolutionary modification and elimination of the tadpole larva. *Sem. Dev. Biol.* **1**, 253-261.
- Jeffery, W. R. and Swalla, B. J. (1991). An evolutionary change in the muscle lineage of an anural ascidian embryo is restored by interspecific hybridization with a urodele ascidian. *Dev. Biol.* **145**, 328-337.
- Jeffery, W. R. and Swalla, B. J. (1992a). Evolution of alternate modes of development in ascidians. *BioEssays* **14**, 219-226.
- Jeffery, W. R. and Swalla, B. J. (1992b). Factors necessary for restoring an evolutionary change in an anural ascidian embryo. *Dev. Biol.* **153**, 194-205.
- Jeffery, W. R., Tomlinson, C. R. and Brodeur, R. D. (1983). Localization of actin messenger RNA during early ascidian development. *Dev. Biol.* **99**, 408-417.
- Johnston, M. (1987). A model fungal gene regulatory mechanism: the GAL genes of *Saccharomyces cerevisiae*. *Microbiol. Rev.* **51**, 458-476.
- Kalderon, D., Roberts, B. L., Richardson, W. D. and Smith, A. E. (1984). A short amino acid sequence able to specify nuclear localization. *Cell* **39**, 499-509.
- Kozak, M. (1991). Structural features in eucaryotic mRNAs that modulate the initiation of translation. *J. Biol. Chem.* **266**, 19867-19870.
- Kyte, J. and Doolittle, R. F. (1982). A simple method for displaying the hydrophobic character of a protein. *J. Mol. Biol.* **157**, 105-132.
- Kusakabe, T., Suzuki, J., Saiga, H., Jeffery, W. R., Makabe, K. W. and Satoh, N. (1991). Temporal and spatial expression of a muscle actin gene during embryogenesis of the ascidian *Halocynthia roretzi*. *Dev. Growth Differ.* **33**, 227-234.
- Landschultz, W. H., Johnson, P. and McKnight, S. L. (1988). The leucine zipper: A hypothetical structure common to a new class of DNA binding proteins. *Science* **240**, 1759-1764.
- Lindsey, G. G. and Thompson, P. (1992). S(P)XX motifs promote the interaction between the extended N-terminal tails of histone H2B with 'linker' DNA. *J. Biol. Chem.* **267**, 14622-14628.
- March, C. J., Mosley, B., Larsen, A., Cerretti, D. P., Braedt, G., Price, V., Gillis, S., Henney, C. S., Kronheim, S. R., Grabstein, K., Conlon, P. J., Hopp, T. P. and Cosman, D. (1985). Cloning, sequence and

- expression of two distinct human interleukin-1 complementary DNAs. *Nature* **315**, 641-647.
- McCormack, K., Campanelli, J. T., Ramaswami, M., Mathew, M. K., Tanouye, M. A., Iverson, L. E. and Rudy, B.** (1989). Leucine-zipper motif update. *Nature* **340**, 103.
- Miller, H. I.** (1984). Primary structure of the *himA* gene of *Escherichia coli*: homology with DNA binding protein HU and association with the phenylalanyl-tRNA synthetase operon. *Cold Spring Harb. Symp. Quant. Biol.* **49**, 691-698.
- Miyamoto, D. M. and Crowther, R. J.** (1985). Formation of the notochord in living ascidian embryos. *J. Embryol. Exp. Morph.* **86**, 1-17.
- Nash, H. A.** (1990). Bending and supercoiling of DNA at the attachment site of bacteriophage lambda. *Trends Biochem. Sci.* **15**, 222-227.
- Nash, H. A. and Granston, A. E.** (1991). Similarity between the DNA-binding domains of IHF protein and TFIID protein. *Cell* **67**, 1037-1038.
- Nikolov, D. B., Hu, S.-U., Lin, J., Gasch, A., Hoffmann, A., Horikoshi, M., Chua, N.-H., Roeder, R. G. and Burley, S. K.** (1992). Crystal structure of TFIID TATA-box binding protein. *Nature* **360**, 40-46.
- Nishida, H.** (1992). Developmental potential for tissue differentiation of fully dissociated cells of the ascidian embryo. *Roux's Arch. Dev. Biol.* **201**, 81-87.
- Nishida, H. and Satoh, N.** (1989). Determination and regulation in the pigment cell lineage of the ascidian embryo. *Dev. Biol.* **132**, 355-367.
- Raff, R. A.** (1987). Constraint, flexibility, and phylogenetic history in the evolution of direct development in sea urchins. *Dev. Biol.* **119**, 6-19.
- Reddy, B. A., Etkin, L. D. and Freemont, P. S.** (1992). A novel zinc finger coiled coil domain in a family of nuclear proteins. *Trends Biochem. Sci.* **17**, 344-345.
- Sambrook, J., Fritsch, E. F. and Maniatis, T.** (1989). *Molecular Cloning: A Laboratory Manual*. 2nd Ed., New York: Cold Spring Harbor.
- Sanger, F., Nicklen, S. and Coulson, A. R.** (1977). DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**, 5463-5467.
- Satoh, N., Deno, T., Nishida, H., Nishikata, T. and Makabe, K. W.** (1990). Cellular and molecular mechanisms of muscle cell differentiation in ascidian embryos. *Int. Rev. Cytol.* **122**, 221-258.
- Sive, H. L. and St. John, T.** (1988). A simple subtractive hybridization technique employing photoactivatable biotin and phenol extraction. *Nuc. Acids Res.* **16**, 10937.
- Suzuki, M.** (1989). SPXX, a frequent sequence motif in gene regulatory proteins. *J. Mol. Biol.* **207**, 61-84.
- Swalla, B. J.** (1992). The role of maternal factors in ascidian muscle development. *Sem. Dev. Biol.* **3**, 287-295.
- Swalla, B. J. and Jeffery, W. R.** (1990). Interspecific hybridization between an anural and urodele ascidian: Differential expression of urodele features suggests multiple mechanisms control anural development. *Dev. Biol.* **142**, 319-334.
- Swalla, B. J. and Jeffery, W. R.** (1991). Vestigial brain melanocyte development during embryogenesis of an anural ascidian. *Dev. Growth Diff.* **34**, 12-25.
- Swalla, B. J., Badgett, M. R. and Jeffery, W. R.** (1991). Identification of a cytoskeletal protein localized in the myoplasm of ascidian eggs: localization is modified during anural development. *Development* **111**, 425-436.
- Tanaka, I., Appelt, K., Dijk, J., White, S. W. and Wilson, K. S.** (1984). 3-Å resolution structure of a protein with histone-like properties in prokaryotes. *Nature* **310**, 376-381.
- Timblin, C., Battey, J. and Kuehl, W. M.** (1990). Amplification for PCR technology to subtractive cDNA cloning: Identification of genes expressed specifically in murine plasmacytoma cells. *Nuc. Acids Res.* **18**, 1587-1593.
- Vallee, B. L., Coleman, J. E. and Auld, D. S.** (1991). Zinc fingers, zinc clusters and zinc twists in DNA binding proteins. *Proc. Natl. Acad. Sci. USA* **88**, 999-1003.
- Van Gelder, R. N., von Zastrow, M. E., Yool, A., Denent, W. C., Barchas, J. D. and Eberwine, J. H.** (1990). Amplified RNA synthesized from limited quantities of heterogeneous cDNA. *Proc. Nat. Acad. Sci. USA* **87**, 1663-1667.
- Venuti, J. M. and Jeffery, W. R.** (1989). Cell lineage and determination of cell fate in ascidian embryos. *Int. J. Dev. Biol.* **33**, 197-212.
- von Heijne, G.** (1986). A new method for predicting signal sequence cleavage sites. *Nuc. Acids Res.* **14**, 4883-4690.
- White, M. K. and Weber, M. J.** (1989). Leucine-zipper motif update. *Nature* **340**, 103-104.
- Whittaker, J. R.** (1979a). Development of vestigial tail muscle acetylcholinesterase in embryos of an anural ascidian species. *Biol. Bull. Mar. Biol. Lab., Woods Hole* **156**, 393-407.
- Whittaker, J. R.** (1979b). Development of tail muscle acetylcholinesterase in ascidian embryos lacking mitochondrial localization and segregation. *Biol. Bull. Mar. Biol. Lab., Woods Hole* **157**, 344-355.
- Yang, C.-C. and Nash, H. A.** (1989). The interaction of *E. coli* IHF protein with its specific binding sites. *Cell* **57**, 869-880.

(Accepted 30 June 1993)