

Electrosensory ampullary organs are derived from lateral line placodes in cartilaginous fishes

J. Andrew Gillis^{1,2,*}, Melinda S. Modrell¹, R. Glenn Northcutt³, Kenneth C. Catania⁴, Carl A. Luer⁵ and Clare V. H. Baker^{1,‡}

SUMMARY

Ampullary organ electroreceptors excited by weak cathodal electric fields are used for hunting by both cartilaginous and non-teleost bony fishes. Despite similarities of neurophysiology and innervation, their embryonic origins remain controversial: bony fish ampullary organs are derived from lateral line placodes, whereas a neural crest origin has been proposed for cartilaginous fish electroreceptors. This calls into question the homology of electroreceptors and ampullary organs in the two lineages of jawed vertebrates. Here, we test the hypothesis that lateral line placodes form electroreceptors in cartilaginous fishes by undertaking the first long-term *in vivo* fate-mapping study in any cartilaginous fish. Using Dil tracing for up to 70 days in the little skate, *Leucoraja erinacea*, we show that lateral line placodes form both ampullary electroreceptors and mechanosensory neuromasts. These data confirm the homology of electroreceptors and ampullary organs in cartilaginous and non-teleost bony fishes, and indicate that jawed vertebrates primitively possessed a lateral line placode-derived system of electrosensory ampullary organs and mechanosensory neuromasts.

KEY WORDS: Lateral line placode, Electroreceptor, Neuromast hair cell, Parvalbumin-3, *Eya4*, Chondrichthyan

INTRODUCTION

Electrosensory 'hair cells' excited by weak cathodal electric fields (Bodznick and Montgomery, 2005; Münz et al., 1984; Teeter et al., 1980) and innervated by lateral line nerves that project to a dorsal octavolateral nucleus in the medulla (Bullock et al., 1983) are present in both lineages of jawed vertebrates: all cartilaginous fishes (sharks, skates, rays and holocephalans) and some bony fishes (the lobe-finned salamanders, caecilians, coelacanths and lungfishes, and the ray-finned sturgeons, paddlefishes and bichirs) (Bullock et al., 1983; New, 1997; Northcutt, 1997; Schlosser, 2002a). These electroreceptors are housed in ampullary sense organs, i.e. sensory epithelia at the base of conductive jelly-filled canals that open to the surface via pores (Jørgensen, 2005). Their similar electrophysiology and innervation suggest they are homologous, but conflicting data on their embryonic origins call this homology into question. We have demonstrated experimentally, by direct cell lineage tracing in axolotl (Northcutt et al., 1995) and paddlefish (Modrell et al., 2011a), that ampullary organs in bony fishes are ancestrally derived from lateral line placodes, i.e. patches of thickened neurogenic cranial ectoderm that elongate in characteristic lines over the head and trunk, and that also form neuromasts containing mechanosensory hair cells (Aman and Piotrowski, 2011; Baker et al., 2008; Ghysen and Dambly-Chaudière, 2007; Ma and Raible, 2009; Sarrazin et al., 2010;

Schlosser, 2002b; Schlosser, 2006). Analysis of gene and antigen expression in developing shark ampullary organs, however, has led to the suggestion that the electrosensory hair cells of cartilaginous fishes derive from the neural crest (Freitas et al., 2006). We sought to test the hypothesis that lateral line placodes give rise to ampullary organ electroreceptors in cartilaginous fishes by using *in vivo* fate-mapping techniques in the little skate, *Leucoraja erinacea*.

MATERIALS AND METHODS

Embryo collection and Dil-labelling

L. erinacea eggs were obtained from the Marine Biological Laboratory (Woods Hole, MA, USA) and maintained in a flow-through seawater system at ambient temperature to approximately stage 25 (Maxwell et al., 2008). A window was cut in the eggcase, and the embryo and yolk sac removed to a Petri dish. Embryos were anaesthetized in a solution of ethyl 3-aminobenzoate methanesulfonate salt (MS-222, Sigma) in seawater. CellTracker CM-Dil (Invitrogen), diluted in 0.3 M sucrose from a 5 µg/µl stock diluted in ethanol, was focally injected into the anterodorsal, anteroventral or posterior lateral line placode using a pulled glass capillary needle and a Picospritzer pressure injector. Embryos were allowed to recover in seawater, then replaced in their eggcases and left to develop in a flow-through seawater table for 6-10 weeks. *Raja eglanteria* embryos for scanning electron microscopy were obtained from Mote Marine Laboratory (Sarasota, FL, USA).

Histology, immunohistochemistry and *in situ* hybridization

L. erinacea embryos were fixed in 4% paraformaldehyde in phosphate-buffered saline (PBS) overnight at 4°C, rinsed three times in PBS and stored at 4°C in PBS with 0.01% sodium azide. For histological analysis, Dil-labelled embryos were embedded in paraffin wax and sectioned at 10 µm as previously described (O'Neill et al., 2007). Immunohistochemistry with anti-parvalbumin 3 (a gift from A. Hudspeth, The Rockefeller University, NY, USA) was performed in whole mount and on sections as previously described (Modrell et al., 2011a). Nuclei were counterstained with DAPI. *In situ* hybridization for *L. erinacea Eya4* (GenBank Accession Number JQ425114) was performed in whole mount and on sections as previously described (O'Neill et al., 2007), except that slides were not treated with proteinase K before hybridization and the colour reaction was performed using BM Purple (Roche).

¹Department of Physiology, Development and Neuroscience, University of Cambridge, Anatomy Building, Downing Street, Cambridge CB2 3DY, UK. ²Marine Biological Laboratory, 7 MBL Street, Woods Hole, MA 02543, USA. ³Scripps Institution of Oceanography and Department of Neurosciences, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA. ⁴Department of Biological Sciences, Vanderbilt University, 465 21st Ave S, Nashville, TN 37240, USA. ⁵Marine Biomedical Research Program, Mote Marine Laboratory, 1600 Ken Thompson Parkway, Sarasota, FL 34236, USA.

*Present address: Department of Biology, Dalhousie University, 1355 Oxford Street, Halifax, NS B3H 4R2, Canada

‡Author for correspondence (cvhb1@cam.ac.uk)

Scanning electron microscopy

After fixation in 2% glutaraldehyde, *R. eglanteria* embryos were dehydrated into ethanol and processed with dimethoxypropane and CO₂ or Freon 113 (1,1,2 trichlorotrifluoroethane) and Freon 13 (chlorotrifluoromethane). Specimens were mounted, sputter-coated with gold palladium alloy (300-400 Å thick) and imaged.

RESULTS AND DISCUSSION

Parvalbumin 3 is a marker of electrosensory and mechanosensory hair cells in *Leucoraja erinacea*

In order to identify ampullary organs more easily in *L. erinacea* embryos, we examined the expression of parvalbumin 3 (Pv3), a Ca²⁺-binding protein thought to be the major Ca²⁺ buffer in mechanosensory hair cells of the inner ear and lateral line (Heller et al., 2002), and which we recently showed also to be expressed in ampullary electroreceptors in the paddlefish, *Polyodon spathula* (Modrell et al., 2011a), and the axolotl, *Ambystoma mexicanum* (Modrell and Baker, 2012). Whole-mount immunostaining for Pv3 in *L. erinacea* at stage 33 (Maxwell et al., 2008) revealed expression in the entire cephalic network of lateral line organs (Fig. 1A), i.e. both in the lines of lateral line neuromasts in canals (Fig. 1B), and in the adjacent fields of ampullary organs located deep within the dermis (Fig. 1C). Hence, Pv3 is expressed in electrosensory hair cells in ampullary organs, as well as in mechanosensory hair cells, in both cartilaginous fishes and bony fishes (Modrell and Baker, 2012; Modrell et al., 2011a), suggesting that it acts as a Ca²⁺ buffer for both types of sensory hair cell in all jawed vertebrates.

Lateral line placodes in *L. erinacea* give rise to cephalic ampullary organs and neuromasts

We aimed to test the hypothesis that lateral line placodes form both ampullary organs and neuromasts in cartilaginous fishes, by performing *in vivo* fate-mapping in the little skate. We used the fluorescent lipophilic dye DiI to label focally the anterodorsal lateral line placode alone, or the anterodorsal plus anteroventral lateral line placodes, in *L. erinacea* embryos at stage 25 ($n=18$). These placodes were selected because they are the largest cranial lateral line placodes, hence the easiest to target, and they give rise to both ampullary organs and neuromasts in the axolotl (Northcutt et al., 1995) and paddlefish (Modrell et al., 2011a). The anterodorsal and anteroventral lateral line placodes can both be recognized in skate embryos morphologically as thickenings of ectoderm caudal to the eye and dorsal to the mandibular and hyoid arches, respectively (Fig. 1D), and molecularly, by expression of the gene encoding the transcription co-factor *Eya4* (Fig. 1E), an established marker of the developing lateral line system in shark, paddlefish and axolotl embryos (O'Neill et al., 2007; Modrell et al., 2011a; Modrell and Baker, 2012). Fig. 1F shows an example of an embryo photographed immediately after receiving a focal injection of DiI in both the anterodorsal and anteroventral lateral line placodes. In different DiI-injected *L. erinacea* embryos, DiI was incorporated into the elongating primordia of the anterodorsal placode (Northcutt, 2005) and/or anteroventral placodes. For example, in one DiI-labelled embryo in which the anterodorsal lateral line placode was labelled, DiI-positive cells were seen at 14 days post-injection within the migrating supraorbital primordium of the anterodorsal lateral line placode, far from the original injection site (Fig. 1G). At 69 days post-injection (Fig. 1H-J), DiI-positive cells were observed on the ventral surface of the same embryo in patterns that recapitulated the normal distribution of ampullary organs and




Fig. 1. Lateral line placodal origin of skate ampullary organs and neuromasts. (A-C) Whole-mount immunostaining for parvalbumin 3 (Pv3) in *L. erinacea* embryos (A) reveals a cephalic network of mechanosensory neuromasts (B) and electrosensory ampullary organs (C). In order to test the lateral line placodal origin of neuromasts and ampullary organs, we fate mapped the anterodorsal or anteroventral lateral line placodes. (D,E) The anterodorsal and anteroventral placodes were recognizable as (D) ectodermal thickenings caudal to the eye and dorsal to the mandibular and hyoid arches, respectively, which (E) express the transcription co-factor gene *Eya4*. (F) An embryo in which the anterodorsal and anteroventral lateral line placodes were focally labelled with DiI. (G) In an embryo in which the anterodorsal lateral line placode was labelled, DiI was observed at 14 days post-injection in the supraorbital lateral line primordium (black arrows), far from the original injection site (*). (H-M) In embryos with DiI-labelled anterodorsal and/or anteroventral lateral line placodes, the distribution of DiI-positive cells at 60-70 days post-injection recapitulated the normal distribution of cephalic neuromasts and ampullary pores (H-J), and ampullary organs (K-M). ad, anterodorsal lateral line placode; ao, ampullary organ; ap, ampullary tubule pore; av, anteroventral lateral line placode; e, eye; nm, neuromast; ot, otic vesicle; so, supraorbital lateral line primordium. Scale bars: 2.5 mm in A,H,K; 0.5 mm in B-F,J,M; 0.8 mm in G.

neuromasts (Fig. 1I,J). We observed similar distributions of DiI in both neuromasts and ampullary organs (Fig. 1K-M) at 60-70 days post-injection in 17/18 injected embryos.




Fig. 2. Lateral line placodal origin of sensory receptor cells, support cells and canal cells in skate ampullary organs and cephalic neuromasts. (A) A ventral view of the head, illustrating the elongation of the sensory ridge of the anterodorsal lateral line placode that will give rise to the infraorbital neuromasts and closely associated ampullary organs. (B) A lateral view of the head, illustrating the elongation of the sensory ridge of the anterodorsal lateral line placode as it passes rostral and ventral to the eye (where it will run medial to the more laterally situated infraorbital line), giving rise to the supraorbital neuromasts and closely associated ampullary organs. (C) Electrosensory ampullary organs (black arrows) are clustered within the dermis, and each sensory ampulla opens externally via a long jelly-filled ampullary canal (*). (D) Mechanosensory neuromasts are distributed within a continuous network of epithelial canals. (E,F) Immunohistochemical localization of parvalbumin 3 (Pv3) in (E) ampullary organs and (F) neuromasts reveals small clusters of sensory receptor cells nested among non-sensory support cells. (G,H) Electrosensory hair cells (G) and mechanosensory hair cells (H) both express *Eya4*, an established marker of the developing lateral line system. (I-L) Fate mapping of the anterodorsal lateral line placode of *L. erinacea* by Dil injection reveals a placodal origin of (I) the Pv3-positive sensory hair cells, non-sensory support cells and (J) canal cells of electrosensory ampullary organs; and (K) the Pv3-positive sensory hair cells, canal cells and (L) non-sensory support cells of mechanosensory neuromasts. Images in I-L are all from different individuals. ap, ampullary tubule pore; e, eye; io, infraorbital sensory ridge; m, mouth; nm, neuromast; olf, olfactory organ; so, supraorbital sensory ridge. Scale bars: 125 μ m in A; 50 μ m in B; 24 μ m in C,D; 10 μ m in E-L.

The skate anterodorsal and anteroventral lateral line placodes elongate to form sensory ridges that are closely associated with lines of mechanosensory neuromasts and fields of electrosensory ampullary organs. For example, the anterodorsal lateral line placode splits to form the infraorbital (Fig. 2A) and supraorbital (Fig. 2B) sensory ridges, which are closely associated with the ampullary organs and neuromasts located between the rostrum and the mouth (Northcutt, 2005). Differentiated ampullary organs in cartilaginous fishes are organized into fields of sensory ampullae (Fig. 2C) at the base of long, jelly-filled ampullary tubules that open externally (Jørgensen, 2005). By contrast, neuromasts are organized as lines and are regularly spaced within a continuous epithelial canal (Fig. 2D). Both ampullary organs (Fig. 2E) and neuromasts (Fig. 2F) contain Pv3-positive sensory receptor cells nested among non-sensory support cells. As in paddlefish and axolotl embryos (Modrell et al., 2011a; Modrell and Baker, 2012), the Pv3-positive sensory receptor cells of *L. erinacea* ampullary organs (Fig. 2G) and neuromasts (Fig. 2H) express *Eya4*. In order to determine the precise fate of Dil-labelled cells that originated from the lateral line placodes, we immunostained sections of six of our experimental embryos for Pv3. In all six embryos, Dil was observed in Pv3-positive sensory receptor cells (Fig. 2I), non-sensory support cells (Fig. 2I) and canal cells (Fig. 2J) of ampullary organs, and Pv3-positive sensory hair cells (Fig. 2K), non-sensory support cells (Fig. 2L) and canal cells (Fig. 2K) of neuromasts.

The posterior lateral line placode in *L. erinacea* gives rise to trunk neuromasts

Ampullary organs are confined to the head of cartilaginous fishes, but neuromasts are also found on the trunk. Grafting experiments performed a century ago revealed the posterior lateral line (PLL) placode origin of trunk neuromasts in the bony fish clade (Harrison, 1904). The skate PLL placode is easily recognized as a thickening of ectoderm dorsal to the gill arches (Fig. 3A) that elongates caudally along the trunk, forming a distinct ridge in the epidermis (Fig. 3B,C); it also expresses *Eya4* (Fig. 3D). To complete our experimental study of the origin of lateral line organs in cartilaginous fishes, we labelled the PLL placode in *L. erinacea* with Dil (Fig. 3E, $n=5$): 50–65 days post-injection (Fig. 3F), all labelled PLL placodes had deposited clusters of Dil-positive cells along the trunk (Fig. 3G,H) in a pattern recapitulating the distribution of trunk neuromasts (as illustrated by whole-mount Pv3 immunostaining; Fig. 3I). *L. erinacea* trunk neuromasts resemble head neuromasts, with regularly spaced clusters of Pv3-positive (Fig. 3J) and *Eya4*-positive (Fig. 3K) hair cells nested among non-sensory support cells in a continuous epithelial canal. In trunk sections of embryos with Dil-labelled PLL placodes, Dil was observed in both Pv3-positive hair cells and support cells (Fig. 3L). Hence, the PLL placode forms trunk neuromasts in cartilaginous fishes, as it does in bony fishes.




Fig. 3. Lateral line placodal origin of both sensory receptor cells and support cells in skate trunk neuromasts.

(A) The skate posterior lateral line placode is recognizable as an elongated ectodermal thickening dorsal to the pharyngeal arches. (B, C) This placode elongates caudally along the entire length of the trunk (B), forming a distinct ridge in the epidermis (C). (D) The posterior lateral line placode expresses the transcription co-factor gene *Eya4*. (E–I) *L. erinacea* embryos in which the posterior lateral line placode was labelled with Dil (E) show Dil-positive cells (F–H) organized in a pattern that recapitulates the normal distribution of parvalbumin 3 (Pv3)-positive trunk mechanosensory neuromasts (I). (J, K) Immunohistochemical localization of Pv3 in trunk neuromasts reveals small clusters of sensory hair cells nested among non-sensory support cells (J); these sensory hair cells also express *Eya4* (K). (L) The presence of Dil in trunk neuromast Pv3-positive sensory hair cells as well as in non-sensory support cells confirms the posterior lateral line placodal origin of these cell types in *L. erinacea*. m, middle lateral line placode; pcf, pectoral fin; pvf, pelvic fin; st, supratemporal lateral line placode; nm, neuromast; pllp, posterior lateral line placode. Scale bars: 250 μ m in A, C–E; 100 μ m in B; 5 mm in F; 500 μ m in G–I; 10 μ m in J–L.

Overall, our *in vivo* fate-mapping data show that lateral line placodes form both ampullary organs and neuromasts, including the sensory receptor cells within them, in cartilaginous fishes. The proposal that shark electroreceptors are neural crest derived (Freitas et al., 2006) was based on the expression of *Sox8* and HNK1 (neither of which exclusively marks neural crest cells) in developing ampullary organs, and not on lineage-tracing data. Taken together with the previous demonstration of a lateral line placode origin for ampullary organs in both lobe-finned and non-

teleost ray-finned bony fishes (Northcutt et al., 1995; Modrell et al., 2011a), our data confirm that ampullary organs in jawed vertebrates are primitively derived from lateral line placodes.

The lateral line placode-derived ampullary organs of jawed vertebrates probably represent the elaboration of an electrosensory system already present in the last common ancestor of vertebrates. Lampreys (jawless fishes) possess epidermal ‘end bud’ electroreceptors (Bodznick and Northcutt, 1981; Jørgensen, 2005). Although not housed in ampullary organs, lamprey electroreceptors share key features with the ampullary electroreceptors of non-teleost jawed fishes: they respond to cathodal stimuli (Bodznick and Preston, 1983) and are innervated by the anterior lateral line nerve, which projects to an electroreceptive dorsal octavolateral nucleus in the medulla (Bodznick and Northcutt, 1981; Bodznick and Preston, 1983; Ronan and Bodznick, 1986). Taken together, these shared characters suggest homology with non-teleost ampullary organ electroreceptors. However, the embryonic origin of lamprey electroreceptors is currently unknown.

Within jawed vertebrates, electroreception has been lost independently at least three times (in the lineages leading to anuran amphibians, amniotes and neopterygian fishes, i.e. gars, bowfins and teleosts) (Bullock et al., 1983; New, 1997; Northcutt, 1997; Schlosser, 2002a), and has also evolved at least twice independently within the teleosts (in siluriforms and gymnotiforms, and in mormyriiforms) (Alves-Gomes, 2001; Bullock et al., 1983; New, 1997; Northcutt, 1997) and convergently as a specialization of trigeminal nerve endings in monotremes (Pettigrew, 1999) and dolphins (Czech-Damal et al., 2012). Teleost electroreceptors are innervated by lateral line nerves projecting to a special ‘electrosensory lateral line lobe’ in the hindbrain (Bodznick and Montgomery, 2005), but in contrast to non-teleost electroreceptors (Lu and Fishman, 1995), teleost electroreceptors are excited by anodal stimuli, and the basal membrane is the voltage sensor (Bodznick and Montgomery, 2005). As neurotransmitter release is triggered in mechanosensory hair cells by opening voltage-gated channels in the basal membrane, electroreceptors may have evolved in some teleosts via genetic modification of the mechanisms underlying neuromast hair cell differentiation (Bodznick and Montgomery, 2005), but this hypothesis remains untested. Although the molecular mechanisms underlying mechanosensory hair cell formation have been intensively studied (Driver and Kelley, 2009; Puligilla and Kelley, 2009), few papers have reported gene expression during ampullary organ development (Freitas et al., 2006; Metscher et al., 1997; Modrell and Baker, 2012; Modrell et al., 2011a; Modrell et al., 2011b; O’Neill et al., 2007). Discovering the extent to which the mechanisms of mechanosensory and electrosensory receptor cell formation are conserved – both for ancestral ampullary organ electroreceptors and for independently evolved teleost electroreceptors – may shed further light on the fascinating evolutionary history of this ancient vertebrate sense.

Acknowledgements

We thank Mark Terasaki for assistance with skate embryo husbandry, Kim Cooper for assistance with skate embryo storage and shipping, and the staff of the Woods Hole Marine Biological Laboratory for technical support.

Funding

This work was funded by a Royal Society Newton International Fellowship (to J.A.G.), Marine Biological Laboratory Spiegel and Colwin Endowed Summer Research Fellowships (to J.A.G.), the National Science Foundation [BRS 8806539 to R.G.N.] and the Biotechnology and Biological Sciences Research Council [BB/F00818X/1 to C.V.H.B.].

Competing interests statement

The authors declare no competing financial interests.

References

- Alves-Gomes, J. A.** (2001). The evolution of electroreception and bioelectrogenesis in teleost fish: a phylogenetic perspective. *J. Fish Biol.* **58**, 1489-1511.
- Aman, A. and Piotrowski, T.** (2011). Cell-cell signaling interactions coordinate multiple cell behaviors that drive morphogenesis of the lateral line. *Cell Adhes. Migr.* **5**, 499-508.
- Baker, C. V. H., O'Neill, P. and McCole, R. B.** (2008). Lateral line, otic and epibranchial placodes: developmental and evolutionary links? *J. Exp. Zool. B Mol. Dev. Evol.* **310B**, 370-383.
- Bodznick, D. and Northcutt, R. G.** (1981). Electroreception in lampreys: evidence that the earliest vertebrates were electroreceptive. *Science* **212**, 465-467.
- Bodznick, D. and Preston, D. G.** (1983). Physiological characterization of electroreceptors in the lampreys *Ichthyomyzon unicuspis* and *Petromyzon marinus*. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **152**, 209-217.
- Bodznick, D. and Montgomery, J. C.** (2005). The physiology of low-frequency electrosensory systems. In *Electroreception* (ed. T. H. Bullock, C. D. Hopkins, A. N. Popper and R. R. Fay), pp. 132-153. New York, NY: Springer.
- Bullock, T. H., Bodznick, D. A. and Northcutt, R. G.** (1983). The phylogenetic distribution of electroreception: evidence for convergent evolution of a primitive vertebrate sense modality. *Brain Res.* **287**, 25-46.
- Czech-Damal, N. U., Liebschner, A., Miersch, L., Klauer, G., Hanke, F. D., Marshall, C., Dehnhardt, G. and Hanke, W.** (2012). Electroreception in the Guiana dolphin (*Sotalia guianensis*). *Proc. R. Soc. B* **279**, 663-668.
- Driver, E. C. and Kelley, M. W.** (2009). Specification of cell fate in the mammalian cochlea. *Birth Defects Res. C Embryo Today* **87**, 212-221.
- Freitas, R., Zhang, G., Albert, J. S., Evans, D. H. and Cohn, M. J.** (2006). Developmental origin of shark electrosensory organs. *Evol. Dev.* **8**, 74-80.
- Ghysen, A. and Dambly-Chaudière, C.** (2007). The lateral line microcosmos. *Genes Dev.* **21**, 2118-2130.
- Harrison, R. G.** (1904). Experimentelle Untersuchung über die Entwicklung der Sinnesorgane der Seitenlinie bei den Amphibien. *Arch. Mikrosk. Anat.* **63**, 35-149.
- Heller, S., Bell, A. M., Denis, C. S., Choe, Y. and Hudspeth, A. J.** (2002). Parvalbumin 3 is an abundant Ca²⁺ buffer in hair cells. *J. Assoc. Res. Otolaryngol.* **3**, 488-498.
- Jørgensen, J. M.** (2005). Morphology of electroreceptive sensory organs. In *Electroreception* (ed. T. H. Bullock, C. D. Hopkins, A. N. Popper and R. R. Fay), pp. 47-67. New York, NY: Springer.
- Lu, J. and Fishman, H. M.** (1995). Ion channels and transporters in the electroreceptive ampullary epithelium from skates. *Biophys. J.* **69**, 2467-2475.
- Ma, E. Y. and Raible, D. W.** (2009). Signaling pathways regulating zebrafish lateral line development. *Curr. Biol.* **19**, R381-R386.
- Maxwell, E. E., Fröbisch, N. B. and Heppleston, A. C.** (2008). Variability and conservation in late chondrichthyan development: ontogeny of the winter skate (*Leucoraja ocellata*). *Anat. Rec. (Hoboken)* **291**, 1079-1087.
- Metscher, B. D., Northcutt, R. G., Gardiner, D. M. and Bryant, S. V.** (1997). Homeobox genes in axolotl lateral line placodes and neuromasts. *Dev. Genes Evol.* **207**, 287-295.
- Modrell, M. S. and Baker, C. V. H.** (2012). Evolution of electrosensory ampullary organs: conservation of *Eya4* expression during lateral line development in jawed vertebrates. *Evol. Dev.* **14**, 277-285.
- Modrell, M. S., Bemis, W. E., Northcutt, R. G., Davis, M. C. and Baker, C. V. H.** (2011a). Electrosensory ampullary organs are derived from lateral line placodes in bony fishes. *Nat. Commun.* **2**, 496.
- Modrell, M. S., Buckley, D. and Baker, C. V. H.** (2011b). Molecular analysis of neurogenic placode development in a basal ray-finned fish. *Genesis* **49**, 278-294.
- Münz, H., Claas, B. and Fritzsche, B.** (1984). Electroreceptive and mechanoreceptive units in the lateral line of the axolotl *Ambystoma mexicanum*. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **154**, 33-44.
- New, J. G.** (1997). The evolution of vertebrate electrosensory systems. *Brain Behav. Evol.* **50**, 244-252.
- Northcutt, R. G.** (1997). Evolution of gnathostome lateral line ontogenies. *Brain Behav. Evol.* **50**, 25-37.
- Northcutt, R. G.** (2005). Ontogeny of electroreceptors and their neural circuitry. In *Electroreception* (ed. T. H. Bullock, C. D. Hopkins, A. N. Popper and R. R. Fay), pp. 112-131. New York, NY: Springer.
- Northcutt, R. G., Brändle, K. and Fritzsche, B.** (1995). Electroreceptors and mechanosensory lateral line organs arise from single placodes in axolotls. *Dev. Biol.* **168**, 358-373.
- O'Neill, P., McCole, R. B. and Baker, C. V. H.** (2007). A molecular analysis of neurogenic placode and cranial sensory ganglion development in the shark, *Scyliorhinus canicula*. *Dev. Biol.* **304**, 156-181.
- Pettigrew, J. D.** (1999). Electroreception in monotremes. *J. Exp. Biol.* **202**, 1447-1454.
- Puligilla, C. and Kelley, M. W.** (2009). Building the world's best hearing aid; regulation of cell fate in the cochlea. *Curr. Opin. Genet. Dev.* **19**, 368-373.
- Ronan, M. C. and Bodznick, D.** (1986). End buds: non-ampullary electroreceptors in adult lampreys. *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **158**, 9-15.
- Sarrazin, A. F., Nuñez, V. A., Sapède, D., Tassin, V., Dambly-Chaudière, C. and Ghysen, A.** (2010). Origin and early development of the posterior lateral line system of zebrafish. *J. Neurosci.* **30**, 8234-8244.
- Schlosser, G.** (2002a). Development and evolution of lateral line placodes in amphibians.-II. Evolutionary diversification. *Zoology (Jena)* **105**, 177-193.
- Schlosser, G.** (2002b). Development and evolution of lateral line placodes in amphibians I. Development. *Zoology (Jena)* **105**, 119-146.
- Schlosser, G.** (2006). Induction and specification of cranial placodes. *Dev. Biol.* **294**, 303-351.
- Teeter, J. H., Szamier, R. B. and Bennett, M. V. L.** (1980). Ampullary electroreceptors in the sturgeon *Scaphirhynchus platyrhynchus* (Rafinesque). *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **138**, 213-223.