

CORRECTION

A transport and retention mechanism for the sustained distal localization of Spn-F-IKK ϵ during *Drosophila* bristle elongation

Tetsuhisa Otani, Kenzi Oshima, Akiyo Kimpara, Michiko Takeda, Uri Abdu and Shigeo Hayashi

There was an error published in *Development* 142, 2338-2351.

Otani et al. reported the genetic interactions between *ikke* and *spn-F*, using the allele *ikke*⁶⁶. This allele was referred to in the Materials and Methods on p. 2349, Fig. 3 on p. 2343 and Table S1. However, they subsequently found that the allele used in the experiments was *ikke*¹ (also known as *ikke*³⁰). This was as a result of misannotation in their laboratory stock list. Both alleles are strong loss-of-function alleles with a missense mutation in the kinase domain and show similar phenotypes (Oshima et al., 2006; Shapiro and Anderson, 2006). Therefore, this error does not affect the conclusions of the paper.

The authors apologise to readers for this mistake.

References

- Oshima, K., Takeda, M., Kuranaga, E., Ueda, R., Aigaki, T., Miura, M. and Hayashi, S. (2006). IKK epsilon regulates F actin assembly and interacts with Drosophila IAP1 in cellular morphogenesis. *Curr. Biol.* 16, 1531-1537.
Shapiro, R. S. and Anderson, K. V. (2006). Drosophila Ik2, a member of the I κ B kinase family, is required for mRNA localization during oogenesis. *Development* 133, 1467-1475.