



Cover: 3D reconstructions of mid-gastrula (left) and mid-neurula (right) *Ciona* embryos electroporated with transgenes driving H2B:Cherry (red) in the epidermis and YFP-CAAX (green) in the neural plate. Nodal and FGF control the cellular behaviours underlying morphogenesis of the neural tube during *Ciona* development. **See Research article by Navarrete and Levine on p. 4665.**

SPOTLIGHT

- 4555 An interview with Doug Melton
Maartens, A.

PRIMER

- 4558 Fox transcription factors: from development to disease
Golson, M. L. and Kaestner, K. H.

REVIEW

- 4571 The many faces of hematopoietic stem cell heterogeneity
Crisan, M. and Dzierzak, E.

STEM CELLS AND REGENERATION

- 4582 Early bioelectric activities mediate redox-modulated regeneration
Ferreira, F., Luxardi, G., Reid, B. and Zhao, M.
- 4595 *Setd5* is essential for mammalian development and the co-transcriptional regulation of histone acetylation
Osipovich, A. B., Gangula, R., Vianna, P. G. and Magnuson, M. A.
- 4608 PRMT5 is essential for the maintenance of chondrogenic progenitor cells in the limb bud
Norrie, J. L., Li, Q., Co, S., Huang, B.-L., Ding, D., Uy, J. C., Ji, Z., Mackem, S., Bedford, M. T., Galli, A., Ji, H. and Vokes, S. A.
- 4620 Transcriptional regulation of intermediate progenitor cell generation during hippocampal development
Harris, L., Zalucki, O., Gobius, I., McDonald, H., Osinki, J., Harvey, T. J., Essebier, A., Vidovic, D., Gladwyn-Ng, I., Burne, T. H., Heng, J. I., Richards, L. J., Gronostajski, R. M. and Piper, M.
- 4631 Phosphorylated Groucho delays differentiation in the follicle stem cell lineage by providing a molecular memory of EGFR signaling in the niche
Johnston, M. J., Bar-Cohen, S., Paroush, Z. and Nystul, T. G.
- 4643 Dynein light chain DLC-1 promotes localization and function of the PUF protein FBF-2 in germline progenitor cells
Wang, X., Olson, J. R., Rasoloson, D., Ellenbecker, M., Bailey, J. and Voronina, E.

RESEARCH ARTICLES

- 4654 Foxn4 promotes gene expression required for the formation of multiple motile cilia
Campbell, E. P., Quigley, I. K. and Kintner, C.

- 4665 Nodal and FGF coordinate ascidian neural tube morphogenesis
Navarrete, I. A. and Levine, M.

- 4676 Integration of actomyosin contractility with cell-cell adhesion during dorsal closure
Duque, J. and Gorfinkiel, N.

- 4687 Phospholipid composition and a polybasic motif determine D6 PROTEIN KINASE polar association with the plasma membrane and tropic responses
Barbosa, I. C. R., Shikata, H., Zourelidou, M., Heilmann, M., Heilmann, I. and Schwechheimer, C.

- 4701 Delayed neurogenesis with respect to eye growth shapes the pigeon retina for high visual acuity
Rodrigues, T., Krawczyk, M., Skowronska-Krawczyk, D., Matter-Sadzinski, L. and Matter, J.-M.

- 4713 Exon- and contraction-dependent functions of titin in sarcomere assembly
Shih, Y.-H., Dvornikov, A. V., Zhu, P., Ma, X., Kim, M., Ding, Y. and Xu, X.

- 4723 Molecular dynamics of Dkk4 modulates Wnt action and regulates meibomian gland development
Sima, J., Piao, Y., Chen, Y. and Schlessinger, D.

- 4736 CFAP157 is a murine downstream effector of FOXJ1 that is specifically required for flagellum morphogenesis and sperm motility
Weidemann, M., Schuster-Gossler, K., Stauber, M., Wrede, C., Hegermann, J., Ott, T., Boldt, K., Beyer, T., Serth, K., Kremmer, E., Blum, M., Ueffing, M. and Gossler, A.

TECHNIQUES AND RESOURCES

- 4749 Insights from imaging the implanting embryo and the uterine environment in three dimensions
Arora, R., Fries, A., Oelerich, K., Marchuk, K., Sabeur, K., Giudice, L. C. and Laird, D. J.

CORRECTION

- 4755 Correction: Mice humanised for the EGF receptor display hypomorphic phenotypes in skin, bone and heart
Sibilia, M., Wagner, B., Hoebertz, A., Elliott, C., Marino, S., Jochum, W. and Wagner, E. F.