

DEVELOPMENT AT A GLANCE

Wnt signaling in development and tissue homeostasis

Zachary Steinhart and Stephane Angers*

ABSTRACT

The Wnt-β-catenin signaling pathway is an evolutionarily conserved cell-cell communication system that is important for stem cell renewal, cell proliferation and cell differentiation both during embryogenesis and during adult tissue homeostasis. Genetic or epigenetic events leading to hypo- or hyper-activation of the Wnt-β-catenin signaling cascade have also been associated with human diseases such as cancer. Understanding how this pathway functions is thus integral for developing therapies to treat diseases or for regenerative medicine approaches. Here, and in the accompanying poster, we provide an overview of Wnt-β-catenin signaling and briefly highlight its key functions during development and adult tissue homeostasis.

KEY WORDS: Wnt protein, Wnt signaling, Epigenetics

Introduction

Wnt proteins are secreted growth factors that regulate the proliferation and differentiation of stem and progenitor cells, both during embryonic development and during adult tissue homeostasis in multicellular animals (Logan and Nusse, 2004). The *Drosophila* Wnt protein Wingless (Wg) and the core members of the intracellular signaling pathway it regulates were originally identified and characterized in flies through genetic screens and phenotypic analyses of embryonic morphogenesis defects (Nüsslein-Volhard and Wieschaus, 1980; 'Sharma and Chopra, 1976). The discovery that Wg was homologous to the mouse *Wnt1* proto-oncogene (Rijsewijk et al., 1987) supported the notion that Wnt1 and possibly other vertebrate Wnt homologs could have important roles during normal vertebrate development, and that Wnt signaling activity might be dysregulated in cancer. A vast number of integrative genetic, biochemical and 'omic-based approaches have since refined our understanding of how Wnt ligands are produced and secreted, how cells recognize and integrate Wnt signals to yield various cellular and physiological responses, and how Wnt signals are modulated, terminated or

University of Toronto, 144 College Street, Toronto, ON M5S 3M2, Canada.

*Author for correspondence (stephane.angers@utoronto.ca)

ID S.A., 0000-0001-7241-9044

amplified depending on spatiotemporal contexts. This work has provided a foundation to recognize and study the pervasive roles of Wnt signaling during development and tissue homeostasis, as well as to understand the frequent defects associated with this pathway in human diseases such as cancer (Nusse and Clevers, 2017).

Multiple functionally divergent Wnt-based signaling pathways have been identified to date. The best characterized of these is the Wnt- β -catenin pathway, which is often referred to as the ‘canonical’ Wnt pathway. This pathway culminates in the regulation of context-specific β -catenin-dependent gene expression programs that direct stem and progenitor cell renewal, proliferation and differentiation. By contrast, the non-canonical pathways operate independently of β -catenin and have been described to impinge on small GTPases of the Rho family to control cell polarity and cell movement, or to act via heterotrimeric G proteins to control Ca^{2+} signaling (Angers and Moon, 2009). Here, and in the accompanying poster, we discuss the key features of the canonical Wnt- β -catenin pathway and briefly highlight its roles in development, stem cells and adult tissue homeostasis.

Wnt signal recognition and pathway initiation

The Wnt- β -catenin signaling pathway centers around the post-translational control of β -catenin protein abundance. In the absence of Wnt proteins, cytoplasmic levels of β -catenin are kept low through ubiquitin-dependent proteasomal degradation, a process governed by a molecular machine called the β -catenin destruction complex (Stamos and Weis, 2012). The destruction complex is composed of the scaffolding proteins Axin (Behrens, 1998; Hart et al., 1998) and APC (Munemitsu et al., 1995; Rubinfeld et al., 1993), and the kinases $\text{CK1}\alpha$ (Amit et al., 2002; Liu et al., 2002; Rubinfeld et al., 1996) and GSK3 α/β (Amit et al., 2002; Liu et al., 2002; Rubinfeld et al., 1996). The destruction complex functions by catalyzing the serine/threonine phosphorylation of a highly conserved phospho-degron at the N terminus of β -catenin (Winston et al., 1999), which earmarks β -catenin for recruitment to the $\text{SCF}^{\beta\text{-TRCP}}$ E3-ubiquitin ligase (Hart et al., 1999) and ensuing proteasome-mediated degradation (Hart et al., 1999). Simply put, in the absence of Wnt proteins, neo-synthesized β -catenin is constitutively targeted for proteolysis, and Wnts function by inhibiting this degradation.

Wnts are proteins of ~40 kDa that are post-translationally modified through palmitoylation and glycosylation. The human genome encodes 19 Wnt proteins (Nusse, 2001), all of which are secreted through a conserved secretory pathway. Wnt proteins are translated in the endoplasmic reticulum (ER), where they are palmitoylated by the ER-bound o-acyl-transferase porcupine (van den Heuvel et al., 1993; Willert et al., 2003). This is an essential step in Wnt protein secretion (Herr and Basler, 2012) and introduces a modification that is required for Wnt activity (Janda et al., 2012; Proffitt and Virshup, 2012; Willert et al., 2003). Following their palmitoylation Wnt proteins interact with the multipass transmembrane escort protein GPR177, the vertebrate homolog of *Drosophila* Wingless/Evi/Sprinter, which mediates endosome trafficking to the plasma membrane and the release of Wnt proteins into the extracellular space (Bänziger et al., 2006; Bartscherer et al., 2006; Goodman et al., 2006).

The cell-surface receptor for Wnts that is responsible for transducing signals through the β -catenin pathway consists of a heterodimer between a Frizzled family member (Bhanot et al., 1996; MacDonald and He, 2012; Yang-Snyder et al., 1996) and one of the Wnt co-receptors LRP5 or LRP6 (LRP5/6) (Pinson et al., 2000; Tamai et al., 2000; Wehrli et al., 2000). Frizzled

proteins are seven-transmembrane receptors, of which there are 10 homologs in humans, which all contain a conserved 120 amino acid extracellular cysteine-rich-domain (CRD) serving as the obligate binding interface for Wnt (Janda et al., 2012; Povelones and Nusse, 2005). Structural studies have revealed that Wnt proteins resemble the general structure of a hand, with the thumb and index fingers pinching the extracellular CRD of Frizzled. The Wnt palmitoleic acid moiety acts as a major binding determinant, inserting itself in a hydrophobic crevice within the CRD (Janda et al., 2012). LRP5/6 Wnt co-receptors, which are single transmembrane domain proteins, contain four extracellular β -propeller domains that are involved in Wnt binding (Chen et al., 2011) and five PPPSP phosphorylation motifs within the intracellular region (Tamai et al., 2004).

The binding of a Wnt ligand to the FZD-LRP5/6 complex first results in recruitment of the cytoplasmic protein Dishevelled (Klingensmith et al., 1994; Krasnow et al., 1995), putatively via direct interaction with Frizzled (Cong et al., 2004; Tauriello et al., 2012), and of Axin-GSK3 complexes via the interaction of Axin with Dishevelled. GSK3 (Tamai et al., 2004) and the Cdk14(PFTK1)-Cyclin Y mitotic kinase complex (Davidson et al., 2009) then phosphorylate PPPSP motifs within the intracellular region of LRP5/6, thereby priming them for further phosphorylation by the membrane-anchored kinase casein kinase I γ (Davidson et al., 2005; Zeng et al., 2005). The phosphorylated C-terminal tail of LRP5/6 was thought to represent a high-affinity binding site for Axin, thereby leading to the titration of active destruction complex molecules and to β -catenin stabilization. However, this model mostly relied on yeast two-hybrid experiments that identified Axin clones using the LRP5 tail as a bait (Mao et al., 2001); subsequent biochemical (Cselenyi et al., 2008; Piao et al., 2008) and structural experiments (Stamos et al., 2014) have demonstrated that phospho-LRP5/6 blocks β -catenin degradation directly by inhibiting GSK3 activity. Whether Axin interacts with the phosphorylated LRP5/6 intracellular tail indirectly through GSK3, or whether direct binding of Axin is needed, remains to be determined.

The subsequent accumulation of β -catenin leads to its translocation to the nucleus, where it acts as a transcriptional co-activator by binding to and modulating the activity of the Tcf/Lef family of DNA-binding proteins (Behrens et al., 1996; Molenaar et al., 1996; van de Wetering et al., 1997) that otherwise repress Wnt-responsive enhancers through their physical interaction with Groucho/Tle family of co-repressors (Cavallo et al., 1998; Roose et al., 1998). How this transcriptional switch functions at the molecular level was recently clarified with the identification of the E3 ubiquitin ligase UBR5 as a required component of the Wnt-dependent response (Flack et al., 2017). UBR5 functions through the Wnt- and ubiquitin-mediated inactivation of Groucho/Tle, thereby leading to de-repression and activation of β -catenin target genes in a context-dependent manner. Wnt- β -catenin target genes vary depending on cell lineage/type (Nakamura et al., 2016), but common targets include genes that function in positive- and negative-feedback regulation of the pathway, genes involved in cell-cycle progression and genes involved in stem cell function.

Wnt signal restriction and termination

Multiple mechanisms exist to restrict and/or terminate Wnt signaling at the ligand, receptor, intracellular and nuclear levels. At the ligand level, the extracellular enzymes Tiki and Notum function as a protease and a carboxylesterase to cleave the N terminus of Wnts (Zhang et al., 2012) or remove their palmitoleate

moiety (Kakugawa et al., 2015; Zhang et al., 2015), respectively. Additionally, Wnt ligands are inhibited through direct interaction with secreted proteins that prevent their interaction with the receptor complex, such as Wnt inhibitory factor (WIF) (Hsieh et al., 1999), or secreted frizzled related proteins (sFRPs) (Hsieh et al., 1999; Leyns et al., 1997; Rattner et al., 1997; Wang et al., 1997), which share homology with the Frizzled CRD.

Several negative-feedback loops exist to terminate or dampen Wnt signals at the receptor level. Dickkopf-1 (*Dkk1*) (Glinka et al., 1998) is a Wnt- β -catenin target gene (González-Sancho et al., 2005) and functions by binding to LRP5/6 in competition with Wnts (Bafico et al., 2001; Semenov et al., 2001). *Rnf43* and *Znrf3* are recently characterized Wnt- β -catenin target genes that encode transmembrane E3-ubiquitin-ligases (Hao et al., 2012; Koo et al., 2012) involved in ubiquitin-dependent Frizzled endocytosis and proteasomal degradation, thereby reducing receptor levels at the plasma membrane.

In the cytoplasm, the best-studied negative feedback for β -catenin levels involves Axin2. Both Axin1 and Axin2 are functionally equivalent proteins (Chia and Costantini, 2005) but, in contrast to its paralog *Axin1*, which is expressed constitutively, *Axin2* is a Wnt target gene (Jho et al., 2002; Lustig et al., 2002). Given that Axin proteins are thought to be limiting components of the β -catenin destruction complex (Lee et al., 2003), the rise in Axin levels in response to Wnt signaling leads to increased destruction complex formation and β -catenin degradation, thereby providing a mechanism for negative feedback (Goentoro and Kirschner, 2009). The Wnt target gene *Nkd1* also provides negative feedback in the cytoplasm but the exact mechanism by which it functions remains unclear. NKD1 was first suspected to bind to and inhibit Dishevelled (Rousset et al., 2001); however, other findings have shown that it interacts directly with β -catenin and prevents its nuclear import (Van Raay et al., 2011). Finally, the 81 amino acid protein inhibitor of β -catenin and T cell factor (ICAT) was found to directly interact with and inhibit β -catenin in the nucleus, acting to inhibit the posteriorizing activity of Wnt- β -catenin signaling in a spatiotemporal manner in the nervous system (Satoh et al., 2004).

Wnt signal amplification

A number of recent studies have revealed a mechanism involving the cell surface protein LGR5 and secreted R-spondin proteins that provides a way of augmenting signaling through the Wnt pathway. *Lgr5*, which is among the best-characterized target genes of the Wnt- β -catenin signaling pathway, is an established stem cell marker (Barker et al., 2007). As such, *Lgr5* reporter mice have proven valuable for the identification, isolation and functional characterization of tissue stem cells (Barker et al., 2007). However, the precise role of LGR5 and its homologs LGR4/LGR6 remained controversial until the discovery that they function as receptors for four R-spondin secreted proteins, RSPO1-RSPO4 (Carmon et al., 2011; de Lau et al., 2011; Glinka et al., 2011). It was further revealed that R-spondins do not possess signaling activity on their own (Janda et al., 2017) but rather amplify Wnt signaling activity, as they require the presence of Wnt ligands. A missing link in our understanding of RSPO-LGR4/5/6 function came with the identification that the E3 ubiquitin ligases RNF43/ZNRF3 act as co-receptors for RSPOs (Chen et al., 2013; Hao et al., 2012; Zebisch et al., 2013). As mentioned above, *Rnf43* and *Znrf3* are themselves target genes of Wnt- β -catenin signaling, and when upregulated are part of a negative-feedback loop that downregulates cell-surface levels of Frizzled-LRP5/6 complexes. Thus, the binding of

R-spondin to the LGR4/5/6-RNF43/ZNRF3 complex leads to endocytosis of the complex and therefore blocks the ability of RNF43 and ZNRF3 to downregulate Frizzled at the cell surface, hence leading to a sensitized Wnt signaling state (Hao et al., 2012). Interestingly, R-spondin may also have roles that are independent of LGR proteins as RSPO2 and RSPO3 were found to have remaining activities in LGR4/5/6 triple-knockout cells (Lebensohn and Rohatgi, 2018). In this case, R-spondin activity is instead mediated by ZNRF3/RNF43 and heparan sulfate proteoglycans.

Interestingly, a recent study revealed that Wnt and R-spondin have non-overlapping roles during stem cell homeostasis (Yan et al., 2017). This study indicated that, whereas R-spondin overexpression is sufficient to expand the *Lgr5*⁺ stem cell population in the intestine, increasing Wnt ligand levels is ineffective. This highlights a unique role for R-spondin in providing the high levels of Wnt signaling activity required to drive stem cell proliferation and self-renewal. These results led the authors to propose a model in which Wnt ligands act as priming factors, enabling expression of the R-spondin receptors LGR5 and RNF43/ZNRF3. Within this model, primed cells exposed to R-spondin would be competent to produce the high Wnt- β -catenin signaling response and activate transcriptional programs required for cell proliferation and self-renewal. On the other hand, cells further away from the source of R-spondin (the stem cell niche) have higher levels of functional RNF43/ZNRF3 at their surface and, as a result, gradually express decreasing amounts of Frizzled receptors, which is translated into decreased levels of Wnt- β -catenin signaling.

Wnt signaling in early vertebrate embryonic development

Cell-cell communication is integral during the embryonic development of multicellular organisms, acting to coordinate stem cell self-renewal, cell fate decisions, cell migration and the organization of cells into tissues. The Wnt- β -catenin signaling pathway is one of the evolutionary conserved communication systems that regulates embryonic development. It involves Wnt ligands that are released by Wnt-producing cells and that act over various ranges to influence neighboring Wnt-responsive cells.

Wnt- β -catenin signaling is instrumental for defining the dorsoventral and anteroposterior body axes in multiple animal species (Genikhovich and Technau, 2017; Niehrs, 2010). For example, following *Xenopus* egg fertilization, cortical rotation leads to dorsal translocation of maternal determinants that are important for β -catenin signaling activation, which in turn leads to specification of the Spemann organizer (Moon and Kimelman, 1998) – an essential embryonic inducer of dorsoventral patterning. Accordingly, the injection of Wnt mRNA or other factors that activate β -catenin signaling into future ventral blastomeres notoriously leads to duplication of the dorsal axis and results in two-headed frog embryos (McMahon and Moon, 1989). Whether Wnt ligands themselves were involved in dorsal axis induction in *Xenopus* remained controversial for a long time, but Wnt-11 was eventually identified as the ligand implicated in this context (Tao et al., 2005). At the onset of gastrulation, Wnt ligand-dependent β -catenin signaling then activates a largely distinct transcriptional program that directs anteroposterior axis development. In this context, Wnt- β -catenin signaling is inhibited anteriorly, leading to the development of head structures, and activated posteriorly to define tail formation (Green et al., 2015). Paramount to these patterning events, the spatiotemporal regulation of Wnt- β -catenin transcriptional programs is achieved through the action of secreted antagonists, epigenetic regulation, the actions of different Tcf/Lef factors and signaling integration with other

developmental pathways. Beyond these early roles in establishing the embryonic body axes, Wnt- β -catenin signaling plays important roles during the morphogenesis of multiple tissues derived from the three germ layers.

Wnt signaling in embryonic stem cells

The derivation of human embryonic stem cells (ESCs) and induced pluripotent stem cells (iPSCs) has provided a system for studying human embryonic development and human diseases, and a framework for the production of large quantities of differentiated cells that can be transplanted for regenerative medicine or tissue engineering applications. The directed differentiation of such pluripotent stem cells (PSCs) into specialized cells in culture follows the same principles underlying embryonic development; as such, insights gained from developmental biology have guided strategies for using growth factors and inhibitors of developmental signaling pathways to maintain pluripotency or steer cell fate towards the desired lineage with increasing efficiencies (Williams et al., 2012).

Functional redundancy between Wnt- β -catenin signaling activation, leukemia inhibitory factor (LIF) signaling activation and MAPK pathway inhibition is important for maintaining the naive pluripotent state in ESCs. Indeed, β -catenin KO ESCs maintain their pluripotency when cultured with LIF but rapidly exit the naive state in its absence (Lyashenko et al., 2011). Reciprocally, the use of GSK3 inhibitors to stimulate β -catenin signaling bypasses the requirement for LIF in this process (Sato et al., 2004). Wnt signaling is therefore not absolutely required for the maintenance of pluripotency, a finding supported by the analysis of porcine KO mESCs that exhibit normal self-renewal properties (Biechele et al., 2013). Paradoxical reports implicating Wnt- β -catenin signaling in the self-renewal versus differentiation of ESCs have been reconciled with its seemingly opposite roles in promoting pluripotency in the naive state (Sato et al., 2004; ten Berge et al., 2011; Xu et al., 2016) and differentiation in the primed state (Davidson et al., 2012; Kurek et al., 2015), which is characterized by a distinct epigenetic and gene expression landscape (Nichols and Smith, 2009). Levels of Wnt- β -catenin signaling therefore appear to control the transition between the naive and primed ESC states. In line with its role in embryonic development, the further differentiation of ESCs into different germ layers requires the temporal activation of Wnt- β -catenin signaling to steer cells towards the mesendoderm lineage or inactivation to obtain neuroectoderm (Tabar and Studer, 2014).

Wnt signaling in adult tissue homeostasis

The self-renewal versus differentiation of stem cells is directed by extrinsic short-ranged signals that emanate from the niche. Confirming the pervasive role of Wnt ligands as niche factors, lineage-tracing experiments using Wnt-target gene reporters have revealed labeled stem cells in several tissues, including the intestine (Barker et al., 2007), the stomach (Barker et al., 2010), the skin (Lim et al., 2013), the liver (Wang et al., 2015) and the mammary gland (van Amerongen et al., 2012). Below, we describe three of the best-studied tissues in which Wnt signaling has been demonstrated to play crucial roles with regard to the function of tissue stem cells.

Intestine

The best-characterized function of Wnt signaling in adult tissues is in the maintenance of stem cell niches, where Wnt ligands promote the proliferation and self-renewal capability of tissue-specific stem cells (Clevers et al., 2014). This has been particularly well studied in

the adult intestinal epithelium, which turns over fully every 4-5 days and thereby requires balanced stem and progenitor cell self-renewal, proliferation and differentiation (Clevers, 2013). The intestinal epithelium is composed of two compartments: the villi, which are responsible for the absorptive and secretion functions of the intestine; and the crypts of Lieberkühn, which are not exposed to the intestinal lumen, and contain tissue stem cells and rapidly proliferating transit amplifying cells. Genetic studies have shown that the disruption of *Tcf4* (Korinek et al., 1998) or β -catenin (Fevr et al., 2007), or the overexpression of *Dkk1* (Fevr et al., 2007; Pinto et al., 2003), leads to robust loss of transit amplifying cells and crypt structures, confirming the functional requirement of Wnt- β -catenin signaling in the intestinal epithelium. These genetic experiments are corroborated by the fact that *APC* loss-of-function mutations induce adenoma formation (Morin, 1997) and occur in 80% of colorectal cancers (Cancer Genome Atlas Network, 2012). Studies on the expression of Wnt target genes have identified a Wnt signaling gradient, with highest expression in the base of the crypt, particularly in the crypt base columnar cells (CBCs) (Kosinski et al., 2007; Van der Flier et al., 2007). Wnts are bound to the plasma membrane of cells at the base of the crypt and are diluted with each cell division away from the base, thereby establishing a gradient (Farin et al., 2016). Underlying the CBCs at the bottom of crypts are myoepithelial cells, which have been shown to be the *in vivo* source of Wnt ligands (Kabiri et al., 2014). In addition, the Wnt target gene *Lgr5* was found to be expressed exclusively in CBCs meaning that it could be used for lineage-tracing experiments; these experiments showed that *Lgr5*⁺ CBCs could differentiate into all epithelial cell types (Barker et al., 2007). Further supporting the possibility that *Lgr5*⁺ CBCs are the intestinal tissue stem cells, dissociated single *Lgr5*⁺ cells were found to give rise to ‘mini-gut’ organoids *in vitro*, which contain all intestinal cell types (Sato et al., 2009).

Bone

Bone tissue remodeling is crucial for maintaining a balance between systemic calcium homeostasis and the biomechanical needs of the skeleton. Human genetics data perhaps best highlight the crucial role of Wnt- β -catenin signaling in bone tissue homeostasis. Indeed, rare pathological mutations within genes encoding components of the Wnt- β -catenin pathway lead to either severely increased or decreased bone mass. For example, *Lrp5*, which encodes a Wnt co-receptor, was revealed as a critical gene directing bone mass regulation when *LRP5* loss-of-function mutations were identified in the low bone mass disorder osteoporosis-pseudoglioma syndrome (Gong et al., 2001) and when heterozygous missense mutations in *LRP5* were observed in individuals with dominantly inherited high bone mass (Boyden et al., 2002; Little et al., 2002). Another example is mutations discovered in *SOST* (sclerostin), a negative regulator of LRP5/6 that was found to be causally linked to high bone density pathology via loss-of-function mutations (Brunkow et al., 2001) or mutations that decrease its expression (Balemans et al., 2002; Loots et al., 2005). Corroborating this human genetic evidence are extensive genetic experiments performed in mice that link Wnt- β -catenin signaling to the regulation of bone homeostasis (Baron and Kneissel, 2013). Leveraging these findings, several pharmaceutical companies have developed anti-SOST antibodies to activate endogenous Wnt- β -catenin circuitries in bone tissues, and these have been shown in clinical trials to augment bone mass density and prevent fractures (Glorieux et al., 2017; Recker et al., 2015; Saag et al., 2017; Williams, 2017). At the cellular level, the process of bone remodeling results from the concerted actions of three different cell types: osteoblasts, which are responsible for

bone matrix formation; osteoclasts, which are responsible for bone resorption; and osteocytes, which are responsible for bone maintenance. The precise roles of Wnt- β -catenin signaling in bone mass homeostasis still remain to be clarified but the prevailing model is that it functions to regulate osteoblast differentiation and function (Bennett et al., 2005; Cui et al., 2011; Day et al., 2005; Hill et al., 2005; Tu et al., 2015).

Skin

The best-studied role of Wnt signaling in adult skin is its role in the hair follicle cycle, where Wnt is an important factor in orchestrating the control of hair follicle stem cells (HFSCs) (Veltri et al., 2018). The adult hair follicle goes through continuous cycles of telogen (rest), anagen (growth) and catagen (degeneration). The essential role of Wnt- β -catenin in hair follicle cycling was identified through the observations that loss of epidermal β -catenin or inhibition of LRP5/6 via ectopic DKK1 expression leads to defects in hair follicle development and/or loss after a single hair cycle (Andl et al., 2002; Huelsken et al., 2001). Specifically, Wnt- β -catenin signaling acts on multiple cell types to induce the telogen-anagen transition (Greco et al., 2009; Kishimoto et al., 2000; Van Mater et al., 2003). During telogen, HFSCs are slow cycling or quiescent, reside in the bulge region of the hair follicle (Cotsarelis et al., 1990; Lyle et al., 1998), are LGR5-positive (Jaks et al., 2008) and rely on autocrine Wnt for maintenance (Lim et al., 2016). During the telogen-anagen transition, Wnt- β -catenin activity in the bulge HFSCs increases (Lien et al., 2014) and cells from the bulge migrate to the base of the hair follicle and begin proliferating rapidly to produce cell lineages of a new hair follicle (Oshima et al., 2001; Tumber et al., 2004). This proliferation is aided by concurrent Wnt- β -catenin activation in the dermal papilla, which induces growth-promoting FGF signals (Enshell-Seijffers et al., 2010).

Perspectives

Wnt ligands are crucial for the development and homeostasis of virtually all tissues. In the past few decades, genetic, biochemical and structural experiments have identified many Wnt signaling components and have revealed their underlying mechanisms. However, important gaps remain, specifically with respect to the mechanisms of signal transduction regulated by Wnt proteins and β -catenin. For example, although structures of Wnt in complex with the Frizzled CRD have provided details about the mode of binding, it remains unclear how this is coupled to potential allosteric changes in the heptahelical bundles of Frizzled to transduce signaling to the intracellular components. How Wnts co-engage Frizzled and LRP5/6 co-receptors, and the precise nature of the activated complex, also remains poorly characterized. Another loosely understood step is the recruitment of Dishevelled proteins by activated Frizzled receptors, and how this leads to inactivation of destruction complex activity. Furthermore, despite close investigation, the biochemical nature of the destruction complex and the precise molecular mechanisms underlying its function and regulation remain ill defined. The role of β -catenin as a cell-adhesion component when bound to cadherin molecules and whether this pool of β -catenin at the plasma membrane interacts functionally with the transcriptional pool in specific contexts also needs to be evaluated. Similarly, β -catenin transcriptional functions, which are independent of Wnt ligands and influenced by signaling crosstalk with other signaling pathways, need to be considered. In addition, how various levels of Wnt signals are ultimately translated into precise changes in nuclear β -catenin, and hence spatiotemporal modulation of gene expression, is incompletely

explained by our current linear view of the Wnt- β -catenin pathway. More work is thus needed to reverse engineer the complex signaling network formed by the 19 Wnt ligands, 10 Frizzled receptors and multiple co-receptors to better understand how, when and where each of these components functions. Finally, elucidating which Wnt-receptor pairs are involved in specific physiological functions will likely provide more selective therapeutic strategies for treating diseases. Advances in cryoelectron microscopy and single-molecule imaging, combined with progress in functional genomics-based approaches, will undoubtedly help us to fill these gaps.

Acknowledgements

We thank members of our laboratory for helpful discussions and Drs Gregory M. Kelly, Daniel Schramek and Bradley Doble for their thoughts and comments to improve the manuscript.

Competing interests

The authors declare no competing or financial interests.

Funding

The authors' research is supported by grants from the Canadian Institute of Health Research (364969) and the Canadian Cancer Society (705045).

Development at a Glance

A high-resolution version of the poster is available for downloading in the online version of this article at <http://dev.biologists.org/content/145/11/dev146589/F1.poster.jpg>

References

- Amit, S., Hatzubai, A., Birman, Y., Andersen, J. S., Ben-Shushan, E., Mann, M., Ben-Neriah, Y. and Alkalay, I. (2002). Axin-mediated CKI phosphorylation of beta-catenin at Ser 45: a molecular switch for the Wnt pathway. *Genes Dev.* **16**, 1066-1076.
- Andl, T., Reddy, S. T., Gaddapara, T. and Millar, S. E. (2002). WNT signals are required for the initiation of hair follicle development. *Dev. Cell* **2**, 643-653.
- Angers, S. and Moon, R. T. (2009). Proximal events in Wnt signal transduction. *Nat. Rev. Mol. Cell Biol.* **10**, 468-477.
- Bafico, A., Liu, G., Yaniv, A., Gazit, A. and Aaronson, S. A. (2001). Novel mechanism of Wnt signalling inhibition mediated by Dickkopf-1 interaction with LRP6/Arrow. *Nat. Cell Biol.* **3**, 683-686.
- Balemans, W., Patel, N., Ebeling, M., Van Hul, E., Wuyts, W., Lacza, C., Dioszegi, M., Dikkers, F. G., Hildering, P., Willems, P. J. et al. (2002). Identification of a 52 kb deletion downstream of the SOST gene in patients with van Buchem disease. *J. Med. Genet.* **39**, 91-97.
- Bänziger, C., Soldini, D., Schütt, C., Zipperlen, P., Hausmann, G. and Basler, K. (2006). Wntless, a conserved membrane protein dedicated to the secretion of Wnt proteins from signaling cells. *Cell* **125**, 509-522.
- Barker, N., van Es, J. H., Kuipers, J., Kujala, P., van den Born, M., Cozijnsen, M., Haegebarth, A., Korving, J., Begthel, H., Peters, P. J. et al. (2007). Identification of stem cells in small intestine and colon by marker gene Lgr5. *Nature* **449**, 1003-1007.
- Barker, N., Huch, M., Kujala, P., van de Wetering, M., Snippert, H. J., van Es, J. H., Sato, T., Stange, D. E., Begthel, H., van den Born, M. et al. (2010). Lgr5 +ve stem cells drive self-renewal in the stomach and build long-lived gastric units in vitro. *Cell Stem Cell* **6**, 25-36.
- Baron, R. and Kneissel, M. (2013). WNT signaling in bone homeostasis and disease: from human mutations to treatments. *Nat. Med.* **19**, 179-192.
- Bartscherer, K., Pelte, N., Ingelfinger, D. and Boutros, M. (2006). Secretion of Wnt ligands requires Evi, a conserved transmembrane protein. *Cell* **125**, 523-533.
- Behrens, J. (1998). Functional interaction of an axin homolog, conductin, with β -catenin, APC, and GSK3. *Science* **280**, 596-599.
- Behrens, J., von Kries, J. P., Kühl, M., Bruhn, L., Wedlich, D., Grosschedl, R. and Birchmeier, W. (1996). Functional interaction of β -catenin with the transcription factor LEF-1. *Nature* **382**, 638-642.
- Bennett, C. N., Longo, K. A., Wright, W. S., Suva, L. J., Lane, T. F., Hankenson, K. D. and MacDougald, O. A. (2005). Regulation of osteoblastogenesis and bone mass by Wnt10b. *Proc. Natl. Acad. Sci. USA* **102**, 3324-3329.
- Bhanot, P., Brink, M., Samos, C. H., Hsieh, J.-C., Wang, Y., Macke, J. P., Andrew, D., Nathans, J. and Nusse, R. (1996). A new member of the frizzled family from Drosophila functions as a Wingless receptor. *Nature* **382**, 225-230.
- Biechele, S., Cockburn, K., Lanner, F., Cox, B. J. and Rossant, J. (2013). Porcn-dependent Wnt signaling is not required prior to mouse gastrulation. *Development* **140**, 2961-2971.

- Boyden, L. M., Mao, J., Belsky, J., Mitzner, L., Farhi, A., Mitnick, M. A., Wu, D., Inogna, K. and Lifton, R. P. (2002). High bone density due to a mutation in LDL-receptor-related protein 5. *N. Engl. J. Med.* **346**, 1513-1521.
- Brunkow, M. E., Gardner, J. C., Van Ness, J., Paepfer, B. W., Kovacevich, B. R., Proll, S., Skonier, J. E., Zhao, L., Sabo, P. J., Fu, Y.-H. et al. (2001). Bone dysplasia sclerosteosis results from loss of the SOST gene product, a novel cysteine knot-containing protein. *Am. J. Hum. Genet.* **68**, 577-589.
- Cancer Genome Atlas Network (2012). Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **487**, 330-337.
- Carmon, K. S., Gong, X., Lin, Q., Thomas, A. and Liu, Q. (2011). R-spondins function as ligands of the orphan receptors LGR4 and LGR5 to regulate Wnt/beta-catenin signaling. *Proc. Natl. Acad. Sci. USA* **108**, 11452-11457.
- Cavallo, R. A., Cox, R. T., Moline, M. M., Roose, J., Polevoy, G. A., Clevers, H., Peifer, M. and Bejsovec, A. (1998). Drosophila Tcf and Groucho interact to repress Wingless signalling activity. *Nature* **395**, 604-608.
- Chen, S., Bubeck, D., MacDonald, B. T., Liang, W.-X., Mao, J.-H., Malinauskas, T., Llorca, O., Aricescu, A. R., Siebold, C., He, X. et al. (2011). Structural and functional studies of LRP6 ectodomain reveal a platform for Wnt signaling. *Dev. Cell* **21**, 848-861.
- Chen, P.-H., Chen, X., Lin, Z., Fang, D. and He, X. (2013). The structural basis of R-spondin recognition by LGR5 and RNF43. *Genes Dev.* **27**, 1345-1350.
- Chia, I. V. and Costantini, F. (2005). Mouse axin and axin2/conductin proteins are functionally equivalent in vivo. *Mol. Cell. Biol.* **25**, 4371-4376.
- Clevers, H. (2013). The intestinal crypt, a prototype stem cell compartment. *Cell* **154**, 274-284.
- Clevers, H., Loh, K. M. and Nusse, R. (2014). Stem cell signaling. An integral program for tissue renewal and regeneration: Wnt signaling and stem cell control. *Science* **346**, 1248012.
- Cong, F., Schweizer, L. and Varmus, H. (2004). Wnt signals across the plasma membrane to activate the beta-catenin pathway by forming oligomers containing its receptors, Frizzled and LRP. *Development* **131**, 5103-5115.
- Cotsarelis, G., Sun, T.-T. and Lavker, R. M. (1990). Label-retaining cells reside in the bulge area of pilosebaceous unit: implications for follicular stem cells, hair cycle, and skin carcinogenesis. *Cell* **61**, 1329-1337.
- Cselenyi, C. S., Jernigan, K. K., Tahinci, E., Thorne, C. A., Lee, L. A. and Lee, E. (2008). LRP6 transduces a canonical Wnt signal independently of Axin degradation by inhibiting GSK3's phosphorylation of -catenin. *Proc. Natl. Acad. Sci. USA* **105**, 8032-8037.
- Cui, Y., Niziolek, P. J., MacDonald, B. T., Zylstra, C. R., Alenina, N., Robinson, D. R., Zhong, Z., Matthes, S., Jacobsen, C. M., Conlon, R. A. et al. (2011). Lrp5 functions in bone to regulate bone mass. *Nat. Med.* **17**, 684-691.
- Davidson, G., Wu, W., Shen, J., Bilic, J., Fenger, U., Stanek, P., Glinka, A. and Niehrs, C. (2005). Casein kinase 1 gamma couples Wnt receptor activation to cytoplasmic signal transduction. *Nature* **438**, 867-872.
- Davidson, G., Shen, J., Huang, Y.-L., Su, Y., Karaulanov, E., Bartscherer, K., Hassler, C., Stanek, P., Boutros, M. and Niehrs, C. (2009). Cell cycle control of wnt receptor activation. *Dev. Cell* **17**, 788-799.
- Davidson, K. C., Adams, A. M., Goodson, J. M., McDonald, C. E., Potter, J. C., Berndt, J. D., Biechele, T. L., Taylor, R. J. and Moon, R. T. (2012). Wnt1 -catenin signaling promotes differentiation, not self-renewal, of human embryonic stem cells and is repressed by Oct4. *Proc. Natl. Acad. Sci. USA* **109**, 4485-4490.
- Day, T. F., Guo, X., Garrett-Beal, L. and Yang, Y. (2005). Wnt/beta-catenin signaling in mesenchymal progenitors controls osteoblast and chondrocyte differentiation during vertebrate skeletogenesis. *Dev. Cell* **8**, 739-750.
- de Lau, W., Barker, N., Low, T. Y., Koo, B.-K., Li, V. S. W., Teunissen, H., Kujala, P., Haegbarth, A., Peters, P. J., van de Wetering, M. et al. (2011). Lgr5 homologues associate with Wnt receptors and mediate R-spondin signalling. *Nature* **476**, 293-297.
- Enshell-Seiffers, D., Lindon, C., Kashiwagi, M. and Morgan, B. A. (2010). beta-catenin activity in the dermal papilla regulates morphogenesis and regeneration of hair. *Dev. Cell* **18**, 633-642.
- Farin, H. F., Jordens, I., Mosa, M. H., Basak, O., Korving, J., Tauriello, D. V. F., de Punder, K., Angers, S., Peters, P. J., Maurice, M. M. et al. (2016). Visualization of a short-range Wnt gradient in the intestinal stem-cell niche. *Nature* **530**, 340-343.
- Fevr, T., Robine, S., Louvard, D. and Huelsken, J. (2007). Wnt/beta-catenin is essential for intestinal homeostasis and maintenance of intestinal stem cells. *Mol. Cell. Biol.* **27**, 7551-7559.
- Flack, J. E., Mieszczynek, J., Novcic, N. and Bienz, M. (2017). Wnt-dependent inactivation of the groucho/TLE co-repressor by the HECT E3 ubiquitin ligase Hyd/UBR5. *Mol. Cell* **67**, 181-193.e5.
- Genikhovich, G. and Technau, U. (2017). On the evolution of bilaterality. *Development* **144**, 3392-3404.
- Glinka, A., Wu, W., Delius, H., Monaghan, A. P., Blumenstock, C. and Niehrs, C. (1998). Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction. *Nature* **391**, 357-362.
- Glinka, A., Dolde, C., Kirsch, N., Huang, Y.-L., Kazanskaya, O., Ingelfinger, D., Boutros, M., Cruciat, C.-M. and Niehrs, C. (2011). LGR4 and LGR5 are R-spondin receptors mediating Wnt/beta-catenin and Wnt/PCP signalling. *EMBO Rep.* **12**, 1055-1061.
- Glorieux, F. H., Devogelaer, J. P. and Durigova, M. (2017). BPS804 anti-sclerostin antibody in adults with moderate osteogenesis imperfecta: results of a randomized Phase 2a trial. *J. Bone Miner Res.* **32**, 1496-1504.
- Goentoro, L. and Kirschner, M. W. (2009). Evidence that fold-change, and not absolute level, of beta-catenin dictates Wnt signaling. *Mol. Cell* **36**, 872-884.
- Gong, Y., Slee, R. B., Fukui, N., Rawadi, G., Roman-Roman, S., Reginato, A. M., Wang, H., Cundy, T., Glorieux, F. H., Lev, D. et al. (2001). LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. *Cell* **107**, 513-523.
- González-Sancho, J. M., Aguilera, O., García, J. M., Pendas-Franco, N., Peña, C., Cal, S., de Herreros, A. G., Bonilla, F. and Muñoz, A. (2005). The Wnt antagonist DICKKOPF-1 gene is a downstream target of beta-catenin/TCF and is downregulated in human colon cancer. *Oncogene* **24**, 1098-1103.
- Goodman, R. M., Thombre, S., Firtina, Z., Gray, D., Betts, D., Roebuck, J., Spana, E. P. and Selva, E. M. (2006). Sprinter: a novel transmembrane protein required for Wg secretion and signaling. *Development* **133**, 4901-4911.
- Greco, V., Chen, T., Rendl, M., Schober, M., Pasolli, H. A., Stokes, N., Dela Cruz-Racelis, J. and Fuchs, E. (2009). A two-step mechanism for stem cell activation during hair regeneration. *Cell Stem Cell* **4**, 155-169.
- Green, D., Whitener, A. E., Mohanty, S. and Lekven, A. C. (2015). Vertebrate nervous system posteriorization: Grading the function of Wnt signaling. *Dev. Dyn.* **244**, 507-512.
- Hao, H.-X., Xie, Y., Zhang, Y., Charlat, O., Oster, E., Avello, M., Lei, H., Mickanin, C., Liu, D., Ruffner, H. et al. (2012). ZNRF3 promotes Wnt receptor turnover in an R-spondin-sensitive manner. *Nature* **485**, 195-200.
- Hart, M. J., de los Santos, R., Albert, I. N., Rubinfeld, B. and Polakis, P. (1998). Downregulation of beta-catenin by human Axin and its association with the APC tumor suppressor, beta-catenin and GSK3beta. *Curr. Biol.* **8**, 573-581.
- Hart, M., Concordet, J.-P., Lassot, I., Albert, I., del los Santos, R., Durand, H., Perret, C., Rubinfeld, B., Margottin, F., Benarous, R. et al. (1999). The F-box protein beta-TrCP associates with phosphorylated beta-catenin and regulates its activity in the cell. *Curr. Biol.* **9**, 207-211.
- Herr, P. and Basler, K. (2012). Porcupine-mediated lipidation is required for Wnt recognition by Wls. *Dev. Biol.* **361**, 392-402.
- Hill, T. P., Später, D., Taketo, M. M., Birchmeier, W. and Hartmann, C. (2005). Canonical Wnt/beta-catenin signaling prevents osteoblasts from differentiating into chondrocytes. *Dev. Cell* **8**, 727-738.
- Hsieh, J.-C., Kodjacobian, L., Rebbert, M. L., Rattner, A., Smallwood, P. M., Samos, C. H., Nusse, R., Dawid, I. B. and Nathans, J. (1999). A new secreted protein that binds to Wnt proteins and inhibits their activities. *Nature* **398**, 431-436.
- Huelsken, J., Vogel, R., Erdmann, B., Cotsarelis, G. and Birchmeier, W. (2001). beta-Catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. *Cell* **105**, 533-545.
- Jaks, V., Barker, N., Kasper, M., van Es, J. H., Snippert, H. J., Clevers, H. and Toftgård, R. (2008). Lgr5 marks cycling, yet long-lived, hair follicle stem cells. *Nat. Genet.* **40**, 1291-1299.
- Janda, C. Y., Waghray, D., Levin, A. M., Thomas, C. and Garcia, K. C. (2012). Structural basis of Wnt recognition by Frizzled. *Science* **337**, 59-64.
- Janda, C. Y., Dang, L. T., You, C., Chang, J., de Lau, W., Zhong, Z. A., Yan, K. S., Marecic, O., Siepe, D., Li, X. et al. (2017). Surrogate Wnt agonists that phenocopy canonical Wnt and beta-catenin signalling. *Nature* **545**, 234-237.
- Jho, E.-H., Zhang, T., Domon, C., Joo, C.-K., Freund, J.-N. and Costantini, F. (2002). Wnt1 -catenin/Tcf signaling induces the transcription of Axin2, a negative regulator of the signaling pathway. *Mol. Cell. Biol.* **22**, 1172-1183.
- Kabiri, Z., Greicius, G., Madan, B., Biechele, S., Zhong, Z., Zaribafzadeh, H., Edison, Aliyev, J., Wu, Y., Bunte, R. et al. (2014). Stroma provides an intestinal stem cell niche in the absence of epithelial Wnts. *Development* **141**, 2206-2215.
- Kakugawa, S., Langton, P. F., Zebisch, M., Howell, S. A., Chang, T.-H., Liu, Y., Feizi, T., Bineva, G., O'Reilly, N., Snijders, A. P. et al. (2015). Notum deacylates Wnt proteins to suppress signalling activity. *Nature* **519**, 187-192.
- Kishimoto, J., Burgeson, R. E. and Morgan, B. A. (2000). Wnt signaling maintains the hair-inducing activity of the dermal papilla. *Genes Dev.* **14**, 1181-1185.
- Klingensmith, J., Nusse, R. and Perrimon, N. (1994). The Drosophila segment polarity gene dishevelled encodes a novel protein required for response to the wingless signal. *Genes Dev.* **8**, 118-130.
- Koo, B.-K., Spit, M., Jordens, I., Low, T. Y., Stange, D. E., van de Wetering, M., van Es, J. H., Mohammed, S., Heck, A. J. R., Maurice, M. M. et al. (2012). Tumour suppressor RNF43 is a stem-cell E3 ligase that induces endocytosis of Wnt receptors. *Nature* **488**, 665-669.
- Korinek, V., Barker, N., Moerer, P., van Donselaar, E., Huls, G., Peters, P. J. and Clevers, H. (1998). Depletion of epithelial stem-cell compartments in the small intestine of mice lacking Tcf-4. *Nat. Genet.* **19**, 379-383.
- Kosinski, C., Li, V. S. W., Chan, A. S. Y., Zhang, J., Ho, C., Tsui, W. Y., Chan, T. L., Mifflin, R. C., Powell, D. W., Yuen, S. T. et al. (2007). Gene expression patterns of human colon tops and basal crypts and BMP antagonists as intestinal stem cell niche factors. *Proc. Natl. Acad. Sci. USA* **104**, 15418-15423.
- Krasnow, R. E., Wong, L. L. and Adler, P. N. (1995). Dishevelled is a component of the frizzled signaling pathway in Drosophila. *Development* **121**, 4095-4102.
- Kurek, D., Neagu, A., Tastemel, M., Tüysüz, N., Lehmann, J., van de Werken, H. J. G., Philipsen, S., van der Linden, R., Maas, A., van IJcken, W. F. J. et al. (2015). Endogenous WNT signals mediate BMP-induced and spontaneous

- differentiation of epiblast stem cells and human embryonic stem cells. *Stem Cell Rep.* **4**, 114-128.
- Lebensohn, A. M. and Rohatgi, R.** (2018). R-spondins can potentiate WNT signaling without LGRs. *Elife* **7**, e33126.
- Lee, E., Salic, A., Krüger, R., Heinrich, R. and Kirschner, M. W.** (2003). The roles of APC and Axin derived from experimental and theoretical analysis of the Wnt pathway. *PLoS Biol.* **1**, e10.
- Leyns, L., Bouwmeester, T., Kim, S.-H., Piccolo, S. and De Robertis, E. M.** (1997). Frzb-1 is a secreted antagonist of Wnt signaling expressed in the spemann organizer. *Cell* **88**, 747-756.
- Lien, W.-H., Polak, L., Lin, M., Lay, K., Zheng, D. and Fuchs, E.** (2014). In vivo transcriptional governance of hair follicle stem cells by canonical Wnt regulators. *Nat. Cell Biol.* **16**, 179-190.
- Lim, X., Tan, S. H., Koh, W. L. C., Chau, R. M. W., Yan, K. S., Kuo, C. J., van Amerongen, R., Klein, A. M. and Nusse, R.** (2013). Interfollicular epidermal stem cells self-renew via autocrine Wnt signaling. *Science* **342**, 1226-1230.
- Lim, X., Tan, S. H., Yu, K. L., Lim, S. B. H. and Nusse, R.** (2016). Axin2 marks quiescent hair follicle bulge stem cells that are maintained by autocrine Wnt/ β -catenin signaling. *Proc. Natl. Acad. Sci. USA* **113**, E1498-E1505.
- Little, R. D., Folz, C., Manning, S. P., Swain, P. M., Zhao, S.-C., Eustace, B., Lappe, M. M., Spitzer, L., Zweier, S., Braunschweiger, K. et al.** (2002). A mutation in the LDL receptor-related protein 5 gene results in the autosomal dominant high-bone-mass trait. *Am. J. Hum. Genet.* **70**, 11-19.
- Liu, C., Li, Y., Semenov, M., Han, C., Baeg, G.-H., Tan, Y., Zhang, Z., Lin, X. and He, X.** (2002). Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell* **108**, 837-847.
- Logan, C. Y. and Nusse, R.** (2004). The Wnt signaling pathway in development and disease. *Annu. Rev. Cell Dev. Biol.* **20**, 781-810.
- Loots, G. G., Kneissel, M., Keller, H., Baptist, M., Chang, J., Collette, N. M., Ovcharenko, D., Plajzer-Frick, I. and Rubin, E. M.** (2005). Genomic deletion of a long-range bone enhancer misregulates sclerostin in Van Buchem disease. *Genome Res.* **15**, 928-935.
- Lustig, B., Jerchow, B., Sachs, M., Weiler, S., Pietsch, T., Karsten, U., van de Wetering, M., Clevers, H., Schlag, P. M., Birchmeier, W. et al.** (2002). Negative feedback loop of Wnt signaling through upregulation of conductin/axin2 in colorectal and liver tumors. *Mol. Cell Biol.* **22**, 1184-1193.
- Lyashenko, N., Winter, M., Migliorini, D., Biechele, T., Moon, R. T. and Hartmann, C.** (2011). Differential requirement for the dual functions of β -catenin in embryonic stem cell self-renewal and germ layer formation. *Nat. Cell Biol.* **13**, 753-761.
- Lyle, S., Christofidou-Solomidou, M., Liu, Y., Elder, D. E., Albelda, S. and Cotsarelis, G.** (1998). The C8/144B monoclonal antibody recognizes cytokeratin 15 and defines the location of human hair follicle stem cells. *J. Cell Sci.* **111**, 3179-3188.
- MacDonald, B. T. and He, X.** (2012). Frizzled and LRP5/6 receptors for Wnt/ β -catenin signaling. *Cold Spring Harb. Perspect. Biol.* **4**, a007880.
- Mao, J., Wang, J., Liu, B., Pan, W., Farr, G. H., III, Flynn, C., Yuan, H., Takada, S., Kimelman, D., Li, L. et al.** (2001). Low-density lipoprotein receptor-related protein-5 binds to Axin and regulates the canonical Wnt signaling pathway. *Mol. Cell* **7**, 801-809.
- McMahon, A. P. and Moon, R. T.** (1989). Ectopic expression of the proto-oncogene int-1 in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* **58**, 1075-1084.
- Molenaar, M., van de Wetering, M., Oosterwegel, M., Peterson-Maduro, J., Godsave, S., Korinek, V., Roose, J., Destree, O. and Clevers, H.** (1996). XTcf-3 transcription factor mediates β -catenin-induced axis formation in *xenopus* embryos. *Cell* **86**, 391-399.
- Moon, R. T. and Kimelman, D.** (1998). From cortical rotation to organizer gene expression: toward a molecular explanation of axis specification in *Xenopus*. *BioEssays* **20**, 536-545.
- Morin, P. J.** (1997). Activation of beta -Catenin-Tcf signaling in colon cancer by mutations in beta -catenin or APC. *Science* **275**, 1787-1790.
- Munemitsu, S., Albert, I., Souza, B., Rubinfeld, B. and Polakis, P.** (1995). Regulation of intracellular beta-catenin levels by the adenomatous polyposis coli (APC) tumor-suppressor protein. *Proc. Natl. Acad. Sci. USA* **92**, 3046-3050.
- Nakamura, Y., de Paiva Alves, E., Veenstra, G. J. C. and Hoppler, S.** (2016). Tissue- and stage-specific Wnt target gene expression is controlled subsequent to β -catenin recruitment to cis-regulatory modules. *Development* **143**, 1914-1925.
- Nichols, J. and Smith, A.** (2009). Naive and primed pluripotent states. *Cell Stem Cell* **4**, 487-492.
- Niehrs, C.** (2010). On growth and form: a Cartesian coordinate system of Wnt and BMP signaling specifies bilaterian body axes. *Development* **137**, 845-857.
- Nusse, R.** (2001). An ancient cluster of Wnt paralogues. *Trends Genet.* **17**, 443.
- Nusse, R. and Clevers, H.** (2017). Wnt/ β -catenin signaling, disease, and emerging therapeutic modalities. *Cell* **169**, 985-999.
- Nüsslein-Volhard, C. and Wieschaus, E.** (1980). Mutations affecting segment number and polarity in *Drosophila*. *Nature* **287**, 795-801.
- Oshima, H., Rochat, A., Kedzija, C., Kobayashi, K. and Barrandon, Y.** (2001). Morphogenesis and renewal of hair follicles from adult multipotent stem cells. *Cell* **104**, 233-245.
- Piao, S., Lee, S.-H., Kim, H., Yum, S., Stamos, J. L., Xu, Y., Lee, S.-J., Lee, J., Oh, S., Han, J.-K. et al.** (2008). Direct inhibition of GSK3 β by the phosphorylated cytoplasmic domain of LRP6 in Wnt/ β -catenin signaling. *PLoS One* **3**, e4046.
- Pinson, K. I., Brennan, J., Monkley, S., Avery, B. J. and Skarnes, W. C.** (2000). An LDL-receptor-related protein mediates Wnt signalling in mice. *Nature* **407**, 535-538.
- Pinto, D., Gregorieff, A., Begthel, H. and Clevers, H.** (2003). Canonical Wnt signals are essential for homeostasis of the intestinal epithelium. *Genes Dev.* **17**, 1709-1713.
- Povelones, M. and Nusse, R.** (2005). The role of the cysteine-rich domain of Frizzled in Wingless-Armadillo signaling. *EMBO J.* **24**, 3493-3503.
- Proffitt, K. D. and Virshup, D. M.** (2012). Precise regulation of porcupine activity is required for physiological Wnt signaling. *J. Biol. Chem.* **287**, 34167-34178.
- Rattner, A., Hsieh, J.-C., Smallwood, P. M., Gilbert, D. J., Copeland, N. G., Jenkins, N. A. and Nathans, J.** (1997). A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors. *Proc. Natl. Acad. Sci. USA* **94**, 2859-2863.
- Recker, R. R., Benson, C. T., Matsumoto, T., Bolognese, M. A., Robins, D. A., Alam, J., Chiang, A. Y., Hu, L., Kregge, J. H., Sowa, H. et al.** (2015). A randomized, double-blind phase 2 clinical trial of bloszumab, a sclerostin antibody, in postmenopausal women with low bone mineral density. *J. Bone Miner. Res.* **30**, 216-224.
- Rijsewijk, F., Schuermans, M., Wagenaar, E., Parren, P., Weigel, D. and Nusse, R.** (1987). The *Drosophila* homolog of the mouse mammary oncogene int-1 is identical to the segment polarity gene wingless. *Cell* **50**, 649-657.
- Roose, J., Molenaar, M., Peterson, J., Hurenkamp, J., Brantjes, H., Moerer, P., van de Wetering, M., Destree, O. and Clevers, H.** (1998). The *Xenopus* Wnt effector XTcf-3 interacts with Groucho-related transcriptional repressors. *Nature* **395**, 608-612.
- Roussel, R., Mack, J. A., Wharton, K. A., Jr, Axelrod, J. D., Cadigan, K. M., Fish, M. P., Nusse, R. and Scott, M. P.** (2001). Naked cuticle targets dishevelled to antagonize Wnt signal transduction. *Genes Dev.* **15**, 658-671.
- Rubinfeld, B., Souza, B., Albert, I., Müller, O., Chamberlain, S. H., Masiarz, F. R., Munemitsu, S. and Polakis, P.** (1993). Association of the APC gene product with beta-catenin. *Science* **262**, 1731-1734.
- Rubinfeld, B., Albert, I., Porfiri, E., Fiol, C., Munemitsu, S. and Polakis, P.** (1996). Binding of GSK3beta to the APC-beta-catenin complex and regulation of complex assembly. *Science* **272**, 1023-1026.
- Saag, K. G., Petersen, J., Brandi, M. L., Karaplis, A. C., Lorentzon, M., Thomas, T., Maddox, J., Fan, M., Meisner, P. D. and Grauer, A.** (2017). Romosozumab or alendronate for fracture prevention in women with osteoporosis. *N. Engl. J. Med.* **377**, 1417-1427.
- Sato, N., Meijer, L., Skaltsounis, L., Greengard, P. and Brivanlou, A. H.** (2004). Maintenance of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling by a pharmacological GSK-3-specific inhibitor. *Nat. Med.* **10**, 55-63.
- Sato, T., Vries, R. G., Snippert, H. J., van de Wetering, M., Barker, N., Stange, D. E., van Es, J. H., Abo, A., Kujala, P., Peters, P. J. et al.** (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* **459**, 262-265.
- Satoh, K., Kasai, M., Ishidao, T., Tago, K., Ohwada, S., Hasegawa, Y., Senda, T., Takada, S., Nada, S., Nakamura, T. et al.** (2004). Anteriorization of neural fate by inhibitor of beta-catenin and T cell factor (ICAT), a negative regulator of Wnt signaling. *Proc. Natl. Acad. Sci. USA* **101**, 8017-8021.
- Seménov, M. V., Tamai, K., Brott, B. K., Kühn, M., Sokol, S. and He, X.** (2001). Head inducer Dickkopf-1 is a ligand for Wnt coreceptor LRP6. *Curr. Biol.* **11**, 951-961.
- Sharma, R. P. and Chopra, V. L.** (1976). Effect of the Wingless (wg1) mutation on wing and haltere development in *Drosophila melanogaster*. *Dev. Biol.* **48**, 461-465.
- Stamos, J. L. and Weis, W. I.** (2012). The -catenin destruction complex. *Cold Spring Harb. Perspect. Biol.* **5**, a007898-a007898.
- Stamos, J. L., Chu, M. L.-H., Enos, M. D., Shah, N. and Weis, W. I.** (2014). Structural basis of GSK-3 inhibition by N-terminal phosphorylation and by the Wnt receptor LRP6. *Elife* **3**, e01998.
- Tabar, V. and Studer, L.** (2014). Pluripotent stem cells in regenerative medicine: challenges and recent progress. *Nat. Rev. Genet.* **15**, 82-92.
- Tamai, K., Semenov, M., Kato, Y., Spokony, R., Liu, C., Katsuyama, Y., Hess, F., Saint-Jeannet, J. P. and He, X.** (2000). LDL-receptor-related proteins in Wnt signal transduction. *Nature* **407**, 530-535.
- Tamai, K., Zeng, X., Liu, C., Zhang, X., Harada, Y., Chang, Z. and He, X.** (2004). A mechanism for Wnt coreceptor activation. *Mol. Cell* **13**, 149-156.
- Tao, Q., Yokota, C., Puck, H., Kofron, M., Birsoy, B., Yan, D., Asashima, M., Wylie, C. C., Lin, X. and Heasman, J.** (2005). Maternal wnt11 activates the canonical wnt signaling pathway required for axis formation in *Xenopus* embryos. *Cell* **120**, 857-871.
- Tauriello, D. V. F., Jordens, I., Kirchner, K., Slootstra, J. W., Kruitwagen, T., Bouwman, B. A. M., Noutsou, M., Rüdiger, S. G. D., Schwamborn, K., Schambony, A. et al.** (2012). Wnt/ β -catenin signaling requires interaction of the

- Dishevelled DEP domain and C terminus with a discontinuous motif in Frizzled. *Proc. Natl. Acad. Sci. USA* **109**, E812-E820.
- ten Berge, D., Kurek, D., Blauwkamp, T., Koole, W., Maas, A., Eroglu, E., Siu, R. K. and Nusse, R.** (2011). Embryonic stem cells require Wnt proteins to prevent differentiation to epiblast stem cells. *Nat. Cell Biol.* **13**, 1070-1075.
- Tu, X., Delgado-Calle, J., Condon, K. W., Maycas, M., Zhang, H., Carlesso, N., Taketo, M. M., Burr, D. B., Plotkin, L. I. and Bellido, T.** (2015). Osteocytes mediate the anabolic actions of canonical Wnt/ β -catenin signaling in bone. *Proc. Natl. Acad. Sci. USA* **112**, E478-E486.
- Tumbar, T., Guasch, G., Greco, V., Blanpain, C., Lowry, W. E., Rendl, M. and Fuchs, E.** (2004). Defining the epithelial stem cell niche in skin. *Science* **303**, 359-363.
- van Amerongen, R., Bowman, A. N. and Nusse, R.** (2012). Developmental stage and time dictate the fate of Wnt/ β -catenin-responsive stem cells in the mammary gland. *Cell Stem Cell* **11**, 387-400.
- van de Wetering, M., Cavallo, R., Dooijes, D., van Beest, M., van Es, J., Loureiro, J., Ypma, A., Hursh, D., Jones, T., Bejsovec, A. et al.** (1997). Armadillo coactivates transcription driven by the product of the Drosophila segment polarity gene dTCF. *Cell* **88**, 789-799.
- van den Heuvel, M., Harryman-Samos, C., Klingensmith, J., Perrimon, N. and Nusse, R.** (1993). Mutations in the segment polarity genes wingless and porcupine impair secretion of the wingless protein. *EMBO J.* **12**, 5293-5302.
- Van der Flier, L. G., Sabates-Bellver, J., Oving, I., Haegebarth, A., De Palo, M., Anti, M., Van Gijn, M. E., Suijkerbuijk, S., Van de Wetering, M., Marra, G. et al.** (2007). The intestinal Wnt/TCF signature. *Gastroenterology* **132**, 628-632.
- Van Mater, D., Kolligs, F. T., Dlugosz, A. A. and Fearon, E. R.** (2003). Transient activation of beta-catenin signaling in cutaneous keratinocytes is sufficient to trigger the active growth phase of the hair cycle in mice. *Genes Dev.* **17**, 1219-1224.
- Van Raay, T. J., Fortino, N. J., Miller, B. W., Ma, H., Lau, G., Li, C., Franklin, J. L., Attisano, L., Solnica-Krezel, L. and Coffey, R. J.** (2011). Naked1 antagonizes Wnt signaling by preventing nuclear accumulation of β -catenin. *PLoS One* **6**, e18650.
- Veltri, A., Lang, C. and Lien, W.-H.** (2018). Concise review: Wnt signaling pathways in skin development and epidermal stem cells. *Stem Cells* **36**, 22-35.
- Wang, S., Krinks, M., Lin, K., Luyten, F. P. and Moos, M. Jr.** (1997). Frzb, a secreted protein expressed in the Spemann organizer, binds and inhibits Wnt-8. *Cell* **88**, 757-766.
- Wang, B., Zhao, L., Fish, M., Logan, C. Y. and Nusse, R.** (2015). Self-renewing diploid Axin2(+) cells fuel homeostatic renewal of the liver. *Nature* **524**, 180-185.
- Wehrli, M., Dougan, S. T., Caldwell, K., O'Keefe, L., Schwartz, S., Vaizel-Ohayon, D., Schejter, E., Tomlinson, A. and DiNardo, S.** (2000). arrow encodes an LDL-receptor-related protein essential for Wingless signalling. *Nature* **407**, 527-530.
- Willert, K., Brown, J. D., Danenberg, E., Duncan, A. W., Weissman, I. L., Reya, T., Yates, J. R., III and Nusse, R.** (2003). Wnt proteins are lipid-modified and can act as stem cell growth factors. *Nature* **423**, 448-452.
- Williams, B. O.** (2017). LRP5: From bedside to bench to bone. *Bone* **102**, 26-30.
- Williams, L. A., Davis-Dusenbery, B. N. and Eggan, K. C.** (2012). SnapShot: directed differentiation of pluripotent stem cells. *Cell* **149**, 1174-1174.e1.
- Winston, J. T., Strack, P., Beer-Romero, P., Chu, C. Y., Elledge, S. J. and Harper, J. W.** (1999). The SCFbeta-TRCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in I κ Balpha and beta-catenin and stimulates I κ Balpha ubiquitination in vitro. *Genes Dev.* **13**, 270-283.
- Xu, Z., Robitaille, A. M., Berndt, J. D., Davidson, K. C., Fischer, K. A., Mathieu, J., Potter, J. C., Ruohola-Baker, H. and Moon, R. T.** (2016). Wnt/ β -catenin signaling promotes self-renewal and inhibits the primed state transition in naïve human embryonic stem cells. *Proc. Natl. Acad. Sci. USA* **113**, E6382-E6390.
- Yan, K. S., Janda, C. Y., Chang, J., Zheng, G. X. Y., Larkin, K. A., Luca, V. C., Chia, L. A., Mah, A. T., Han, A., Terry, J. M. et al.** (2017). Non-equivalence of Wnt and R-spondin ligands during Lgr5(+) intestinal stem-cell self-renewal. *Nature* **545**, 238-242.
- Yang-Snyder, J., Miller, J. R., Brown, J. D., Lai, C.-J. and Moon, R. T.** (1996). A frizzled homolog functions in a vertebrate Wnt signaling pathway. *Curr. Biol.* **6**, 1302-1306.
- Zebisch, M., Xu, Y., Krastev, C., MacDonald, B. T., Chen, M., Gilbert, R. J. C., He, X. and Jones, E. Y.** (2013). Structural and molecular basis of ZNRF3/RNF43 transmembrane ubiquitin ligase inhibition by the Wnt agonist R-spondin. *Nat. Commun.* **4**, 2787.
- Zeng, X., Tamai, K., Doble, B., Li, S., Huang, H., Habas, R., Okamura, H., Woodgett, J. and He, X.** (2005). A dual-kinase mechanism for Wnt co-receptor phosphorylation and activation. *Nature* **438**, 873-877.
- Zhang, X., Abreu, J. G., Yokota, C., MacDonald, B. T., Singh, S., Coburn, K. L. A., Cheong, S.-M., Zhang, M. M., Ye, Q.-Z., Hang, H. C. et al.** (2012). Tiki1 is required for head formation via Wnt cleavage-oxidation and inactivation. *Cell* **149**, 1565-1577.
- Zhang, X., Cheong, S.-M., Amado, N. G., Reis, A. H., MacDonald, B. T., Zebisch, M., Jones, E. Y., Abreu, J. G. and He, X.** (2015). Notum is required for neural and head induction via Wnt deacylation, oxidation, and inactivation. *Dev. Cell* **32**, 719-730.