

# Autosomal *P[ovo<sup>DI</sup>]* dominant female-sterile insertions in *Drosophila* and their use in generating germ-line chimeras

Tze-Bin Chou, Elizabeth Noll and Norbert Perrimon

Howard Hughes Medical Institute, Department of Genetics, Harvard Medical School, 200 Longwood Avenue, Boston, Massachusetts 02115, USA

## SUMMARY

The 'dominant female-sterile' technique used to generate germ-line mosaics in *Drosophila* is a powerful tool to determine the tissue specificity (germ line versus somatic) of recessive female-sterile mutations as well as to analyze the maternal effect of recessive zygotic lethal mutations. This technique requires the availability of germ-line-dependent, dominant female-sterile (DFS) mutations that block egg laying but do not affect viability. To date only one X-linked mutation, *ovo<sup>DI</sup>* has been isolated that completely fulfills these criteria. Thus the 'DFS technique' has been largely limited to the X-chromosome. To extend this technique to the autosomes,

we have cloned the *ovo<sup>DI</sup>* mutation into a P-element vector and recovered fully expressed *P[ovo<sup>DI</sup>]* insertions on each autosomal arm. We describe the generation of these *P[ovo<sup>DI</sup>]* strains as well as demonstrate their use in generating germ-line chimeras. Specifically, we show that the *Gap1* gene, which encodes a *Drosophila* homologue of mammalian GTPase-activating protein, is required in somatic follicle cells for embryonic dorsoventral polarity determination.

Key words: *Drosophila*, oogenesis, mosaic, female-sterile

## INTRODUCTION

Formation of the *Drosophila* egg involves the contribution of two different ovarian cell types, the nurse cell-oocyte complex of germ-line origin derived from the embryonic pole cells, and the follicle cells derived from the embryonic mesoderm (King, 1970). The early separation of these two cell types makes it possible to generate chimeras that have germ-line and follicle cells of different genotypes. Generation of such chimeras has been important for two sets of experiments. First, they allow the determination of the tissue specificity of recessive female-sterile (fs) mutations (Wieschaus et al., 1981; Perrimon and Gans, 1983). Saturation screens have provided us with a large collection of fs mutations involved during oogenesis and embryonic patterning (Gans et al., 1975; Mohler, 1977; Komitopoulou et al., 1983; Perrimon et al., 1986; Schupbach and Wieschaus, 1989). Due to the dual origin of the cell types that form the egg and their complex interactions in oogenesis, it is critical to determine the tissue specificity of these fs mutations by generating germ-line mosaics. Second, germ-line chimeras allow one to analyze the maternal effect of zygotic lethal mutations (Perrimon et al., 1984, 1989). Approximately 85% of the genes in *Drosophila* are zygotic lethals when mutated and one direct way to determine the maternal function of these mutations is to analyze the phenotype of eggs derived from mosaic mothers that carry an homozygous mutant germ line.

Two techniques are currently used to generate germ-line

chimeras: pole cell transplantation in which mutant pole cells are transferred into an otherwise wild-type donor (Illmensee, 1973) and mitotic recombination (Wieschaus and Szabad, 1979). A critical aspect of both techniques is the ability to distinguish the eggs derived from homozygous mutant cells from the otherwise wild-type eggs. In the case of pole cell transplantation, this can be easily done by injecting the mutant pole cells into an animal that does not develop germ cells. Females lacking germ cells can be generated by using recessive or dominant female-sterile mutations that prevent pole cell formation or affect germ cell development. In the case of mitotic recombination, the most successful approach has been the use of germ-line-dependent DFS that do not lay eggs (the 'DFS' technique; Wieschaus, 1980; Perrimon and Gans, 1983). A mitotic recombination event occurring in the germ cells of females heterozygous for DFS will result in recombinant germ-line daughter cells that have eliminated the DFS mutation and thus can develop into normal eggs. If the female is *trans*-heterozygous for both a specific mutation (fs or zygotic lethal) and the DFS mutation, the mitotic exchange will result in simultaneous loss of the DFS mutation and homozygosity of the mutation. The tissue specificity of a fs mutation or the maternal effect phenotype of a zygotic lethal mutation can then be analyzed by examining the phenotypes of eggs and embryos derived from these germ-line clones.

The most widely used DFS mutation is the X-linked *ovo<sup>DI</sup>* mutation (or *Fs(1)K1237*, Busson et al., 1983; Perrimon, 1984). Females heterozygous for *ovo<sup>DI</sup>* never lay

eggs because they have rudimentary ovaries in which oogenesis is blocked at an early stage. The female sterility associated with the *ovo<sup>D1</sup>* mutation is strictly germ-line dependent such that flies carrying this mutation are perfectly viable and wild-type germ cells in an *ovo<sup>D1</sup>* heterozygous female produce wild-type eggs. In addition, germ cells heterozygous for *ovo<sup>D1</sup>* develop poorly such that, when a germ cell has eliminated *ovo<sup>D1</sup>*, it has a growth advantage and the resulting clone can usually give rise to a fully developed ovary (Busson et al., 1983; Perrimon, 1984). This overproliferation of wild-type cells in an *ovo<sup>D1</sup>* background has been a major advantage in using *ovo<sup>D1</sup>* to generate germ-line mosaics.

The frequency at which germ-line clones are recovered is an important consideration in designing experiments. Mitotic recombination events induced in *ovo<sup>D1</sup>* heterozygous animals following X-ray irradiation are generated at a frequency of approximately 5% of mosaic females among heterozygous *ovo<sup>D1</sup>* females (Perrimon, 1984). Recently, we have combined the use of the site-specific recombination system (Golic and Lindquist, 1989; Golic, 1991) to develop the 'FLP-DFS' technique (Chou and Perrimon, 1992) where the frequency of mosaics reaches almost 100%.

Screens for autosomal mutations associated with female sterility have identified a few DFS mutations, some of which have been used occasionally to generate germ-line mosaics (Yarger and King, 1971; Erdelyi and Szabad, 1989; Szabad et al., 1989). These DFS mutations in some cases affect viability or lay some eggs making their use difficult in detecting germ-line chimeras. In addition, clones induced using DFS mutations such as *Fs(2)D* (Yarger and King, 1971) and *Fs(2)I* (Szabad et al., 1987, 1989) are small because none show the regulation phenomena observed in the background of the *ovo<sup>D1</sup>* mutation (N. P. unpublished data).

To overcome the lack of appropriate autosomal DFS mutations to generate germ-line chimeras, we first attempted to create DFS mutations by selectively expressing the A chain of diphtheria toxin (DT-A) in the female germ line. DT-A has been shown to induce cell death in a cell autonomous manner (Palmiter et al., 1987) such that dominant female sterility could be achieved by expressing *DT-A* selectively in female germ cells. We tested a number of female germ-line-specific promoter elements including part of the promoter of the *Drosophila hsp26* gene (Frank et al., 1992) and an element from the *ovarian-tumor* promoter (Comer et al., 1992) to drive the expression of *DT-A* in the female germ line. In addition, we used the germ-line-specific splicing intron of the P-element transposase (Laski and Rubin 1989) or the sex-specific splicing intron of the *transformer* gene (Sosnowski et al., 1989) to further restrict female germ-line cell expression of the *DT-A* gene. However, we failed to recover P-element transformants of these constructs due to the strong toxicity of DT-A resulting from transient expression in injected embryos. After laborious attempts to obtain transformants of these constructs, we decided that transposition of the *ovo<sup>D1</sup>* mutation may be a more straightforward approach. Since early attempts to translocate genetically the *ovo<sup>D1</sup>* mutation failed due to the presence of regions of unusual genetic properties around the *ovo* gene (Oliver et al., 1988), we decided that

transposition of a cloned *ovo<sup>D1</sup>* mutation following P-element transformation might constitute a better approach. These experiments were possible since the *ovo* gene has been cloned (Mével-Ninio et al., 1991; Garfinkel et al., 1992). In this paper, we report the successful recovery of autosomal P-element insertions that carry the *ovo<sup>D1</sup>* mutation and demonstrate their use in generating germ-line mosaics.

## MATERIAL AND METHODS

### Construction and isolation of an *ovo<sup>D1</sup>* cosmid

*ovo<sup>D1</sup>* genomic DNA was isolated from *ovo<sup>D1</sup> v<sup>24</sup>/Y* males and prepared by standard methods (Sambrook et al., 1989). The construction of the *ovo<sup>D1</sup>* cosmid library followed the protocol of Ish-Horowicz and Burke (1981). *ovo<sup>D1</sup>* genomic DNA was partially digested with *Sau3A* and ligated into the pCosPeR vector (V. Pirrotta, personal communication) digested with *Bam*HI. In vitro packaging was performed using Gigapack II XL packaging extract from Stratagene. After titration of the cosmid library, the primary colonies were screened without amplification using 4 different PCR probes which cover the entire *ovo* transcription unit (Fig. 1). Coordinates for these primers followed that described in Mevel-Ninio et al. (1991). The 4 pairs of oligomers are as follows. *ovo*-1 primer: 5'-GTCGACTGCAACAGTTGGCC-3' and *ovo*-740 primer: 5'-CGAATCAACAAGTGATGAAA-3' produce a 740bp fragment; *ovo*-1768 primer: 5'-ATCAGCTAGGACCCGAGCAG-3' and *ovo*-2422 primer: 5'-GATGCTCCACCACCTCCGTT-3' produce a 654bp fragment; *ovo*-4481 primer: 5'-GGTTATC-GATGGCGCTTACC-3' and *ovo*-5119 primer: 5'-CGACGAATC-CTCATTGGCTC-3' produce a 638bp fragment; *ovo*-5719 primer: 5'-GAATAGGAACCATTTCGATCG-3' and *ovo*-6654 primer: 5'-ATGTTGTGTTTAGTATTTTG-3' produce a 935bp fragment. The 638bp PCR fragment was used first as a probe for primary colony screening. Direct colony PCR screening (Hamilton et al., 1991) was performed thereafter. Colonies surrounding the positive signals were scooped into 100 µl LB solution. Using the primers listed above, PCR reactions were performed immediately in 25 µl solution containing 1 µl colony suspension. Primers were used at 250 nM. Amplifications were done first with 2 minutes 30 seconds at 94°C for bacterial cell lysis followed by 30 cycles at 94°C for 1 minute; 54°C for 1 minute; 72°C for 3 minutes. Colonies that were positive with all 4 PCR products were used for the secondary screen using both 935bp and 654bp fragments separately as probes. One positive colony from this screen was confirmed again by PCR and used for the preparation of cosmid DNA.

### Subcloning of *ovo<sup>D1</sup>* genomic DNA fragment

A 7.2 kb genomic DNA fragment containing the putative 5 kb *ovo<sup>+</sup>* mRNA transcription unit can rescue the female sterility phenotype of *ovo<sup>-</sup>* mutations (Mével-Ninio et al., 1991). A 10.5 kb *ovo<sup>D1</sup>* genomic DNA fragment containing the *ovo<sup>D1</sup>* counterpart of the 7.2 kb rescuing fragment and approximately 3 kb downstream was first cloned into the pCaSpeR2 vector (Thummel and Pirrotta, 1988). The *ovo<sup>D1</sup>* cosmid DNA was partially digested with *Bg*III followed by complete digestion with *Eco*RI to generate the expected 10.5 kb fragment. The ligation reaction was done in Nuseive-agarose gel (Sheen, 1993) containing both the 10.5 kb *Bg*III-*Eco*RI insert and the pCaSpeR2 vector digested with *Bg*III and *Eco*RI. This *ovo<sup>D1</sup>* subclone, pD1B2R, was partially digested with *Not*I and end-filled with the Klenow fragment of DNA polymerase. The resulting clone, pD1B2R#225, has no *Not*I site in the pCaSpeR2 vector sequences. A 12 kb *Not*I-*Bg*III DNA fragment isolated from *Bg*III partial and *Not*I complete digestion of *ovo<sup>D1</sup>* cosmid DNA, Cos-P[*ovo<sup>D1</sup>*]<sup>S23-1</sup>, was cloned into *Not*I and *Bg*III

digested pD1B2R#225 vector. The resulting 18 kb *ovo<sup>D1</sup>* clone, pD1B2NR, contains 8kb upstream of the 7.2 kb *SalI-HindIII* region and approximately 3kb downstream. Another 7.2 kb subclone, pD1B2H, containing only the 7.2 kb *ovo<sup>D1</sup> SalI-HindIII* counterpart was cloned. The *ovo<sup>D1</sup>* cosmid DNA was partially digested with *HindIII* and completely digested with *BglIII*. The purified 7.2 kb fragment was introduced into pCaSperR2 vector partially digested with *HindIII* and completely digested with *BglIII*. Basic molecular biology manipulation were done as described in Sambrook et al. (1989).

### P-element transformation

DNA constructs used for microinjection were purified through an EtBr-CsCl gradient. The EtBr-DNA band was extracted four times with 1:1 isopropanol/Tris-EDTA solution saturated with NaCl to remove EtBr. DNA was then precipitated by EtOH and re-suspended in TE. Further extraction was performed at least three times with phenol/chloroform. The purified DNA was then re-precipitated with EtOH and used for microinjection. Modified microinjection was done as described by Chou et al. (1987). The host strains used for injection were *y w* or *y w; 2-3, Sb/TM6* (Robertson et al., 1988). Both transposase helper plasmids, p 25.7wc (Karess and Rubin, 1984) and p 2-3 (Laski et al., 1986), were used in different injection experiments. Transformants were identified by rescue of the white eye color.

### Transposase stocks

The 'jumpstarter' strain used is *2-3, P[ry<sup>+</sup>; 2-3]*, which carries a defective P-element on the third chromosome at 99B which constitutively expresses high levels of transposase but can not itself transpose (Robertson et al., 1988). The stocks, *ry<sup>506</sup>, 2-3* and *CyO/Sp; ry<sup>506</sup>, 2-3, Sb/TM6*, that provide a source of transposase activity were obtained from the Bowling Green Stock Center. The stock carrying *2-3* on the *TM3* balancer chromosome (*TM3, Sb ry 2-3/Df(3R)C7,ry<sup>506</sup>*) was obtained from G. Reuter and J. Szabad.

Flies were raised on standard *Drosophila* medium at 25°C. Descriptions of balancers and mutations that are not described in the text can be found in Lindsley and Zimm (1992).

### Autosomal *P[ovo<sup>D1</sup>]* stocks

Autosomal *P[ovo<sup>D1</sup>]* dominant female mutations are kept as stocks using dominant male-sterile mutations. Second chromosome *P[ovo<sup>D1</sup>]* insertions are kept as *w/Y; P[ovo<sup>D1</sup>]/CyO* males crossed with *S Sp Ms(2)M bw<sup>D</sup>/CyO* females. Third chromosome *P[ovo<sup>D1</sup>]* insertions are kept as *w/Y; P[ovo<sup>D1</sup>]/TM3, Sb* males crossed with *ru h st B2<sup>D</sup> ss e<sup>s</sup>/TM3, Sb* females. The dominant male-sterile stocks *Ms(2)M* and *B2<sup>D</sup>* were obtained from D. Lindsley and M. Fuller, respectively (D. Lindsley, personal communication; Kempthues et al., 1980).

Concurrently, *P[ovo<sup>D1</sup>]* dominant female insertions are also kept as *w/Y; P[ovo<sup>D1</sup>]/CyO* and *w/Y; P[ovo<sup>D1</sup>]/Sco* males crossed with *w/w; CyO/Sco* females (for the second chromosome), and *w/Y; P[ovo<sup>D1</sup>]/TM3, Sb* and *w/Y; P[ovo<sup>D1</sup>]/Cx<sup>D</sup>* males crossed with *w/w; TM3, Sb/Cx<sup>D</sup>* females (for the third chromosome).

### In situ hybridization

In situ hybridization to polytene chromosomes was performed as described in Chou and Perrimon (1992). The probe used is the pP[>w<sup>hs</sup>>] plasmid containing FRT sequences, the *mini-w* (*w<sup>hs</sup>*) gene and the P-element terminal inverted repeat sequences (Golic and Lindquist, 1989).

### Egg-laying ability test

10-20 females, heterozygous for a *P[ovo<sup>D1</sup>]* insertion, were mated to five males in a vial for at least 6 days and examined for their egg-laying ability. When eggs were found in a vial, the number of

females laying eggs was determined by looking for the presence of mature egg chambers either by visual examination of the abdomen or following ovary dissection.

### Induction of germ-line clones

To induce germ-line clones, first instar larvae (24 to 48 hours old) from the appropriate cross were irradiated at a constant dose of 1000 rads (Torrex 120D X-ray machine; 100 kV, 5 mA, 3 mm aluminum filter) as described in Perrimon et al. (1984).

### Examination of embryonic phenotypes

Females that carry germ-line clones were detected by either examining their egg-laying ability or by the presence of developed ovaries. Eggs were scored as unfertilized if no embryonic development was detectable following dechorionation in 50% bleach. Cuticles were prepared in Hoyers' mountant as previously described (van der Meer, 1977). Embryos were examined by both phase-contrast and dark-field microscopy.

### Mutations used for germ-line clonal analysis


*rho<sup>P 5</sup>/TM6B* and *rho<sup>P 38</sup>/TM6B* (Freeman et al., 1992), *y w; rho<sup>TM43</sup> e/TM6B, Tb* (Mayer and Nusslein-Volhard, 1988), *sev<sup>E4</sup>; Sos<sup>X122</sup>/SM6a* (Rogge et al., 1991), *st e Tr<sup>444</sup>/TM3* (Anderson et al., 1985), *Gap1<sup>B1</sup>/TM6* and *w; Gap1<sup>B2</sup>/TM3, Sb* (Gaul et al., 1992), and *w; mip<sup>w+</sup>/TM6C* (Buckles et al., 1992) were used for germ-line clonal analyses.

## RESULTS

### Recovery of *P[ovo<sup>D1</sup>]* transformants

The *ovo* gene is located at cytological position 4E1-2. Genetic analysis of this region has shown that *ovo* may be part of a gene complex that includes the *shaven-baby* (*svb*) gene. Many *ovo<sup>D1</sup>* revertants are also mutant at the *svb* locus, which maps less than 0.5 map unit from *ovo*, suggesting that both genes may share common regulatory or coding sequences (Oliver et al., 1988; Mevel-Ninio et al., 1991; Garfinkel et al., 1992). Since the nature of the *ovo<sup>D1</sup>* mutation, as well as the regulatory regions of *ovo*, have not yet been defined, we decided to generate a P-element transposon that contains a fragment of *ovo<sup>D1</sup>* genomic DNA as large as possible for germ-line transformation. Since recovery of transformants using a Cos-P vector that contains close to 40 kb of DNA has been previously described (Haenlin et al., 1985), we attempted to introduce a Cos-P[*ovo<sup>D1</sup>*] into the fly genome. A cosmid, Cos-P[*ovo<sup>D1</sup>*]<sup>S23-1</sup>, was isolated as the result of a direct PCR screen (Materials and Methods). This cosmid encompasses the *ovo<sup>D1</sup>* counterpart of the *ovo<sup>+</sup>* 7.2 kb fragment, which can rescue the female sterility phenotype of *ovo* mutation (Mével-Ninio et al., 1991) along with approximately 20 kb of upstream and 10 kb of downstream DNA (Fig. 1). 6,000 embryos were injected with Cos-P[*ovo<sup>D1</sup>*]<sup>S23-1</sup>, but no transformants were recovered (Table 1).

Three subclones of the *ovo<sup>D1</sup>* gene, pD1B2NR, pD1B2R and pD1B2H (Fig. 1), were subsequently derived from Cos-P[*ovo<sup>D1</sup>*]<sup>S23-1</sup> and used for P-element-mediated transformation. All of these plasmids were associated with high toxicity as shown by the poor recovery of larvae and adults from the injected embryos (see Table 1). We were not able to recover transformants of pD1B2R and pD1B2H (Table 1); however, three independent G<sub>0</sub> males derived from pD1B2NR injec-



**Fig. 1.** Cloning of *ovo<sup>D1</sup>*. The position of the *ovo<sup>D1</sup>* genomic fragments used in the P-element transformation experiments are shown with respect to the extent of the *ovo* 7.2 kb rescuing fragment. This fragment contains the proposed 5 kb *ovo<sup>+</sup>* mRNA transcription unit and rescues the female sterility phenotype of *ovo<sup>-</sup>* mutations (Mével-Ninio et al., 1991). The 7.2 kb *ovo* fragment and the position of the four PCR products used for screening the cosmid library are indicated. Only the restriction sites used in the plasmid subcloning (see Material and Methods) are shown. For a more detailed map see Mével-Ninio et al. (1991) and Garfinkel et al. (1992). The zero

point demarcates the *SalI* site as described in Mével-Ninio et al. (1991). The locations of *SalI*, *BglII*, *NotI*, *HindIII*, *EcoRI* and *PstI* restriction sites in the Cos-P[*ovo<sup>D1</sup>*]<sup>S23-1</sup> clone are similar to those reported previously (Mével-Ninio et al., 1991; Garfinkel et al., 1992). The Cos-P[*ovo<sup>D1</sup>*]<sup>S23-1</sup> DNA contains 20 kb of DNA upstream of and 18 kb of DNA downstream of the *SalI* site. The extent of the three subclones derived from Cos-P[*ovo<sup>D1</sup>*]<sup>S23-1</sup> is shown.

**Table 1. Recovery of *P[ovo<sup>D1</sup>]* transformants**

Construct (insert size)	Strain/helper	Number injected		Number recovered	
		E	L	G <sub>0</sub>	G <sub>1</sub>
Control (5 kb)	y w/pΔ2-3	150	50	24	5
S23-1 (38 kb)	y w/pΔ2-3	4,971	1,436	385	0
	Δ2-3	1,098	245	24	0
pD1B2NR (18 kb)	Δ2-3	2,354	281	21	2 <i>P[ovo<sup>D1</sup>]<sup>1,3</sup></i>
	y w/pπ25.1wc	1,891	376	158	1 <i>P[ovo<sup>D1</sup>]<sup>2</sup></i>
	Δ2-3*	4,976	364	123	0
pD1B2R (10.5 kb)	Δ2-3	510	75	7	0
pD1B2H (7.2 kb)	y w/pΔ2-3	407	19	7	0
	Δ2-3	2,119	37	3	0

Microinjections were performed as described in Chou et al. (1987). In control experiments, using a 5 kb plasmid at a concentration of 200 μg/ml, 30% of the injected embryos (E) hatch and 50% of the larvae (L) give rise to fertile G<sub>0</sub> adults. 20% of the G<sub>0</sub> adults are usually transformed. We found that *ovo<sup>D1</sup>* plasmid subclones are toxic to injected embryos especially when injected into the Δ2-3 strain (y w; Δ2-3, *Sb/TM6*) or using the pΔ2-3 helper injected into a y w strain (Laski et al., 1986). The toxicity is apparent from the low hatching rate (8 to 12%) and from the low eclosion rate (less than 10% of the hatched larvae). The toxicity effect was lower when the germ-line-specific helper, pπ25.7wc, was used (Karess and Rubin, 1984).

DNA concentrations for injection are: 400-500 μg/ml for both the S23-1 cosmid and pD1B2NR subclone, 100-200 μg/ml for both pD1B2R and pD1B2H subclones, and 100-200 μg/ml for the transposase helpers in the final injection solution.

\*In this experiment, 30 μg/ml of pD1B2NR was injected.

The number of G<sub>0</sub> adults includes both males and females. Lines were not established from the G<sub>0</sub> females since the presence of the *ovo<sup>D1</sup>* mutation in the germ line of G<sub>0</sub> females is expected to prevent egg formation.

tions led to some G<sub>1</sub> w<sup>+</sup> transformed flies. The first G<sub>0</sub> males produced 10 w<sup>+</sup> females without any w<sup>+</sup> males, suggesting that the insertion was X-linked. Six of these females were not able to lay eggs, while others laid eggs and gave rise to adult progeny. The line *P[ovo<sup>D1</sup>]<sup>1</sup>* established from this X-linked insertion contained multiple *P[ovo<sup>D1</sup>]* insertions and chromosomal rearrangements and could not be used in subsequent experiments. From the second G<sub>0</sub> male, we were

able to establish successfully the line *P[ovo<sup>D1</sup>]<sup>2</sup>* that was associated with dominant female sterility (see below). *P[ovo<sup>D1</sup>]<sup>2</sup>* segregates with the third chromosome and localized to 70B-C on 3L (Table 2). No line was derived from the third G<sub>0</sub> male, *P[ovo<sup>D1</sup>]<sup>3</sup>*, since it produced only a single w<sup>+</sup> female that did not lay any eggs.

### Egg-laying ability of *P[ovo<sup>D1</sup>]<sup>2</sup>*

Most females (94%) that carry one copy of the *P[ovo<sup>D1</sup>]<sup>2</sup>* transposon have atrophic ovaries, a phenotype similar to females heterozygous for *ovo<sup>D1</sup>*. However, the remaining 6% females lay some abnormal eggs which never hatch (Table 3). These eggs have fused dorsal appendages and are usually flaccid, a phenotype also associated with two dominant alleles at the *ovo* locus, *ovo<sup>D2</sup>* and *ovo<sup>D3</sup>* (Busson et al., 1983; Perrimon, 1984). *ovo<sup>D2</sup>/+* females lay very few eggs which are always flaccid, while *ovo<sup>D3</sup>/+* females lay a lot of eggs with fused filaments which usually become flaccid. Both *ovo<sup>D2</sup>* and *ovo<sup>D3</sup>* behave as antimorphic mutations. First, their female sterility can be suppressed by the presence of two copies of the *ovo<sup>+</sup>* gene; and, second, both mutations in *trans* with a loss of function *ovo<sup>-</sup>* mutation have atrophic ovaries, a phenotype also found in homozygous *ovo<sup>-</sup>* females (Busson et al., 1983; Oliver et al., 1987). The nature of the *ovo<sup>D1</sup>* mutation is less straightforward. *ovo<sup>D1</sup>* females that carry up to three copies of the *ovo<sup>+</sup>* gene are sterile; however, their atrophic ovaries are larger than those of *ovo<sup>D1</sup>* females that carry 1 or 2 copies of the *ovo<sup>+</sup>* gene. These dosage experiments suggest that *ovo<sup>D1</sup>*, like *ovo<sup>D2</sup>* and *ovo<sup>D3</sup>*, is an antimorphic mutation (Busson et al., 1983).

Since females that carry the autosomal *P[ovo<sup>D1</sup>]<sup>2</sup>* insertion carry two X-linked copies of the *ovo<sup>+</sup>* gene, we reasoned that the incomplete sterility of *P[ovo<sup>D1</sup>]<sup>2</sup>* may be due to a lower expressivity of the transposed *P[ovo<sup>D1</sup>]<sup>2</sup>* gene. A reduced amount of toxic *ovo<sup>D1</sup>* product may lead to a weaker DFS phenotype reminiscent of the sterility phenotypes associated with both the *ovo<sup>D2</sup>* and *ovo<sup>D3</sup>* mutations. To test this hypothesis, we generated flies that carry the autosomal *P[ovo<sup>D1</sup>]<sup>2</sup>* insertion in the presence of a single copy of the *ovo<sup>+</sup>* gene. This was achieved by using the loss-

**Table 2. Recovery of autosomal *P[ovo<sup>D1</sup>]* insertions**

Original <i>P[ovo<sup>D1</sup>]</i>	Chromosomal location	New <i>P[ovo<sup>D1</sup>]</i> insertions			<i>P[ovo<sup>D1</sup>]</i> selected
		Chrm.	N	N hyb.	
<i>P[ovo<sup>D1</sup>]<sup>2</sup></i>	3L; 70B-C	II	12	6	<i>P[ovo<sup>D1</sup>]<sup>13</sup></i> (2L; 27E-28A) <i>P[ovo<sup>D1</sup>]<sup>32</sup></i> (2R; 55D-E) <i>P[ovo<sup>D1</sup>]<sup>48</sup></i> (3L; 70A-E)
		III	>80	8	
		X	5	0	
<i>P[ovo<sup>D1</sup>]<sup>59</sup></i>	X	II, III	0	0	
<i>P[ovo<sup>D1</sup>]<sup>96</sup></i>	X	II, III	0	0	
<i>P[ovo<sup>D1</sup>]<sup>116</sup></i>	X	II, III	0	0	
<i>P[ovo<sup>D1</sup>]<sup>13</sup></i>	2L; 27E-28A	II	21	1	<i>P[ovo<sup>D1</sup>]<sup>13X13</sup></i> (2L; 28A and 30D) <i>P[ovo<sup>D1</sup>]<sup>13X6</sup></i>
		III	1	0	
		II(CyO)	1	0	
<i>P[ovo<sup>D1</sup>]<sup>32</sup></i>	2R; 55D-E	II	18	1	<i>P[ovo<sup>D1</sup>]<sup>131</sup></i> <i>P[ovo<sup>D1</sup>]<sup>32X9</sup></i> (2R; 55B-D) <i>P[ovo<sup>D1</sup>]<sup>32X8</sup></i>
		III	0	0	
		II	13	0	
<i>P[ovo<sup>D1</sup>]<sup>131</sup></i>	CyO	III	6	3	<i>P[ovo<sup>D1</sup>]<sup>C13</sup></i> (3R; 98A-B) <i>P[ovo<sup>D1</sup>]<sup>C13X3</sup></i> (3R; ND) <i>P[ovo<sup>D1</sup>]<sup>C13X8</sup></i>
		III	6	3	
		III	>25	0	

The number of novel *P[ovo<sup>D1</sup>]* insertions (N), their chromosomal segregation (Chrm.) and the number mapped to polytene chromosomes (N hyb), generated during the course of the various transposition experiments (see Fig. 2) is indicated. See Fig. 2 and text for further details. ND, Not determined.

**Table 3. Egg-laying abilities associated with *P[ovo<sup>D1</sup>]* autosomal insertions**

Insertions	+/+; <i>D1</i> /+	+/+; <i>D1</i> / <i>Bal</i>	Total	<i>SI</i> /+; <i>D1</i> /+	<i>SI</i> /+; <i>D1</i> / <i>Bal</i>	Total
<b>2L</b>						
<i>P[ovo<sup>D1</sup>]<sup>13</sup></i>	3/124	6/106	9/230	0/359	2/299	2/658
<i>P[ovo<sup>D1</sup>]<sup>13X6</sup></i>	0/667	0/225	0/892			
<i>P[ovo<sup>D1</sup>]<sup>13X13</sup></i>	1/374	0/263	1/637			
<b>2R</b>						
<i>P[ovo<sup>D1</sup>]<sup>32</sup></i>	12/199	8/186	20/385	0/320	1/393	1/713
<i>P[ovo<sup>D1</sup>]<sup>32X8</sup></i>	1/566	0/381	1/947			
<i>P[ovo<sup>D1</sup>]<sup>32X9</sup></i>	0/458	0/345	0/803			
<b>3L</b>						
<i>P[ovo<sup>D1</sup>]<sup>2</sup></i>	6/103	6/103		1/310	1/112	2/422
<i>P[ovo<sup>D1</sup>]<sup>2X48</sup></i>	0/31	0/1576	0/1607	0/362	0/362	
<b>3R</b>						
<i>P[ovo<sup>D1</sup>]<sup>C13</sup></i>			5/295*			
<i>P[ovo<sup>D1</sup>]<sup>C13X3</sup></i>	0/822	0/325	0/1197			
<i>P[ovo<sup>D1</sup>]<sup>C13X8</sup></i>	2/593	0/433	2/1026			

The egg-laying ability of females that carry *P[ovo<sup>D1</sup>]* insertions in various mutant backgrounds is indicated. Female sterility is represented by the number of females that lay eggs/the total number of females tested. *D1* represents the *P[ovo<sup>D1</sup>]* insertion used, *SI* is the loss-of-function *ovo<sup>SI</sup>* allele and *Bal* represents the balancer chromosomes (*CyO* for the second and *TM3, Sb* for the third).

A '+/+; *D1*/+' genotype corresponds to a female carrying a *P[ovo<sup>D1</sup>]* insertion in the presence of two wild-type copies of the *ovo* gene on the X-chromosome. Similarly, a '*SI*/+; *D1*/+' genotype corresponds to a female carrying a *P[ovo<sup>D1</sup>]* insertion in the presence of a single wild-type copy of *ovo*. Females heterozygous for a *P[ovo<sup>D1</sup>]* insertion sometimes lay eggs due to the occurrence of spontaneous mitotic recombination events in their germ line. These events occur at the frequency of 0.1-1% (Wieschaus and Szabad, 1979; Perrimon and Gans, 1983) and can be prevented by the presence of an appropriate balancer chromosome (*Bal*). The egg-laying ability of some *P[ovo<sup>D1</sup>]/+* females that carry a fully expressed insertion can be accounted for by these events.


Flies that carry a *P[ovo<sup>D1</sup>]* insertion are perfectly viable and morphologically normal. Leaky *P[ovo<sup>D1</sup>]* females can have either one or two ovaries which can range from a few ovarioles with vitellogenic oocytes to a fully developed ovary. The morphology of the eggs laid by *P[ovo<sup>D1</sup>]* females can be wild-type, flaccid or with fused dorsal appendages. These eggs can in some cases produce normal larvae and adults.

\*In this experiment we did not determine the genotype of the females that produced eggs.

of-function mutation, *ovo<sup>SI</sup>*, which behaves as a null *ovo*-mutation (Oliver et al., 1987). *ovo<sup>SI</sup> /+; P[ovo<sup>D1</sup>]<sup>2</sup>/+* females had atrophic ovaries and significantly reduced egg-laying ability (Table 3), indicating that +/+; *P[ovo<sup>D1</sup>]<sup>2</sup>/+* females laid eggs due to incomplete expressivity of the transposed *ovo<sup>D1</sup>* mutation. This result is consistent with the proposed antimorphic nature of the *ovo<sup>D1</sup>* mutation (Busson, et al., 1983; Perrimon, 1984).

### New *P[ovo<sup>D1</sup>]* insertions are associated with incomplete expressivity

P-element transposons can be mobilized using the 2-3 transposase (Robertson et al., 1988). These transposition events can be followed using the *w<sup>+</sup>* marker carried by the transposon. Flies heterozygous for the *P[ovo<sup>D1</sup>]<sup>2</sup>* insertion have pale-orange eyes. To recover additional *P[ovo<sup>D1</sup>]* autosomal insertions, we first mobilized the original



**Fig. 2.** Recovery of autosomal *P[ovoD1]* insertions. Round 1 transpositions: beginning with *P[ovoD1]<sup>j2</sup>*, as an ammunition chromosome, 500 individual crosses between one *y w/Y*; *P[ovoD1]<sup>j2</sup>/Δ2-3, Sb* male and three *C(1)DX, y f/Y* females were set up. 44 new independent jumps were recovered as *y w/Y* males with the *Sb* marker and *w<sup>+</sup>* eye color. They are on the X, 2nd and Δ2-3, *Sb* chromosomes. Tests for egg-laying ability of all 44 new insertions showed that none of them fully expressed the *ovoD1* DFS phenotype since approximately 5%-100% of females heterozygous for any specific *P[ovoD1]* insertion laid eggs. Six of the tighter insertions that segregated with the second chromosome were localized by in situ hybridization (Table 3). Among these, we selected two of them, *P[ovoD1]<sup>j13</sup>* (27E-28A) and *P[ovoD1]<sup>j32</sup>* (55D-E), for further analysis. More than 80 lines were also established from transposition of *P[ovoD1]<sup>j2</sup>*. These *y w/Y* males without the *Sb* marker showed eye colors that are significantly darker than the pale-orange eye color of *P[ovoD1]<sup>j2</sup>*. These lines segregated with the original *P[ovoD1]<sup>j2</sup>* third chromosome and eight of them were localized by in situ hybridization, four of which were found to have an insertion at 70A-E, which is very close to the position of *P[ovoD1]<sup>j2</sup>*. One of these, *P[ovoD1]<sup>j2x48</sup>*, showed a fully expressed DFS phenotype (Table 3). From this transposition, 5 X-linked insertions were also recovered. Two of them did not express the *ovoD1* DFS phenotype. Further efforts to destabilize the three insertions, *P[ovoD1]<sup>j59</sup>*, *P[ovoD1]<sup>j96</sup>* and *P[ovoD1]<sup>j116</sup>* showed that these X-linked insertions could not be mobilized.

Round 2 transpositions: the *P[ovoD1]<sup>j13</sup>* and *P[ovoD1]<sup>j32</sup>* insertions were used as ammunition to recover *P[ovoD1]* insertions on 3R. Individual crosses of one *+/Y*; *P[ovoD1]<sup>j13</sup> or <sup>j32</sup>/CyO; TM3, Δ2-3, Sb/+* male crossed to three *y w* females were established. From 200 *P[ovoD1]<sup>j13</sup>* dysgenic males, 4 independent jumps were recovered as *CyO* males with *w<sup>+</sup>* eye color. One of these, *P[ovoD1]<sup>j134</sup>*, segregated with the third chromosome, and another, *P[ovoD1]<sup>j131</sup>*, with the *CyO* chromosome. From 70 *P[ovoD1]<sup>j32</sup>* dysgenic males tested, no transposition events to other chromosomes could be recovered. Taking advantage of the previous observation that the fully expressed *ovoD1* DFS phenotype could be

recovered among flies with darker eye colors, we selected insertions associated with darker eye colors that segregated with the *P[ovoD1]<sup>j13</sup>* or *P[ovoD1]<sup>j32</sup>* chromosomes. From the above transposition of the *P[ovoD1]<sup>j13</sup>* element, 21 lines were recovered with dark eye colors. Two of them, *P[ovoD1]<sup>j13x6</sup>* and *P[ovoD1]<sup>j13x13</sup>*, showed the fully expressed *ovoD1* DFS phenotype. In situ hybridization detected two signals on the *P[ovoD1]<sup>j13x13</sup>* chromosome, one at 28A, which most likely corresponds to the original site of *P[ovoD1]<sup>j32</sup>*, and one at 30D. This result suggests that the tight *ovoD1* DFS phenotype of *P[ovoD1]<sup>j13x13</sup>* is the result of two *P[ovoD1]* insertions. Due to the low frequency of *P[ovoD1]<sup>j32</sup>* transposition, an additional large scale mobilization experiment was performed to recover flies with darker eye color insertions. Of 18 lines that were recovered, two of them, *P[ovoD1]<sup>j32x8</sup>* and *P[ovoD1]<sup>j32x9</sup>*, were associated with a fully expressed *ovoD1* DFS phenotype (Table 3). *P[ovoD1]<sup>j32x9</sup>* localized to the same position as that of *P[ovoD1]<sup>j32</sup>* suggesting that it is the result of a local transposition.

Round 3 transpositions: to recover additional insertions on the third chromosome, we mobilized *P[ovoD1]<sup>j131</sup>* on the *CyO* balancer chromosome. From the large scale cross between *CyO, P[ovoD1]<sup>j131</sup>/+; Δ2-3, Sb/+* males with *y w* females, new transpositions were recovered among the progeny as males that segregate the *w<sup>+</sup>* eye color from the *CyO* chromosome. From 28 lines recovered, 6 segregated with the third chromosome and expressed the DFS *ovoD1* phenotype. Germ-line clonal analysis of *T<sup>tr444</sup>* was performed using 5 of these 6 insertions to determine on which arm they resided. In situ hybridization was performed for 3 of these 6 insertions to determine their locations. One of them, *P[ovoD1]<sup>jC13</sup>*, mapped to position 98A-B and was selected for further analyses since it is associated with an expressed *P[ovoD1]* DFS phenotype (1.7% leakiness; Table 3).

Round 4 transpositions: a final set of transpositions was undertaken to recover derivatives of *P[ovoD1]<sup>jC13</sup>* that fully express the DFS *ovoD1* phenotype. From large scale crosses between *y w/Y; P[ovoD1]<sup>jC13</sup>/Δ2-3, Sb* males and *y w* females, more than 25 lines of *Sb<sup>+</sup>* males with darker eye colors were recovered. Two of these lines, *P[ovoD1]<sup>jC13x3</sup>* and *P[ovoD1]<sup>jC13x8</sup>*, show the fully expressed *ovoD1* DFS phenotype (Table 3).

**Table 4. Frequency of females with germ-line clones generated using the autosomal DFS technique**

Mutation	Location	$P[ovo^{D1}]$	#Females	#GLC	Percentage
<i>Toll</i> <sup>r444</sup>	97D	<i>C13</i> *	ND	5	ND
<i>Sos</i> <sup>X122</sup>	34D	<i>13</i> †	1060	12	1.13
<i>rho</i> <sup>P 5</sup>	62A	<i>2X48</i> ‡	1302	19	1.46
<i>rho</i> <sup>P 38</sup>			510	11	2.16
<i>rho</i> <sup>7m43</sup>			850	11	1.29
<i>Gap1</i> <sup>B2</sup>	67D	<i>2X48</i>	368	6	1.63
<i>Gap1</i> <sup>mipw+</sup>			799	12	1.5

#Females: number of *mutant/P[ovo<sup>D1</sup>]* females examined.

#GLC: number of *mutant/P[ovo<sup>D1</sup>]* females with germ-line clones.

Percentage : [ #GLC / #Female ] × 100.

ND : the number of females was not determined in this experiment.

\*Germ-line clonal analysis of *Toll<sup>r444</sup>* was performed by using *P[ovo<sup>D1</sup>]<sup>C13</sup>* without introducing the *ovo<sup>S1</sup>* allele on the X-chromosome. Eggs not derived from homozygous germ-line clones have two fused dorsal appendages (Table 3).

†*P[ovo<sup>D1</sup>]<sup>13</sup>* was used as the DFS mutation for the analysis of *Sos<sup>X122</sup>* and *ovo<sup>S1</sup>* was introduced onto the X chromosome to ensure full expressivity of *P[ovo<sup>D1</sup>]*. The germ-line clone phenotype of *Sos<sup>X122</sup>* was described by Lu et al. (1993).

‡*P[ovo<sup>D1</sup>]<sup>2X48</sup>* shows a fully expressed *ovo<sup>D1</sup>* DFS phenotype (Table 2). It was used directly as a DFS without introducing the *ovo<sup>S1</sup>* mutation on the X-chromosome. The *rho<sup>P 5</sup>* germ-line clone phenotype was described by Ruohola-Baker et al. (1993). Like *rho<sup>P 5</sup>*, neither *rho<sup>P 38</sup>* nor *rho<sup>7M43</sup>* function is required in the germ line for oogenesis (Ruohola-Baker et al. unpublished data).

We also used fully expressed *P[ovo<sup>D1</sup>]* insertions on 2L and 2R to analyze the maternal effects of different *Enhancer of sevenless* mutations (Simon et al., 1991). *P[ovo<sup>D1</sup>]<sup>13X13</sup>* was used in the analysis of various *Sos* alleles and *P[ovo<sup>D1</sup>]<sup>32X9</sup>* for the analysis of *drk<sup>e0A</sup>*. In both cases these *P[ovo<sup>D1</sup>]* insertions behaved as expected (data not shown).

*P[ovo<sup>D1</sup>]<sup>2</sup>* insertion. New transpositions were recovered on the X, 2nd and 3rd chromosomes (Fig. 2; Table 2).

We focused our efforts on the 12 new insertions that segregated with the 2nd chromosome. All of these insertions showed varying degrees of egg-laying ability. In no case did we find a 2nd chromosome *P[ovo<sup>D1</sup>]* insertion that was associated with a fully expressed *ovo<sup>D1</sup>* DFS phenotype. Among these, we selected those associated with a more severe DFS sterility. We mapped six of them by in situ hybridization and further selected two of them, *P[ovo<sup>D1</sup>]<sup>13</sup>* located at position 27E-28A on the 2L arm, and *P[ovo<sup>D1</sup>]<sup>32</sup>* located at 55D-E on the 2R arm, for further transpositions (Table 2). Approximately 5% of females heterozygous for these two *P[ovo<sup>D1</sup>]* insertions display a leaky *ovo<sup>D1</sup>* DFS phenotype and lay some eggs. In the presence of only one copy of the wild-type *ovo* gene, the leakiness of *P[ovo<sup>D1</sup>]<sup>13</sup>* and *P[ovo<sup>D1</sup>]<sup>32</sup>* is reduced significantly (Table 3).

### Isolation of insertions that exhibit a fully expressed *ovo<sup>D1</sup>* DFS phenotype

In addition to the new insertions described above, we also recovered in the same experiment a large number of flies associated with darker eye colors, which segregated with the original *P[ovo<sup>D1</sup>]<sup>2</sup>* chromosome. A darker eye color can be recovered when multiple copies of the transposon are present or when the P-element is inserted into some chromosomal regions that provide high expression of the transgene. More than 80 of these insertions segregated with the original *P[ovo<sup>D1</sup>]<sup>2</sup>* third chromosome. The female sterility of 10 of these lines was examined in detail and one

of them *P[ovo<sup>D1</sup>]<sup>2X48</sup>* was found to exhibit a complete DFS sterility similar to that of females heterozygous for *ovo<sup>D1</sup>* (Table 3). Therefore, it is possible that the *P[ovo<sup>D1</sup>]<sup>2X48</sup>* line carries either a single insertion in a chromosomal site that increases the expression of the inserted gene or multiple insertions that will increase the amount of the *ovo<sup>D1</sup>* gene product. We performed in situ hybridization to determine whether this chromosome contains multiple insertions. A single site was detected at 70B-C, a position similar to the location of *P[ovo<sup>D1</sup>]<sup>2</sup>*, therefore not allowing us to distinguish between these two possibilities (see Discussion).

### Recovery of other autosomal *P[ovo<sup>D1</sup>]* insertions

Transposition schemes (Fig. 2) similar to the one previously described for *P[ovo<sup>D1</sup>]<sup>2</sup>* transposition were used to recover insertions on the 3R arm. Because insertions on the X-chromosome did not show any sign of transposition, we used *P[ovo<sup>D1</sup>]<sup>13</sup>* and *P[ovo<sup>D1</sup>]<sup>32</sup>* on the 2nd chromosome to start a series of transpositions. One insertion, *P[ovo<sup>D1</sup>]<sup>C13</sup>*, that localized to position 98A-B was selected as the 3R *P[ovo<sup>D1</sup>]*. As observed with the 2nd chromosomal insertions, females that carry this insertion sometimes laid eggs (Table 3).


To recover fully expressed *P[ovo<sup>D1</sup>]* insertions located on 2L, 2R and 3R, we selected transposition events that were associated with darker eye colors. As expected from the experiments that led to the recovery of the 3L *P[ovo<sup>D1</sup>]<sup>2X48</sup>* insertion, these screens allowed us to recover insertions that were associated with a fully expressed *ovo<sup>D1</sup>* DFS phenotype: *P[ovo<sup>D1</sup>]<sup>13X6</sup>* and *P[ovo<sup>D1</sup>]<sup>13X13</sup>* on 2L, *P[ovo<sup>D1</sup>]<sup>32X8</sup>* and *P[ovo<sup>D1</sup>]<sup>32X9</sup>* on 2R, and *P[ovo<sup>D1</sup>]<sup>C13X3</sup>* and *P[ovo<sup>D1</sup>]<sup>C13X8</sup>* on 3R. Females heterozygous for these insertions do not lay eggs (Table 3).

In conclusion, through a series of transpositions, starting with *P[ovo<sup>D1</sup>]<sup>2</sup>*, we have recovered *P[ovo<sup>D1</sup>]* insertions on each autosomal arm which are associated with fully expressed DFS phenotypes that are similar to the original *ovo<sup>D1</sup>* X-linked mutation. We hence designate *P[ovo<sup>D1</sup>]<sup>13X13</sup>* as *P[ovo<sup>D1</sup>]<sup>2L</sup>*, *P[ovo<sup>D1</sup>]<sup>32X9</sup>* as *P[ovo<sup>D1</sup>]<sup>2R</sup>*, *P[ovo<sup>D1</sup>]<sup>2X48</sup>* as *P[ovo<sup>D1</sup>]<sup>3L</sup>*, and *P[ovo<sup>D1</sup>]<sup>C13X3</sup>* as *P[ovo<sup>D1</sup>]<sup>3R</sup>*.

### The germ-line dependence of autosomal *P[ovo<sup>D1</sup>]* insertions

The sterility of *ovo<sup>D1</sup>* is a germ-line-dependent phenomenon (Busson et al., 1983; Perrimon and Gans, 1983; Perrimon, 1984), and we expected that the autosomal *P[ovo<sup>D1</sup>]* insertions that we recovered would display the same germ-line dependence. To test this, we examined the eggs derived from females of genotype *P[ovo<sup>D1</sup>]/Tl<sup>r444</sup>*, X-ray irradiated as larvae. *Tl<sup>r444</sup>* is a loss-of-function *Toll* allele, which produces dorsalized embryos as a recessive maternal effect (Anderson et al., 1985). This maternal effect is strictly due to the germ-line-dependent expression of *Tl* (Schupbach and Wieschaus, 1986). We used *P[ovo<sup>D1</sup>]<sup>C13</sup>* as a DFS mutation to generate *Tl<sup>r444</sup>* homozygous germ-line clones. As expected, fertilized eggs derived from these females were all dorsalized (data not shown).

The frequency of mosaics recovered using these *P[ovo<sup>D1</sup>]* insertions was determined following a mosaic analysis of several zygotic lethal mutations (Table 4). We found that



**Fig. 3.** Somatic dependence of *Gap1* mutations. Two alleles at the *Gap1* locus, *mip<sup>w+</sup>* and *Gap1<sup>B2</sup>* were used in this analysis. *mip<sup>w+</sup>* is denoted as *Gap1<sup>mip<sup>w+</sup></sup>* since *mip* and *Gap1* are allelic (Buckles et al., 1992; Gaul et al., 1992). While *Gap1<sup>mip<sup>w+</sup></sup>* is a strong allele of the *mip* gene (Buckles et al., 1992), *Gap1<sup>B2</sup>* is more likely to represent a complete loss-of-function mutation (Gaul et al., 1992). Both *Gap1<sup>mip<sup>w+</sup></sup>* and *Gap1<sup>B2</sup>* homozygous females are associated with semi-sterility. When crossed to wild-type males approximately 80% of the eggs do not hatch. (A) Wild-type egg. (B) Weak dorsialized egg laid by *Gap1<sup>B2</sup>* females. The basal appendage material is enlarged towards the ventral side of the egg. The degree of dorsalization of *Gap1<sup>mip<sup>w+</sup></sup>* eggs is slightly weaker than in *Gap1<sup>B2</sup>* eggs. (C) Wild-type embryo. (D) Weak partially dorsialized embryo produced by *Gap1<sup>mip<sup>w+</sup></sup>* females. Variability in the degree of dorsalization is apparent in embryos derived from *Gap1<sup>mip<sup>w+</sup></sup>* and *Gap1<sup>B2</sup>* homozygous females. In some cases, embryos were completely dorsialized (data not shown).

approximately 1-2 % of heterozygote females irradiated at the end of the first larval instar stage (see Materials and Methods) carried a germ-line clone. Such frequencies of germ-line mosaicism are similar to those reported previously using *Fs(2)D* and *Fs(2)I* (MacDonald and Struhl, 1986; Wieschaus et al., 1984; N. P. unpublished).

#### Germ-line clonal analysis of *Gap1* gene

Components of the p21<sup>ras</sup> signaling pathway have been implicated in the signaling of many receptor protein tyrosine kinases such as sevenless and Torso (reviewed in Perrimon, 1993). Both *Ras1* and *Sos*, an exchange factor encoded by *Son of sevenless*, have been shown to operate in signaling of both Torso (Lu et al., 1993; Doyle and Bishop, 1993) and Sevenless (Rogge et al., 1991; Simon et al., 1991; Bonfini et al., 1992). To determine whether *Gap-1*, which encodes a Ras-GTPase-activating protein implicated in Sevenless signaling (Gaul et al., 1992), also operates in Torso

signaling, we examined the maternal effect phenotype of two *Gap1* mutations, *Gap1<sup>mip<sup>w+</sup></sup>* (Buckles et al., 1992) and *Gap1<sup>B2</sup>* (Gaul et al., 1992). Homozygous *Gap1<sup>mip<sup>w+</sup></sup>* and *Gap1<sup>B2</sup>* females lay eggs that show a weak dorsialized egg shell morphology (Fig. 3B). The base of the appendages is expanded towards the ventral side of the egg (Fig. 3B). In addition, embryos derived from *Gap1* homozygous females are partially dorsialized (Fig. 3D). Since the weak dorsalization phenotype of both the egg and the embryo may result from *Gap1* gene function in either somatic follicle cells or germ-line cells, we decided to analyze the tissue specificity of *Gap1* mutations. If the combination of mutant germ line and wild-type soma leads to the mutant phenotype, then the function of this gene is necessary in the germ line. On the other hand, if under these conditions wild-type instead of the mutant phenotype is produced, then the mutation is due to gene function in the somatic follicle tissue. When crossed to either heterozygous *Gap1* or wild-type males, all 6 *Gap1<sup>B2</sup>*



and 11 out of 12 *Gap1<sup>mipw+</sup>* mosaic females showed no maternal effect. Eggs with wild-type chorion morphology were laid and larvae hatched. Some of the larvae died during early larval stages without obvious cuticle defects. One *Gap1<sup>mipw+</sup>* mosaic female laid a few embryos showing a maternal effect lethal phenotype, which in no case resembled dorsalized embryos (data not shown).

We conclude that the weak dorsalization of both the egg shell and embryo are not germ-line dependent and *Gap1* function in follicle cells is required for proper egg shell and embryonic dorsal-ventral polarity.

## DISCUSSION

To extend the DFS technique to the autosomes, we have cloned the *ovo<sup>D1</sup>* mutation into a P-element vector and recovered fully expressed *P[ovo<sup>D1</sup>]* insertions on each autosomal arm. Using the autosomal DFS technique, we have analyzed the germ-line clone phenotypes of several zygotic lethal mutations. This analysis indicates that the DFS sterility of *P[ovo<sup>D1</sup>]* insertions is germ-line dependent and that these *P[ovo<sup>D1</sup>]* insertions are excellent tools for generating germ-line chimeras.

### Incomplete expressivity of *P[ovo<sup>D1</sup>]* insertions

Surprisingly, the 18 kb *P[ovo<sup>D1</sup>]* insertions that contain 8 kb of upstream and 3kb of downstream DNA along with the counterpart of the 7.2 kb *ovo<sup>+</sup>* rescuing fragment showed a weak *ovo<sup>D1</sup>* DFS phenotype. We tested more than 100 lines that carried a newly-inserted *P[ovo<sup>D1</sup>]* and did not recover an insertion associated with complete expressivity. This suggests that the *ovo* gene may be larger and more complex than originally proposed (Mével-Ninio et al., 1991; Garfinkel et al., 1992). It is only following local transposition that we were able to recover *P[ovo<sup>D1</sup>]* insertions that had similar genetic properties to the X-linked *ovo<sup>D1</sup>* mutation.

We envision two possibilities for the production of tight *P[ovo<sup>D1</sup>]* insertions. First, they may have been generated by duplication of the original insertion. Precedents for such duplication events have been reported previously (Roiha et al., 1988; Salz et al., 1987; Hawley et al., 1988). Increased expressivity of the DFS phenotype is most likely achieved by expressing two copies of *P[ovo<sup>D1</sup>]* so that two copies of the wild-type X-linked *ovo* gene are fully antagonized. With the exception of *P[ovo<sup>D1</sup>]<sup>13X13</sup>*, only one *in situ* signal was detectable in all tight *P[ovo<sup>D1</sup>]* insertions examined. Recently it was reported that P-elements preferentially transpose into the genomic region very close to their original insertion sites (Tower et al., 1993; Zhang and Spradling, 1993). These observations suggest that duplication of the P-element transposon may have occurred locally in the *P[ovo<sup>D1</sup>]* lines that are associated with a darker eye color and full DFS expressivity as the result of the expression of two copies of *P[ovo<sup>D1</sup>]*. Another alternative is that following local transposition, higher levels of *P[ovo<sup>D1</sup>]* expression from the new insertion site. Further characterization of *P[ovo<sup>D1</sup>]* insertions that display either strong or weak DFS phenotypes will be needed to resolve this issue.

### *Gap1* gene is required in the follicle cells for dorsoventral polarity

During oogenesis, signaling between the oocyte and follicle cells lead to the asymmetric differentiation of the follicle cells along the dorsal-ventral axis (reviewed in Schupbach et al., 1991; Govind and Steward, 1991). Mutations that disrupt these processes alter the polarity of both the egg shell and the embryo. It has been proposed that a dorsalizing signal is sent from the germ-line vesicle/oocyte nucleus to the surrounding dorsal follicle cells and induces a dorsal cell fate (Schupbach, 1987; Manseau and Schupbach, 1989; Kelly, 1993; Ruohola-Baker et al., 1993). This signal may serve as the ligand for Torpedo, which is a *Drosophila* EGF-receptor homologue functioning in the soma (Schupbach, 1987; Price et al., 1989; Schejter and Shilo, 1989; Clifford and Schupbach, 1989). Up to now, components that might transduce signals from activated Torpedo have not yet been characterized.

Different receptor tyrosine kinases use common intracellular signaling components that include *Sos*, *Ras1*, and *drk* (Simon et al., 1993; Olivier et al., 1993; Lu et al., 1993; Doyle and Bishop, 1993; Sturtevant et al., 1993; Perrimon, 1993). Our germ-line clonal analysis of *Gap1* mutations suggests that *Gap1* acts in the establishment of dorsal/ventral polarity of the egg chamber. Based on these observations we propose that components of the *Gap1/Ras1* pathway function in transducing the signal from an activated Torpedo receptor tyrosine kinase. In this model *Gap1* loss-of-function mutations will give rise to dorsalized egg shell and embryo by increasing the level of Ras-GTP in the follicle cells.

Since the two *Gap1* mutations we tested in our germ-line clone analysis are not associated with a germ-line-dependent maternal effect, it suggests that *Gap1* is not involved in Torso signaling. This result suggests the presence of at least a second Ras-GAP *Drosophila* protein. However, a formal possibility is that the two *Gap1* mutations we tested are not complete null mutations. Further work will be required to clarify this issue.

### The autosomal FLP-DFS technique

In this paper, we have described the recovery of *P[ovo<sup>D1</sup>]* insertions that allow the easy generation of germ-line mosaics for autosomal mutations. However, the frequency of X-ray induced germ-line clone induction is low. On the X-chromosome, approximately 5% of chimeras can be recovered in females heterozygous for *ovo<sup>D1</sup>* (Perrimon, 1984). On the autosomes, a lower frequency of only 1 to 2% was achieved using *P[ovo<sup>D1</sup>]* insertions as DFS mutations.

To increase the efficiency of germ-line mosaic production, we previously developed the 'FLP-DFS' technique (Chou and Perrimon, 1992). This technique relies on the ability of the yeast FLP-recombinase to catalyze and promote mitotic exchange between chromosomes specifically at FRT (FLP-Recombinase Target) sites (Golic, 1991). In this method, the site of recombination is determined by the position of the FRT sequences and the efficiency of mitotic recombination is controlled by the heat shock inducible FLP-recombinase. We have shown that greater than 90% of germ-line mosaics can be easily recovered in

heterozygous *ovo<sup>DI</sup>* females following heat shock induction during larval stage.

The availability of the autosomal *P[ovo<sup>DI</sup>]* insertions with similar genetic properties to the *ovo<sup>DI</sup>* mutation will allow us to extend this 'FLP-DFS' technique to the autosomes. The construction of chromosomes that carry FRT elements located at the base of each chromosome with a proximal *P[ovo<sup>DI</sup>]* insertion will provide the tools necessary for large scale analyses of the maternal effects of essential genes on the autosomes.

We thank R. Cohen for providing *hsp26* BB enhancer elements, F. Laski for *P* element DNA sequences, M. McKeown for *tra* DNA sequences and A. Brand for allowing us to try Gal4-UAS binary system in the early stage of this work. We thank V. Pirrotta for pCoSpeR vector, U. Gaul, F. Katz, D. Lindsley, M. Fuller, J. Szabad and G. Reuter and the Bowling Green and Bloomington Stock Centers for essential stocks. Finally, we thank R. Binari, E. Siegfried and L. Lorenz for reading the manuscript and P. Gould for help with the figures and the manuscript. This work was supported by the Howard Hughes Medical Institute.

## REFERENCES

- Anderson, K. V., Bokla, L. and Nusslein-Volhard C. (1985). Establishment of dorsal-ventral polarity in the *Drosophila* embryo: the induction of polarity by the *Toll* gene product. *Cell* **42**, 791-798.
- Bonfini, L., Karlovich, C. A., Dasgupta, C. and Banerjee, U. (1992). The *Son of sevenless* gene product: a putative activator of Ras. *Science* **255**, 603-606.
- Buckles, G. R., Smith, Z. D. J. and Katz, F. N. (1992) *mip* causes hyperinnervation of a retinotopic map in *Drosophila* by excessive recruitment of R7 photoreceptor cells. *Neuron* **8**, 1015-1029.
- Busson, D., Gans, M., Komitopoulou, K. and Masson, M. (1983). Genetic analysis of three dominant female sterile mutations located on the X chromosome of *Drosophila melanogaster*. *Genetics* **105**, 309-325.
- Chou T.-B. and Perrimon, N. (1992). Use of a yeast site-specific recombinase to produce female germline chimeras in *Drosophila*. *Genetics* **131**, 643-653.
- Chou, T.-B., Mims, I., Belanich, M., Zachar, Z. and Bingham, P. M. (1987). Procedural improvements in injections for P-mediated germline gene transfer. *Dros. Inf. Serv.* **66**, 156-157.
- Clifford, R. T. and Schupbach, T. (1989). Coordinately and differentially mutable activities of *torpedo*, the *Drosophila melanogaster* homologue of the vertebrate EGF receptor gene. *Genetics* **123**, 771-787.
- Comer, A. R., Searles, L. L. and Kalfayan, L. J. (1992). Identification of a genomic DNA fragment containing the *Drosophila melanogaster* ovarian tumor gene (*otu*) and localization of regions governing its expression. *Gene* **118**, 171-179.
- Doyle, H. J. and Bishop, J. M. (1993). Torso, a receptor tyrosine kinase required for embryonic pattern formation, shares substrates with Sevenless and EGF-R pathways in *Drosophila*. *Genes Dev.* **7**, 633-646.
- Erdelyi, M. and Szabad, J. (1989). Isolation and characterization of dominant female sterile mutations of *Drosophila melanogaster*. I. Mutations on the third chromosome. *Genetics* **122**, 111-127.
- Frank, L. H., Cheung, H.-K., Cohen, R. S. (1992). Identification and characterization of *Drosophila* female germ line transcriptional control elements. *Development* **114**, 481-491.
- Freeman, M., Kimmel, B. E. and Rubin, G. M. (1992). Identifying targets of the rough homeobox gene of *Drosophila*: evidence that *rhomboid* functions in eye development. *Development* **116**, 335-346.
- Gans, M., Audit, C. and Masson, M. (1975). Isolation and characterization of sex-linked female sterile mutations in *Drosophila melanogaster*. *Genetics* **81**, 683-704.
- Garfinkel, M. D., Lohe, A. R. and Mahowald, A. P. (1992). Molecular genetics of the *Drosophila melanogaster* *ovo* locus, a gene required for sex determination of germline cells. *Genetics* **130**, 791-803.
- Gaul, U., Mardon, G. and Rubin, G. (1992). A putative Ras GTPase activating protein acts as a negative regulator of signaling by the sevenless receptor tyrosine kinase. *Cell* **68**, 1007-1019.
- Golic, K. and Lindquist, S. (1989). The FLP recombinase of yeast catalyzes site-specific recombination in the *Drosophila* genome. *Cell* **59**, 499-509.
- Golic, K. G. (1991). Site-specific recombination between homologous chromosomes in *Drosophila*. *Science* **252**, 958-961.
- Govind, S. and Steward, R. (1991). Dorsoventral pattern formation in *Drosophila*: Signal transduction and nuclear targeting. *Trend. Gen.* **7**, 119-125.
- Haenlin, M., Steller, H., Pirrotta, V. and Mohier, E. (1985). A 43 kilobase cosmid P transposon rescues the *fs(1)K10* morphogenetic locus and three adjacent *Drosophila* developmental mutants. *Cell* **40**, 827-837.
- Hamilton, B. A., Palazzolo, M. J. and Meyerowitz, E. M. (1991). Rapid isolation of long cDNA clones from existing libraries. *Nucleic Acids Res.* **19**, 1951-1952.
- Hawley, R. S., Steuber, R. A., Marcus, C. H., Sohn, R., Baronas, D. M., Gameron, M. L., Zitron, A. E. and Chase, J. W. (1988). Molecular analysis of an unstable P element insertion at the *singed* locus of *Drosophila melanogaster* evidence for intracistonic transposition of a P element. *Genetics* **119**, 85-94.
- Illmensee, K. (1973). The potentialities of transplanted early gastrula nuclei of *Drosophila melanogaster*. *Wilhelm Roux's Arch. Entwicklungsmech. Org.* **171**, 331-343.
- Ish-Horowitz, D. and Burke, J. F. (1981). Rapid and efficient cosmid cloning. *Nucleic Acid Res.* **9**, 2989.
- Karess, R. E. and Rubin, G. M. (1984). Analysis of P transposable element functions in *Drosophila*. *Cell* **38**, 135-146.
- Kelley, R. L. (1993). Initial organization of the *Drosophila* dorsoventral axis depends on an RNA-binding protein encoded by the *squid* gene. *Genes Dev.* **7**, 948-960.
- Kemphues, K. J., Raff, E. C., Raff, R. A. and Kaufman, T. C. (1980). Mutation in a testis-specific  $\beta$ -tubulin in *Drosophila*: analysis of its effects on meiosis and map location of the gene. *Cell* **21**, 445-451.
- King, R. C. (1970). *Ovarian development in Drosophila melanogaster*. New York: Academic Press.
- Klingler, M., Erdelyi, M., Szabad, J. and Nusslein-volhard, C. (1988). The function of *torso* in determining the termini Anlagen of the *Drosophila* embryo. *Nature* **335**, 275-277.
- Komitopoulou, K., Gans, M., Margaritis, L. M., Kafatos, F.C. and Masson, M. (1983). Isolation and characterization of sex-linked female-sterile mutations in *Drosophila melanogaster* with special attention to eggshell mutants. *Genetics* **105**, 897-920.
- Laski, F. A., Rio, D. C. and Rubin, G. M. (1986). Tissue specificity of *Drosophila* P element transposition is regulated at the level of mRNA splicing. *Cell* **44**, 7-19.
- Laski, F. A. and Rubin, G. M. (1989). Analysis of the cis-acting requirements for germ-line-specific splicing of the P-element ORF2-ORF3 intron. *Genes Dev.* **3**, 720-728.
- Lindsley, D. L. and Zimm, G. G. (1992). *The Genome of Drosophila melanogaster*. New York: Academic Press.
- Lu, X., Chou, T.-B., Williams, N. G., Roberts, T. and Perrimon, N. (1993). Control of cell fate determination by p21<sup>ras</sup>/Ras1, an essential component of *torso* signaling in *Drosophila*. *Genes Dev.* **7**, 621-632.
- Manseau, L. J. and Schupbach, T. (1989). *Cappuccino* and *spire*: two unique maternal-effect loci required for both the anteroposterior and dorsal-ventral patterns of the *Drosophila* embryo. *Genes Dev.* **3**, 1437-1452.
- MacDonald P. M. and Struhl, G. (1986). A molecular gradient in early *Drosophila* embryos and its role in specifying the body pattern. *Nature* **324**, 537-545.
- Mayer, U. and Nusslein-Volhard, C. (1988). A group of genes required for pattern formation in the ventral ectoderm of the *Drosophila* embryo. *Genes Dev.* **2**, 1496-1511.
- Mevel-Ninio, M., Terracol, R. and Kafatos, F.C. (1991). The *ovo* gene of *Drosophila* encodes a zinc finger protein required for female germ line development *EMBOJ.* **10**, 2259-2266.
- Mohler, J. D. (1977) Developmental genetics of the *Drosophila* egg. I. Identification of 50 sex-linked cistrons with maternal effects on embryonic development. *Genetics* **85**, 259-272.
- Oliver, B., Perrimon, N. and Mahowald, A.P. (1987). The *ovo* locus is required for sex-specific germ line maintenance in *Drosophila*. *Genes Dev.* **1**, 913-923.
- Oliver, B., Perrimon, N. and Mahowald, A.P. (1988). Genetic evidence

- that the sans fille locus is involved in *Drosophila* sex determination. *Genetics* **120**, 159-171.
- Olivier, J. P., Rabbe, T., Henkemeyer, M., Dickson, B., Mbamalu, G., Margolis, B., Schlessinger, J., Hafen, E. and Pawson, T.** (1993). A *Drosophila* SH2-SH3 adaptor protein implicated in coupling the Sevenless tyrosine kinase to an activator of Ras Guanine nucleotide exchange, Sos. *Cell* **73**, 179-191.
- Palmiter, R. D., Behringer, R. R., Quaife, C. J., Maxwell, F., Maxwell, I. H. and Brinster, R. L.** (1987). Cell lineage ablation in transgenic mice by cell-specific expression of a toxin gene. *Cell* **50**, 435-443.
- Perrimon, N. and Gans, M.** (1983). Clonal analysis of the tissue specificity of recessive female sterile mutations of *Drosophila melanogaster* using a dominant female sterile mutation *Fs(1)K1237*. *Dev. Biol.* **100**, 365-373.
- Perrimon, N.** (1984). Clonal analysis of dominant female sterile, germline-dependent mutations in *Drosophila melanogaster*. *Genetics* **108**, 927-939.
- Perrimon, N., Engstrom, L. and Mahowald, A.P.** (1984). The effects of zygotic lethal mutations on female germ-line functions in *Drosophila*. *Dev. Biol.* **105**, 404-414.
- Perrimon, N., Mohler, J. D., Engstrom, L. and Mahowald, A.P.** (1986). X-linked female sterile loci in *Drosophila melanogaster*. *Genetics* **113**, 695-712.
- Perrimon, N., Engstrom, L. and Mahowald, A. P.** (1989). Zygotic lethals with specific maternal effect phenotypes in *Drosophila melanogaster*. I. Loci on the X-chromosome. *Genetics* **121**, 333-352.
- Perrimon, N.** (1993) The Torso receptor tyrosine kinase signaling pathway: An endless story. *Cell*, in press.
- Price, J. V., Clifford, R. J. and Schupbach, R.** (1989) The maternal ventralizing locus *torpedo* is allelic to *faint little ball*, an embryonic lethal, and encodes the *Drosophila* EGF receptor homolog. *Cell* **56**, 1085-1092.
- Robertson, H. M., Preston, C. R., Phillis, R. W., Johnson-Schlitz, D., Benz, W. K. and Engels, W. R.** (1988). A stable source of P-element transposase in *Drosophila melanogaster*. *Genetics* **118**, 461-470.
- Rogge, R. D., Karlovich, C. A. and Banerjee, U.** (1991). Genetics dissection of a neurodevelopmental pathway: *Son of sevenless* functions downstream of the *sevenless* and EGF receptor tyrosine kinase. *Cell* **64**, 39-48.
- Roiha, H., Rubin, G. M. and O'Hare, K.** (1988). P element insertions and rearrangements at the singed locus of *Drosophila melanogaster*. *Genetics* **119**, 75-83.
- Ruohola-Baker, H., Grell, E., Chou, T.-B., Baker, D., Jan, L. Y. and Jan, Y. N.** (1993) Spatially localized Rhomboid is required for establishment of the dorsal-ventral axis in *Drosophila* oogenesis. *Cell* **73**, 953-965.
- Salz, H. K., Cline, T. W. and Schedl, P.** (1987). Functional changes associated with structural alterations induced by mobilization of a P element inserted in the sex-lethal gene of *Drosophila*. *Genetics* **117**, 221-231.
- Sambrook, J., Fritsch, E. F. and Maniatis, T.** (1989). *Molecular cloning: A Laboratory Manual*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press.
- Schupbach, T. and Wieschaus, E.** (1986). Germline autonomy of maternal-effect mutations altering the embryonic body pattern of *Drosophila*. *Dev. Biol.* **113**, 443-448.
- Schupbach, T. and Wieschaus, E.** (1989). Female sterile mutations on the second chromosome of *Drosophila melanogaster*. I. Maternal effect mutations. *Genetics* **121**, 101-117.
- Schupbach T.** (1987). Germline and soma cooperate during oogenesis to establish the dorso-ventral pattern of egg shell and embryo in *Drosophila melanogaster*. *Cell* **49**, 699-707.
- Schupbach, T., Clifford, R. J., Manseau, L. J. and Price, J. V.** (1991). *Dorso-ventral signaling processes in Drosophila oogenesis*. In *Cell-Cell interactions in early development* (ed. J. Gerhart). pp 163-174. New York: Wiley-Liss.
- Schejter, E. D. and Shilo, B.-Z.** (1989). The *Drosophila* EGF receptor homolog (DER) gene is allelic to *faint little ball*, a locus essential for embryonic development. *Cell* **56**, 1093-1104.
- Sheen, J.** (1993). Protein phosphatase activity required for light-inducible gene expression in maize. *EMBO J.* in press.
- Simon, A. M., Bowtell, D. D. L., Dodson, G. S., Laverty, T. R. and Rubin, G. M.** (1991). Ras1 and a putative guanine nucleotide exchange factor perform crucial steps in signaling by the sevenless protein tyrosine kinase. *Cell* **67**, 701-716.
- Simon, A. M., Dodson, G.S. and Rubin, G. M.** (1993). An SH3-SH2-SH3 protein is required for p21<sup>Ras1</sup> activation and binds to Sevenless and Sos proteins *in vitro*. *Cell* **73**, 169-177.
- Sosnowski, B. A., Belote, J. M. and McKeown, M.** (1989). Sex-specific, alternative splicing of RNA from the transformer gene results from sequence-dependent splice site blockage. *Cell* **58**, 449-459.
- Sturtevant, M. A., Roark, M. and Bier, E.** (1993) The *Drosophila rhomboid* gene mediates the localized formation of wing veins and interacts genetically with components of the EGF-R signaling pathway. *Genes Dev.* **7**, 961-973.
- Szabad, J., Erdelyi, M. and Szidonya, J.** (1987). Characterization of *Fs(2)1*, a germ line dependent dominant female sterile mutation of *Drosophila*. *Acta Biol. Hung.* **38**, 257-266.
- Szabad, J., Erdelyi, M. G., Hoffmann, M. G., Szidonya, J. and Wright, T. R. F.** (1989). Isolation and characterization of dominant female sterile mutations of *Drosophila melanogaster*. II. Mutations on the second chromosome. *Genetics* **122**, 823-835.
- Thummel, C. S. and Pirrotta, V.** (1992). New pCaSpeR P element vectors. *Dros. Inf. Service* **71**, 150.
- Tower, J., Karpen, G. H., Craig, N. and Spradling, A. C.** (1993). Preferential transposition of *Drosophila* P elements to nearby chromosomal sites. *Genetics* **133**, 347-359.
- van der Meer, J.** (1977). Optical clean and permanent whole mount preparations for phase contrast microscopy of cuticular structures of insect larvae. *Dros. Inf. Serv.* **52**, 160.
- Wieschaus, E.** (1980). A Combined Genetic and mosaic approach to the study of oogenesis. In '*Development and Neurobiology of Drosophila*', pp. 85-94. New York/London: Plenum Press.
- Wieschaus, E. and Szabad, J.** (1979). The developmental and function of the female germ line in *Drosophila melanogaster*: A cell lineage study. *Dev. Biol.* **68**, 29-46.
- Wieschaus, E., Audit, C. and Masson, M.** (1981). A clonal analysis of the roles of somatic cells and germ line during oogenesis in *Drosophila*. *Dev. Biol.* **88**, 92-103.
- Wieschaus, E., Nusslein-Volhard, C. and Kluding, H.** (1984) *Kruppel*, a gene whose activity is required early in the zygotic genome for normal embryonic segmentation. *Dev. Biol.* **104**, 172-186.
- Yarger, R. G. and King, R. C.** (1971). The phenogenetics of a temperature sensitive, autosomal dominant, female sterile gene in *Drosophila melanogaster*. *Dev. Biol.* **24**, 166-177.
- Zhang, P. and Spradling, A. C.** (1993) Efficient and dispersed local P element transposition from *Drosophila* females. *Genetics* **133**, 361-373.