

wingless signaling in the *Drosophila* eye and embryonic epidermis

Kenneth M. Cadigan and Roel Nusse

Howard Hughes Medical Institute and the Department of Developmental Biology, Stanford University School of Medicine, Stanford, California 94305, USA

SUMMARY

After the onset of pupation, sensory organ precursors, the progenitors of the interommatidial bristles, are selected in the developing *Drosophila* eye. We have found that *wingless*, when expressed ectopically in the eye via the *sevenless* promoter, blocks this process. Transgenic eyes have reduced expression of *acheate*, suggesting that *wingless* acts at the level of the proneural genes to block bristle development. This is in contrast to the wing, where *wingless* positively regulates *acheate* to promote bristle formation. The *sevenless* promoter is not active in the *acheate*-positive cells, indicating that the *wingless* is acting in a paracrine manner. Clonal analysis revealed a requirement for the genes *porcupine*, *dishevelled* and *armadillo* in mediating the *wingless* effect. Overexpression of *zeste white-3* partially blocks the ability of *wingless* to inhibit bristle formation, consistent with the notion that *wingless*

acts in opposition to *zeste white-3*. Thus the *wingless* signaling pathway in the eye appears to be very similar to that described in the embryo and wing. The *Notch* gene product has also been suggested to play a role in *wingless* signaling (J. P. Couso and A. M. Martinez Arias (1994) *Cell* 79, 259-72). Because *Notch* has many functions during eye development, including its role in inhibiting bristle formation through the neurogenic pathway, it is difficult to assess the relationship of *Notch* to *wingless* in the eye. However, we present evidence that *wingless* signaling still occurs normally in the complete absence of *Notch* protein in the embryonic epidermis. Thus, in the simplest model for *wingless* signalling, a direct role for *Notch* is unlikely.

Key words: *wingless*, signal transduction, *Notch*, *Drosophila*, neurogenesis, segment polarity

INTRODUCTION

The *wingless* (*wg*) gene is the best characterized member of the *Wnt* family, which contains over fifty genes in organisms ranging from nematodes to humans (Nusse and Varmus, 1992). *Wnt* genes encode cysteine-rich proteins containing signal sequences and several members, including *wg*, have rigorously been shown to be secreted (Bradley and Brown, 1990; Fradkin et al., 1995; González et al., 1991; Papkoff and Schryver, 1990; Van den Heuvel et al., 1989; Van Leeuwen et al., 1994).


In *Drosophila melanogaster*, *wg* is required throughout embryogenesis and larval development for a wide range of patterning events (Klingensmith and Nusse, 1994; Siegfried and Perrimon, 1994). Some of these include specifying cell fate in the embryonic epidermis (Baker, 1988; Bejsovec and Martinez-Arias, 1991; Dougan and Dinardo, 1992), CNS (Chu-Lagraff and Doe, 1993), mesoderm (Baylies et al., 1995; Wu et al., 1995) and endoderm (Hoppler and Bienz, 1995). In larval development, *wg* is required for patterning in leg (Couso et al., 1993; Diaz-Benjumea and Cohen, 1994; Struhl and Basler, 1993; Wilder and Perrimon, 1995) and wing (Couso et al., 1994; Diaz-Benjumea and Cohen, 1995; Phillips and Whittle, 1993) imaginal discs. In the eye, *wg* has recently been shown to be necessary for proper spacing of morphogenetic furrow initiation (Ma and Moses, 1995; Treisman and Rubin, 1995). How one signal can produce so many responses

remains an important unanswered question in developmental biology.

Consistent with being a secreted molecule, *wg* is thought to execute most of its functions in a paracrine manner. In the best documented cases, the range of *wg* action can vary from one (Vincent and Lawrence, 1994) to several (Hoppler and Bienz, 1995) cell diameters, though the exact limits of *wg* diffusion remain unclear (Axelrod et al., 1996; Peifer et al., 1991; Theisen et al., 1994). In a few cases, *wg* regulates gene expression in the same cells in which it is expressed, e.g. the activation of *cut* expression at the wing margin (Couso et al., 1994) and the regulation of its own expression in the embryo (Bejsovec and Wieschaus, 1993; Hooper, 1994; Yoffe et al., 1995). This embryonic autoregulation has been referred to as 'autocrine *wg* signaling' but it is not clear whether *wg* works in a truly autocrine manner. However, recent evidence indicates that *wg* autoregulation may have different genetic requirements than the paracrine signaling pathway of *wg* (Hooper, 1994; Manoukian et al., 1995; see discussion).

Three genes with embryonic phenotypes very similar to that of *wg* have been described (Klingensmith et al., 1989; Peifer and Wieschaus, 1990; Perrimon et al., 1989; Perrimon and Mahowald, 1987), *porcupine* (*porc*), *dishevelled* (*dsh*) and *armadillo* (*arm*). Another gene, *zeste white-3* (*zw3*; also known as *shaggy*) has a mutant phenotype (Perrimon and Smouse, 1989; Siegfried et al., 1992) very similar to that of embryos

where *wg* has been expressed ubiquitously (Noordermeer et al., 1992). Genetic epistasis (Noordermeer et al., 1994; Peifer et al., 1994b; Siegfried et al., 1994) have ordered these genes in the following genetic pathway:



porc has been shown to be involved in either secretion or subsequent diffusion of the *wg* protein (Siegfried et al., 1994; van den Heuvel et al., 1993a) and the other three genes are thought to be required for receiving the *wg* signal (Klingensmith and Nusse, 1994; Siegfried and Perrimon, 1994).

Recent work has revealed that many aspects of this embryonic *wg* signaling pathway are conserved in larval *Drosophila* tissues as well as in other organisms. Analysis of *dsh*, *zw3* and *arm* mutations in leg and wing imaginal discs indicates that these genes are required for *wg* signaling (Couso et al., 1994; Diaz-Benjumea and Cohen, 1994; Klingensmith et al., 1994; Peifer et al., 1991; Theisen et al., 1994). This has been best shown in the developing wing margin, where these genes mediate *wg* regulation of the *acheate* (*ac*) gene (Couso et al., 1994; Blair, 1994). The vertebrate homologs of these three genes have been shown to play a role in inducing dorsal mesoderm in *Xenopus* in a manner consistent with functioning in a *Wnt* signaling pathway (Dominguez et al., 1995; He et al., 1995; Heasman et al., 1994; Pierce and Kimelman, 1995; Rothbacher et al., 1995; Sokol et al., 1995).

The *wg* signaling pathway described above was first postulated based on extensive genetic analysis, but recent work indicates that some of the gene products may function directly with *wg* in a biochemical pathway. The *arm* gene encodes the *Drosophila* homolog of β -catenin (Peifer and Wieschaus, 1990), a component of vertebrate adherens junctions (Kemler, 1993). A similar junctional complex is found in flies (Peifer, 1993) but a substantial pool of cytoplasmic arm protein also exists (Peifer et al., 1994b; Van Leeuwen et al., 1994). *wg* signaling causes an accumulation of cytoplasmic arm protein (Peifer et al., 1994b; Van Leeuwen et al., 1994) caused by a dramatic decrease in arm protein turnover (Van Leeuwen et al., 1994). This accumulation is correlated with a reduction in phosphorylation of arm (Peifer et al., 1994a). This increase in arm protein is thought to somehow transduce the *wg* signal to the nucleus (Klingensmith and Nusse, 1994; Siegfried and Perrimon, 1994).

Consistent with the proposed genetic pathway, mutations in the other components of the *wg* pathway affect arm protein levels. The normal segmentally repeated accumulation of arm protein is absent in *wg*, *porc* and *dsh* mutants (Peifer et al., 1994b; Riggelman et al., 1990), while *zw3* mutants have uniformly high levels of arm protein (Peifer et al., 1994b; Siegfried et al., 1994). The *dsh* gene encodes a novel protein (Klingensmith et al., 1994; Theisen et al., 1994) containing a PDZ domain (Kennedy, 1995) that is phosphorylated in response to *wg* in embryos and cultured cells, and this phosphorylation is correlated with the ability of *dsh* to stabilize the arm protein (Yanagawa et al., 1995). *zw3* encodes a serine-threonine protein kinase that is homologous with mammalian glycogen synthase kinase-3 (Ruel et al., 1993a; Siegfried et al., 1992). At the present time, it is not clear whether any of the regulatory steps in the pathway are direct or how many missing components remain to be identified.

One new candidate for functioning in the *wg* pathway is the product of the *Notch* (*N*) gene, which encodes a transmem-

brane protein found on the surface of cells. N protein is thought to act as the receptor for the *Delta* (*Dl*) gene product in a signaling pathway involved in many aspects of development (Muskavitch, 1994; Artavanis-Tsakonas et al. 1995). Its potential role in the *wg* pathway is based on strong genetic interactions between *N* and *wg* mutations in several tissues, but primarily in the wing (Couso and Martinez Arias, 1994; Hing et al., 1994). It is possible that the role of *N* in the separate but oft-used pathway with *Dl* could mask a requirement for *N* in *wg* signaling when *N* mutant embryos or clones are examined. Because *N* is expressed at the cell surface and appears to act as a receptor, it has been postulated that *wg* encodes a ligand for the N protein (Couso and Martinez Arias, 1994).

This report describes a phenotype created by ectopic expression of *wg* during eye development. These transgenic animals lack the mechanosensory bristles normally surrounding each facet of the compound eye. This is the exact opposite effect seen in the wing, where *wg* is required for bristle formation (Couso et al., 1994; Phillips and Whittle, 1993). Despite this difference in regulation, the *wg* signal transduction machinery found in the embryo and wing also functions in the eye. Finally, the role of *N* in *wg* signaling was examined in the eye and in the embryonic epidermis, where, in the complete absence of N protein, *wg* signaling appears to occur normally. These data argue against a direct role for *N* in *wg* signaling.

MATERIALS AND METHODS

Fly stocks

The mutant alleles in components of the *wg* signaling pathway used in this study were: *wg^{LL}*, *wg^{LN}*, *wg^{CX4}*, *porc^{D8}*, *porc^{2E}*, *dsh^{V26}*, *dsh⁴⁷⁷*, *arm^{XM19}*, *arm^{25B}*, *sgg^{D127}* and *zw3^{M11}*. *wg^{CX4}* (van den Heuvel et al., 1993a,b), *dsh^{V26}* (Yanagawa et al., 1995) and *sgg^{D127}* (Ruel et al., 1993b) are null alleles, *wg^{LN}* encodes a non-secreted *wg* protein (van den Heuvel et al., 1993a,b), *wg^{LL}* is a temperature-sensitive allele (Baker, 1988) and the rest are characterized phenotypically as strong alleles (Klingensmith, 1993; Siegfried et al., 1992), except for the *arm* alleles, which are hypomorphs but are the strongest alleles that are cell viable when homozygous (Peifer et al., 1991). Two null alleles of *N*, *N^{264.40}* and *N⁵⁴¹⁹* (S. Artavanis-Tsakonas, personal comm.) and the temperature-sensitive alleles *N^{ts1}* (Cagan and Ready, 1989b), *Dl^{6E}* (Dietrich and Campos-Ortega, 1984) and *Dl^{RF}* (Parody and Muskavitch, 1993) were also used. For further information, see Lindsley and Zimm (1992).

A P-element construct placing the *wg* ORF under the control of the *sevenless* (*sev*) promoter (P[*sev-wg*]) was made by inserting the *XbaI/Clal* (blunt ended) fragment of the *wg* cDNA, pCV (Rijsewijk et al., 1987) into the *XbaI* and *BglIII* (blunt ended) sites of pSEW_a (Fortini et al., 1992), between the *sev* proximal promoter and 3' processing elements. pSEW_a also contains three tandem repeats of the *sev* enhancer 5' of the promoter. *yw⁶⁷* embryos were coinjected with P[*sev-wg*] and p τ 25.7 as described previously (Rubin and Spradling, 1982) and several independent lines were established using standard balancer stocks. A stock containing the *lacZ* coding sequences under the control of the *sev* enhancer (three tandem repeats) and *hsp70* proximal promoter (P[*sev-lacZ*]; R. Carthew, personal communication) was obtained from Todd Laverty (UC Berkeley, CA).

The following heat-shock strains were used: P[*hs-wg*] (Noordermeer et al., 1992), P[*hs-zw3*] (Siegfried et al., 1992) and P[*hs-dsh*] (Axelrod et al., 1996). P[*hs-wg*] is on the third chromosome, the other two on the second. The following chromosomes were created by recombination. P[*sev-wg*; *w⁻*], P[*hs-zw3*; *w⁺*] (the *white* (*w*) gene in

the P[*sev-wg*; *w*⁺] transgene was inactivated by EMS mutagenesis). A P[*sev-wg*] insert on chromosome 3L was recombined with a *Dl*^{RF} mutation to make P[*sev-wg*], *Dl*^{RF}. Two different P[*hs-dsh*; *w*⁺], *wg*^{LL} recombinants were created, one using a *wg*^{LL} *cn bw sp* chromosome and the other a *wg*^{LL} *br pr*, since both chromosomes contain a different lethal mutation unrelated to *wg* (Couso et al., 1994). Both P[*hs-dsh*], *wg*^{LL} recombinants were placed over a SM5a-TM6B compound chromosome, so that homozygotes could be identified by the absence of the *Tubby* pupal marker.

Whole-mount stainings of pupal eyes and embryos

Pupal eyes were dissected and then immunostained as described (Blochlinger et al., 1993). Embryo stainings were performed essentially as previously described [Frasch et al., 1987; Grossniklaus et al., 1992]. Affinity-purified rat α -*cut* antisera was generously provided by K. Blochinger (Fred Hutchinson Institute, WA), mouse α -*ac* monoclonal antibody was a gift of Sean Carroll (University of Wisconsin at Madison). Rabbit α -*lacZ* antisera was from Cappel and affinity purified rabbit α -*wg* antisera was kindly provided by C. Harryman-Samos (Stanford University, CA). Mouse α -*N* monoclonal antibody was provided by S. Artavanis-Tsakonas (Yale Univ. CT) and mouse α -*en* antisera by T. Kornberg (UCSF, CA). The primary antibodies were used at the following dilutions: *ac*, 1:3 to 1:5, *wg*, 1:20, *N*, 1:100, *cut* and *en*, 1:300, *lacZ*, 1:500. For histochemistry, secondary antibodies were either biotinylated (goat α -mouse, horse α -rabbit and rabbit α -rat; all from the Elite ABC kit, Vectastain, used at a 1:500 dilution) or goat α -rabbit conjugated to alkaline phosphatase (from Vector, used at 1:300). For fluorescence microscopy either donkey FITC α -mouse (1:100) or donkey Cy3 α -rabbit (1:200) were used (Jackson Immunochemicals). Confocal images were collected with a Bio-Rad MRC 1000 confocal laser setup attached to a Zeiss Axioscope microscope. Images were imported into Adobe Photoshop for presentation.

In situ hybridization to whole-mount embryos using digoxigenin-labeled probes (Tautz and Pfeiffle, 1989) and antibody/in situ double stainings (Manoukian and Krause, 1992) were performed as described (detailed protocol available upon request).

All whole-mount stainings were photographed with a Nikon Microphot-FXA microscope and slides were scanned into Adobe Photoshop for presentation.

Production of mosaic animals

Mutant alleles of *dsh*, *zw3* and *arm* were recombined onto a P[*hs-neo*; *FRT*]18A chromosome, *porc* onto P[*hs-neo*; *FRT*]19A, *wg* onto P[*hs-neo*; *FRT*]40A and a P[*sev-wg*; *w*⁺] mapping to 3L onto P[*hs-neo*; *FRT*]80A, all in a *w* background. *w* clones were induced in animals heterozygous with the appropriate P[*mini-w*⁺], P[*FRT*] chromosome: P[*mini-w*⁺; *hs- π M*]5A, 10D, P[*hs-neo*; *FRT*]18A; P[*mini-w*⁺]18A, P[*hs-neo*; *FRT*]19A; P[*mini-w*⁺; *hs- π M*]21C, 36F, P[*hs-neo*; *FRT*]. All *FRT* derivatives are as described (Xu and Rubin, 1993) except for P[*mini-w*⁺]18A, which is from the Jan lab enhancer detection collection (Bier et al. 1989). FLP recombinase was provided from the FLP-99 chromosome (Chou and Perrimon, 1992). Clones were induced by a one hour heat shock (37°C) 24-48 hours (at 25°C) after egg laying and scored for the absence of pigmentation in the adult eye.

For production of *N* germ-line clones, the *N* null alleles were recombined onto a P[*mini-w*⁺; *FRT*]101 chromosome (Chou and Perrimon, 1992). *N*, P[*mini-w*⁺; *FRT*]101/EM7 females were crossed to a *w* ovo^{D1}, P[*mini-w*⁺; *FRT*]101/Y; P[*hs-FLP*]38 stock (Chou and Perrimon, 1992) and progeny were heat shocked late 3rd instar/early pupation for 2 hours at 37°C (earlier heat shocks resulted in high lethality due to somatic clones). Mosaic mothers were crossed to P[*ftz-lacZ*]C males (Hiromi and Gehring, 1987) or P[*ftz-lacZ*]C; P[*hs-wg*]TM3 males. Embryos with no β -gal staining lacked both maternal and zygotic expression of *N*.

Heat shocks and other temperature shifts

The P[*hs-wg*] phenotype was induced by multiple heat shocks as pre-

viously described (Noordermeer et al., 1992). Late larval/early pupal temperature shifts were performed by submerging glass vials in a water bath of the appropriate temperature (37°C for heat shocks). At all other times, larvae and pupae were kept at 25°C. Formation of white pupae was used as the reference point (0 hours APF).

Histology

Flies were prepared for scanning electron microscopy by serial dehydration in ethanol and Freon 113 (EM Sciences) as described (Kimmel et al., 1990). Dried samples were mounted with colloidal graphite, and a 10 nm gold-platinum coat was applied with a Hummer sputter coater. The samples were viewed with an AMR1000 SEM and photographed using Polapan 400 film (Kodak). Pupal eyes were surface stained with Co(NO₃)₂·6H₂O and (NH₄)₂S as described (Kimmel et al., 1990).

RESULTS

wg blocks SOP formation in the eye

During the course of our attempts to create a dominant adult *wg* mutant through limited misexpression of *wg* during larval development, we found a highly penetrant phenotype when *wg* was placed under the control of the eye-specific promoter *sev*. As shown in Fig. 1, the eyes of P[*sev-wg*] flies appear normal, except that the interommatidial bristles, normally found at alternating vertices in the compound eye's hexagonal array, are almost completely missing. Sections through adult eyes (data not shown) and surface staining of pupal eyes with cobalt sulfide (Fig. 1E,F) revealed no other detectable abnormality in




Fig. 1. P[*sev-wg*] flies lack interommatidial bristles. SEM images of parental *yw*⁶⁷ (A,C) or P[*sev-wg*] (B,D) eyes showing lack of both the base and shaft of the bristles. Note that the hexagonal array and surface of each facet are unaffected in the transgenic eyes. Cobalt sulfide staining of pupal eyes (36 hours APF at 25°C) of control (E) and P[*sev-wg*] (F). Cone cells (c) and the 1°, 2° and 3° pigment cells appear normal in transgenic eyes, but a 3° pigment cell is found in place of each bristle (b).

adult eyes. The bristles are replaced in the repeated structure of the eye with tertiary pigment cells. Thus, at the level of ectopic *wg* expressed from the P[*sev-wg*] transgene, the effect of *wg* on eye development is very specific.

Interommatidial bristles are mechanosensory organs composed of four cells that are derived from a single sensory organ precursor (SOP; (Cagan and Ready, 1989a). Larval SOP determination has been best described in the wing imaginal disc (Campuzano and Modolell, 1992; Jan and Jan, 1993b). The process begins with small groups of cells expressing basic helix-loop-helix proteins such as *acheate* (*ac*) and *scute* (Cubas et al., 1991; Skeath and Carroll, 1991). All the cells in these proneural clusters have the ability to become the SOP, however, in a wild-type background, only one does. This cell is thought to become the SOP by reaching a threshold level of *ac* and/or *scute* after which it inhibits these genes' expression in its neighbors (Ghysen et al., 1993; Simpson, 1990). This lateral inhibition is mediated by the neurogenic pathway, in which the products of the *Dl* and *N* genes are thought to act as ligand and receptor, respectively (Artavanis-Tsakonas et al., 1995; Muskavitch, 1994). The initiation of SOP development is correlated with the expression of a new set of genes, such as *neuralized* (Huang and Dambly-Chaudière, 1991) and for some SOPs, *cut* (Blochlinger et al., 1993). The SOP undergoes to two divisions to generate the four cells that will give rise to the mature bristle organ (Bodmer et al., 1989; Hartenstein and Posakony, 1989).

The events leading to SOP formation in the eye have many similarities to those occurring in other tissues. *ac* protein becomes detectable shortly after white prepupa formation (data not shown). At 3 hours after the white prepupa stage (3 hours APF), the *ac* gene is expressed in small clusters of cells throughout the eye (Fig. 2C). Unlike the photoreceptors and cone cells, the appearance of the *ac*-positive cells is not related to the distance from the morphogenetic furrow, although the cells anterior of the furrow do not express *ac* (see arrows in Fig. 2C). By 6 hours APF, only one cell per cluster still expresses *ac*, again with the anterior-most portion of the eye showing a less mature pattern (data not shown). At 15 hours APF, after the eye disc everts, *ac* protein is gone, but the daughters of the SOPs can be observed by staining with α -*cut* antisera (Fig. 2A). Because of the complicated morphogenetic movements associated with the eye/head disc eversion, we have been unable to stain tissue between 6 and 15 hours APF.

In the P[SEV-*wg*] eyes, *ac* expression is greatly reduced compared to controls though not completely absent (Fig. 2D,F). After disc eversion, no SOPs are found, as judged by *cut* staining (Fig. 2B) and an enhancer detector line for the *neuralized* gene (data not shown). Thus, *wg* appears to act at the level of the proneural genes, i.e., *ac*, to inhibit SOP formation.

wg-dependent SOP inhibition is a paracrine effect

The activity of the *sev* promoter has been well studied in third instar larva, by monitoring endogenous *sev* expression (Tomlinson et al., 1987) and with chimeric constructs (Bowtell et al., 1989) using *sev* enhancer and promoter elements similar to the ones in P[*sev-wg*]. The enhancer is active in the cone cells and in a subset of the underlying photoreceptor precursors. No description of *sev* expression has been reported after pupation, so the possibility existed that *wg* was expressed in




Fig. 2. P[*sev-wg*] eyes have lower than normal levels of *ac* protein and no SOPs. *yw*⁶⁷ (A,C,E) and P[*sev-wg*] (B,D,F) pupal eyes were stained with antibodies against *cut* (A,B; 15 hours APF) or *ac* (C-F; 3 hours APF). E and F are confocal images. Pairs of *cut*-positive SOP daughter cells (see arrows) can be seen in controls (A) but not in the transgenic eyes (the arrows point to cone cells, which also express *cut* and lie in a slightly more apical focal plane). *ac* is expressed basally in small clusters up until the morphogenetic furrow (C; arrows indicate the approximate position of the furrow. Anterior is down). The *ac*-positive clusters usually consist of two or three cells (E; see arrows). *ac* protein remains in P[*sev-wg*] eyes to varying degrees (the image in D lies in the middle of the range; F shows a close up of an area with relatively high levels of *ac* expression), but staining is always significantly less than controls (E).

the proneural cells of P[*sev-wg*] eyes, suggesting a possible autocrine effect.

This question was addressed by examining the distribution of *wg* protein in P[*sev-wg*] eyes. Though *wg* is a secreted protein, it is found at the highest levels on the surface of the same cells that synthesize it (Bejsovec and Wieschaus, 1995; Couso et al., 1994; van den Heuvel et al., 1993). In P[*sev-wg*] eyes, the highest levels of *wg* protein were found around the four cone cells (Fig. 3A) and accumulated on their apical surface (Fig. 3B). In more basal sections of the eye, *wg* protein was associated with the photoreceptors, which extend basally to the same plane as the *ac*-positive cells (Fig. 3C). There was no significant overlap between *wg* protein and the remaining cells expressing *ac*.

To confirm that the *sev* enhancer was not active in the proneural clusters, we stained eyes of flies that contained a P[*sev-lacZ*] transgene (see Materials and Methods) for products of *lacZ* and *ac*. As found for *wg* in P[*sev-wg*] eyes, most of the β -gal was found in the cone cells (data not shown). In the same focal plane as the *ac*-expressing cells, there is no overlap (Fig. 3D). Thus, the inhibitory effect of *wg* on *ac* expression is paracrine in nature.




Fig. 3. The *sev* enhancer is not active in the *ac*-positive cells. P[*sev-wg*] (A-C) or P[*sev-lacZ*] (D) pupal eyes (all at 3 hours APF) were stained with antibodies against *wg* (A,B), *wg* and *ac* (C) or β -gal and *ac* (D) protein. All panels are confocal images, with *wg* and *lacZ* signals always in red and *ac* always in green. In control eyes, *wg* protein was detected in a ring around the periphery of the eye (data not shown), but no *wg* protein was detected in the eye proper. In P[*sev-wg*] eyes, *wg* is found primarily around the cone cells (A) and accumulated on their apical surface (B) but not in the few remaining *ac*-positive cells (C). The focal plane in C is about 15–20 μ m basal of those in A and B. In the P[*sev-lacZ*] eyes (where *ac* expression is not affected) no β -gal protein is seen in the proneural clusters (D).

The *wg* signal transduction pathway in the eye

Extensive genetic analysis, confirmed by recent biochemical experiments, has identified four genes that encode probable components of the *wg* signaling pathway, *porc*, *dsh*, *zw3* and *arm* (Klingensmith and Nusse, 1994; Siegfried and Perrimon, 1994; see introduction). Mosaic analysis (using the *w* gene as a marker) was performed to determine if these genes were required for the P[*sev-wg*]-dependent bristle inhibition. Control clones still lack bristles (Fig. 4A), as do clones mutant for the endogenous *wg* gene (Fig. 4C). In clones that lack the P[*sev-wg*] transgene, bristles are found almost to the clonal boundary (Fig. 4B). Likewise, 89% of the mutant clones for *porc*, *dsh* and *arm* had the full array of bristles within the clone (Fig. 4D–F and Table 1) and an additional 9% had a partial rescue of the bristleless phenotype. The remaining 2% that still lacked bristles were small in size and probably not completely mutant since the absence of the *w* gene cannot be detected on the surface of the eye at the cellular level. These experiments indicate that *porc*, *dsh* and *arm* are required for *wg*-dependent bristle inhibition.

zw3 is unique among the known genes required for *wg* signaling because it must be inhibited for the *wg* signal to be transduced (Klingensmith and Nusse, 1994; Siegfried and Perrimon, 1994). Thus, loss of *zw3*

should be equivalent to activation of *wg* signaling. Therefore, a *zw3* mutant clone in the eye might be expected to lack bristles. This straightforward analysis cannot be employed because the cells in *zw3* clones in the eye imaginal disc do not differentiate into eye tissue (Treisman and Rubin, 1995; data not shown). This is probably due to the fact that high levels of *wg* signaling activity prevent the morphogenetic furrow from progressing, blocking any subsequent differentiation (Treisman and Rubin, 1995).

If *zw3* must be inhibited for the *wg* signal to be transduced, then flooding cells with *zw3* protein might titrate out the signal. This has been shown to be the case in *Xenopus* where overexpression of the homologue of *zw3*, glycogen-synthase kinase 3, blocks *Wnt* gene induction of dorsal mesoderm (Dominguez et al., 1995; He et al., 1995). We attempted a similar experiment by creating flies with one copy of P[*sev-wg*] (we chose one of the weaker P[*sev-wg*] lines, which at one copy has approximately 20 bristles/eye) and one or two copies of a heat-shock construct expressing the *zw3* gene, P[*hs-zw3*] (Siegfried et al., 1992). *zw3* was induced by heat shock shortly before and twice after the onset of pupation (see Fig. 5 legend for details). Though the results were not entirely conclusive (Fig. 5), many pupal eyes showed a significant response especially when the ratio of P[*hs-zw3*]/P[*sev-wg*] is two (Fig. 5C). Other heat-shock regimes were not as effective at suppressing the P[*sev-wg*] phenotype. These results are consistent with the current model for *zw3* function in *wg* signaling.

Overexpression of *dsh* has previously been found to mimic *wg* signaling in cultured cells (Yanagawa et al., 1995), frog embryos (Sokol et al. 1995; Rothbacher et al. 1995) and in the wing imaginal disc (Axelrod et al., 1996). The same P[*hs-dsh*] transgenic stock used in the wing can also duplicate the effect of *wg* in the eye. Induction of *dsh* at 3 hours (data not shown) or 6 hours APF (Fig. 6B) could block bristle formation, but heat shock at 9 hours APF (Fig. 6C) failed to inhibit bristles in the interior of the eye, though inhibition still occurred toward the periphery. This can be explained by previous work




Fig. 4. The *porc*, *dsh* and *arm* genes are required for the P[*sev-wg*] phenotype, but the endogenous *wg* gene is not. Clones were induced in P[*sev-wg*] eyes as described in Materials and Methods. Clones were detected by the absence of pigmentation (from the *w* gene) in adult eyes. Bristles were still absent in control (A) or *wg*^{CX4} clones (C), but not in clones lacking the transgene (B) or homozygous for *porc* (D), *dsh* (E) and *arm* (F). A summary of all the data can be found in Table 1.

Table 1. Summary of the clonal analysis in a P[sev-wg] background (see Materials and Methods for details)

| Chromosome | Bristle density inside clone | | |
|----------------------------|------------------------------|---------|------|
| | Bare | Partial | Full |
| P[sev-wg; w ⁺] | 0 | 1 | 17 |
| w | 25 | 0 | 0 |
| yw | 21 | 1 | 0 |
| wg ^{CX4} | 23 | 0 | 0 |
| yw porc ^{2E} | 0 | 1 | 26 |
| yw porc ^{I8} | 2 | 5 | 25 |
| yw dsh ⁴⁷⁷ | 0 | 1 | 29 |
| yw dsh ^{V26} | 1 | 4 | 14 |
| w arm ^{25B} | 0 | 0 | 15 |
| w arm ^{XM19} | 0 | 1 | 11 |

The P[sev-wg; w⁺] clones are w;+/+ clones surrounded by w; P[sev-wg; w⁺] tissue. The rest are clones of the homozygous genotype indicated and the entire eye, including the cells in the clone, are P[sev-wg; w⁻]/+. Bare means no bristles found within the clone and full means the normal wild-type bristle density.

(Cagan and Ready, 1989a,b), which showed that SOP determination occurs first in the center of the eye and radiates outward concentrically. The same time requirements were seen when the bristles were inhibited using P[hs-wg] (data not shown).

Genetic and biochemical evidence places dsh downstream of wg in the signal transduction pathway (Klingensmith et al., 1994; Noordermeer et al., 1994; Theisen et al., 1994; Yanagawa et al., 1995), suggesting that the overexpression of dsh can bypass wg function. However, in the wing, where dsh overexpression causes an expansion of the wing margin, it appears that wg gene activity is needed to see the dsh effect (Axelrod et al., 1996). In the eye, the opposite appears to be true. In pupa homozygous for a wg temperature-sensitive mutation, induction of dsh after 6 hours at the restrictive temperature still inhibited SOP formation (Fig. 6E). Thus it appears that dsh in the eye can act independently of wg, though caveats remain (see discussion).

The role of N in wg signaling in the eye

A strong interaction between mutations in the N and wg genes has been described (Couso and Martinez Arias, 1994; Hing et al., 1994), which suggests that the two genes have common developmental targets in some tissues. One report suggested that wg encodes a ligand for N, based on these genetic interactions and the fact that N encodes a transmembrane receptor-like protein (Couso and Martinez Arias, 1994). In the eye, N activity is required for almost every differentiated cell type (Cagan and Ready, 1989b), so examining N clones in a P[sev-wg] background was not possible. Therefore, we utilized N^{ts1}, a temperature-sensitive allele (Cagan and Ready, 1989b). When these flies were reared at the restrictive temperature for 3-11 hours APF in a P[sev-wg] background, a strong suppression of the wg bristleless phenotype was seen (Fig. 7B). This is consistent with a proposed role for N in transducing the wg signal. However, removal of Dl activity for the same time period also suppresses the P[sev-wg] phenotype (Fig. 7C).

N and Dl are key components in the lateral inhibition pathway (functioning as receptor and ligand, respectively) that insures the proper number of bristles in the eye (Cagan and

Ready, 1989b; Parody and Muskavitch, 1993; note the abnormally high bristle density in Fig. 7B and C). This pathway is independent of wg, since mutant clones of wg, porc, dsh and arm in an otherwise wild-type background have the normal number of bristles (data not shown; see also Fig. 4D-F). Thus, the observation that loss of Dl activity can suppress the P[sev-wg] phenotype as well if not better than loss of N raises the possibility that the interaction between N and wg in the eye is due to the role of N in the lateral inhibition pathway.

If a higher level of wg expression is used (via a heat-shock promoter) all the bristles in the N^{ts1} background can be inhibited (data not shown; pupa were placed at the restrictive temperature for 6 hours before a 30 minute heat-shock pulse was given at 6 hours APF). However, it is known that the N^{ts1} allele does not completely remove N activity (Couso and Arias, 1994; Hartenstein et al., 1992) so this result is inconclusive. In the eye, it is not possible to determine whether wg works





Fig. 5. Overexpression of zw3 can suppress the P[sev-wg] phenotype. Three 1 hour heat shocks (37°C separated by two 4 hour recovery periods at 25°C) were given to animals containing one copy of P[sev-wg] and zero, one or two copies of P[hs-zw3] (the genotypes of the three groups were P[sev-wg; w⁺]/+, P[sev-wg; w⁺]/P[hs-zw3; w⁺] and P[sev-wg; w⁻], P[hs-zw3; w⁺]/P[hs-zw3; w⁺], respectively; all combinations were created from crosses of stocks described in Materials and Methods). The first heat shock was given at 1-2 hours prior to white pupa formation. An example of a control with about 25 SOPs (A) and a 1× P[hs-zw3] eye with about 180 SOPs (B) are shown. SOPs were detected with cut immunostaining. (C) The total data are summarized in a scatter plot. The mean number of SOPs are shown above each group, with the n value below. The standard deviation for the 0×, 1× and 2× groups were 15, 38 and 84, respectively. The differences between the 0× and the other two groups are significant at P<0.001 using a Student's t-test.

Fig. 6. Overexpression of *dsh* can inhibit bristle formation independently of *wg*. (A-C), SEM micrographs of P[*hs-dsh*] eyes given no heat shock (A) or a 30 minute heat shock (37°C) at 6 hours APF (B) or 9 hours APF (C). When *dsh* was induced at 6 hours APF, more than half the eyes had no or only a few bristles in the center of the eye ($n=11$) and the rest had a small patch of bristles in the center ($n=8$). At 9 hours APF, bristles were found over the interior two thirds of the eye but bristles were still missing toward the periphery ($n=17$). (D,E), cut stainings of P[*hs-dsh*], *wg^{LL}* homozygotes that were raised at 17°C (the permissive temperature) and then incubated at 29°C for 0-12 hours APF, without (D) or with (E) a 30 minute heat shock at 6 hours APF. Antibody stainings were done at ~30h APF. The cut-positive SOPs (now at the 4-cell stage) are completely absent in the heat shocked eyes ($n=8$). *wg^{LL}* homozygotes were identified as described in Materials and Methods.



through *N* or in a parallel pathway converging at proneural gene expression.

Role of *N* in *wg* signaling in the embryo

In order to more rigorously test the requirement of *N* for *wg* signaling, a tissue is needed where a putative *N*-*wg* connection can be separated from the *wg*-independent functions of *N*. One suitable place is the embryonic epidermis. Embryos mutant for *N* undergo a dramatic neural hyperplasia; almost all of the cells of the epidermis delaminate and become neuroblasts (Campos-Ortega, 1993). However, the epidermis remains relatively intact until full germ-band extension, after significant *wg* signaling has already occurred. Null *N* embryos were generated by making germ-line clones (Chou and Perrimon, 1992; see Materials and Methods). Antibody staining revealed no detectable *N* protein in *N* germline clones that have received a paternal Y chromosome (Fig. 8F). Thus we can examine *wg* signaling in a tissue that has never contained *N* protein.

Two well-characterized targets of *wg* signaling in the embryo are the *engrailed* (*en*) gene (DiNardo et al., 1988; Martinez-Arias et al., 1988) and the *wg* gene itself (Bejsovec and Wieschaus, 1993; Hooper, 1994; Yoffe et al., 1995). Careful analysis of expression of both genes has revealed that, in *wg* mutants, *wg* transcripts begin to fade before the embryo reaches full germ-band extension (stage 9; all stages according to (Campos-Ortega and Hartenstein, 1985), and is gone by the beginning of stage 10 (Manoukian et al., 1995). *en* protein in the adjacent posterior cells fades shortly thereafter. By mid-stage 10, both *en* protein and *wg* transcripts are completely gone from *wg^{IN}* homozygous embryos (Fig. 8B). In *N* null embryos at early stage 10, *wg* and *en* patterns are indistinguishable from wild type (data not shown). At mid-stage 10, both sets of stripes are still clearly present (Fig. 8C,D). The stripes do appear a little ragged, and we believe this is a con-

sequence of the beginning of the disintegration of the epidermis, which is well underway by late stage 10 (about 15-20 minutes later than the embryos shown in Fig. 8).

Despite the results in Fig. 8, it might be argued that in *N* mutants, perhaps *wg* and *en* expression no longer depended on *wg* activity. To address this, we examined the affect of global *wg* expression on *en* transcript distribution in a *N* mutant background. As previously reported (Noordermeer et al., 1992, 1994), overexpression of *wg* via a heat-shock promoter in an otherwise wild-type background causes a dramatic posterior expansion of the *en* stripes so that they are about twice as wide as normal (compare Fig. 9A and B). This expansion is still seen in embryos lacking *N* protein (Fig. 9D) and is dependent on the presence of the P[*hs-wg*] transgene (Fig. 9C). In the complete absence of *N* protein, *wg* signaling appears normal as late as we can reliably assay for it.




Fig. 7. Removal of *N* or *Dl* activity can suppress the P[*sev-wg*] bristleless phenotype. SEM micrographs of P[*sev-wg*]/+ (A), *N^{ts1}*/Y; P[*sev-wg*]/+ (B) and P[*sev-wg*], *Dl^{RF}*/*Dl^{6E}* (C) flies that were reared at 17°C and incubated at 32°C for 3 to 11 hours APF (7 hours APF at 17°C corresponds to 3 hours APF at 25°C) and then kept at 17°C until eclosion or dissection of pharates from pupal cases. Control and *N^{ts1}* hemizygotes were made by crossing P[*sev-wg*] males to either *w* or *w^{Nts1}* females. All males then had the desired genotype. P[*sev-wg*], *Dl^{RF}*/TM6C and *Dl^{6E}*/TM6C flies were crossed and appropriate animals identified by the absence of the dominant Tubby marker (found on TM6C). All *N^{ts1}* hemizygotes ($n=20$) and *Dl^{RF}*/*Dl^{6E}* transheterozygotes ($n=9$) showed the dramatic increase in bristle number. Note the higher than normal bristle density, indicative of the role these genes play in lateral inhibition. The *Dl* mutant combination consistently gave a more severe bristle hyperplasia than *N^{ts1}* in both a P[*sev-wg*] and non-transgenic background.

DISCUSSION

wg inhibits SOP formation at the level of the proneural genes

The interommatidial bristle is a 4-cell sensory organ that arises from a single SOP which is selected from a group of cells expressing proneural basic helix-loop-helix proteins (Campuzano and Modolell, 1992; Jan and Jan, 1993a). Our data strongly suggests that P[*sev-wg*]-derived *wg* protein blocks SOP formation in the eye by inhibiting proneural gene expression. Levels of *ac* protein are much lower in P[*sev-wg*] eyes (at 3 hours APF) compared to controls (Fig. 2C-F). 12 hours later, after the eye disc has everted, no SOP daughter cells are seen in the transgenic eyes (Fig. 2A,B). Though disc eversion prevents us from directly showing that no SOPs ever form in P[*sev-wg*] eyes, the time window when P[*hs-wg*] or P[*hs-dsh*] can inhibit bristle formation (no later than 6 hours APF for the central portion of the eye; Fig. 6 and results) is consistent with the model that, once an SOP is determined, *wg* signaling activity can no longer influence its fate.

The *ac* protein is the only proneural gene product monitored in this study and we are by no means suggesting that the *wg* signaling pathway acts directly on the *ac* promoter. In fact, loss of the *ac* gene alone does not result in complete elimination of interommatidial bristles; a related gene, *scute* (*sc*) must also be removed (Brown et al., 1991). The expression patterns of *ac* and *sc* are nearly identical (Cubas et al., 1991; Skeath and Carroll, 1991). This is most likely achieved by a combination of shared enhancer elements (Gómez-Skarmeta et al., 1995) and auto- and transactivation between the two genes (Martinez and Modolell, 1991; Skeath and Carroll, 1991; Van Doren et al., 1992). In addition, there are important negative inputs from other bHLH proteins such as *extramacrocheate* (Cubas and Modolell, 1992; Van Doren et al., 1992) and *hairy* (Brown et al., 1991; Van Doren et al., 1994). *wg* could be acting to inhibit *ac* (and presumably *sc*) expression at any of these regulatory levels. Further studies are needed to address this issue.

The P[*sev-wg*] bristleless phenotype was unexpected, because in the wing imaginal disc, *wg* has been shown to have the opposite effect, i.e., it promotes bristle development. In the absence of *wg* activity, the proneural *ac*-positive clusters fail to form (Couso et al., 1994; Phillips and Whittle, 1993). It is not clear why *wg* activates *ac* in one tissue and inhibits it in another, but this is a simple example of how one signal can generate different responses in various tissues.

wg is not normally expressed in the interior of the eye, but it is present at the periphery, forming a ring around the pupal eye (Cadigan and Nusse, unpublished data). Interestingly, the edge of the eye lacks bristles (Cagan and Ready, 1989b; Fig. 1A). Clones of *arm* at the

periphery contain ectopic bristles (Cadigan and Nusse, unpublished data), suggesting that *wg* normally inhibits bristles there. However, large *wg* clones do not show this effect. We are currently examining this in more detail.

The *wg* signal transduction pathway in the eye

A genetic pathway for *wg* signal transduction has been elucidated in which the gene products work in the following order: *porc* → *wg* → *dsh* —| *zw3* —| *arm* (Klingensmith and Nusse, 1994; Siegfried and Perrimon, 1994). Studies in the wing and leg imaginal disc have indicated that *dsh*, *zw3* and *arm* are also required there for *wg* signaling (Couso et al., 1994; Diaz-Benjumea and Cohen, 1994; Klingensmith et al., 1994; Peifer et al., 1991; Theisen et al., 1994; Wilder and Perrimon, 1995). This study extends these findings; *porc*, *dsh* and *arm* are clearly required for the ability of *wg* to inhibit eye bristles (Fig. 4; Table 1). The overexpression experiments with *zw3*, while not as conclusive (Fig. 5), are entirely consistent with the favored model, where *wg* acts by antagonizing *zw3* gene activity. While there may be exceptions (see below), it seems that most tissues use the same *wg* signaling components to achieve a variety of effects.

The mammalian counterpart of *zw3*, glycogen synthase kinase-3, has been shown to function in *ras*-dependent signaling (Stambolic and Woodgett, 1994). This raises the possibility that members of the *ras* and *wg* pathways share components in flies. In the eye, differentiation of photoreceptor cells is absolutely dependent on *ras*-dependent signaling (Simon et al., 1991). However, in clones of *dsh* and *arm*, all photoreceptors are present (S. Kaech, K.M. Cadigan and R. Nusse, unpublished observations). In the wing, clonal analysis with members of the *ras* pathway demonstrated that, unlike *wg*, they were not required for wing margin development (Diaz-Benjumea and Hafen, 1994). Thus, no interaction between these two pathways has yet been observed in *Drosophila*.




Fig. 8. *wg* signaling appears to be normal in *N* null mutant embryos. (A-D) Whole-mount staining for *wg* transcripts (blue) and/or *en* protein (brown) in wild-type (A), *wg*^{DN} (B) or *N*⁵⁴¹⁹ (C,D) mutant embryos. All embryos are at mid-stage 10 (Campos-Ortega and Hartenstein, 1985). Both *wg* and *en* are absent at this stage from the epidermis of the *wg* mutants, but remain robust in the *N* mutant background (these embryos were also stained for β -gal protein, to unambiguously identify maternal and zygotic *N* mutants (see Material and Methods). (E,F) Confocal images of *N* antibody staining with a monoclonal antibody directed against the intracellular domain of *N* (Fehon et al., 1990) in *N*⁵⁴¹⁹ germ-line clones receiving a paternal P[*fitz-lacZ*] (E) or Y (F) chromosome. *N* signal is completely lacking in the embryos that are negative for β -gal protein. Similar results in *wg*, *en* and *N* expression were obtained with a second *N* null allele, *N*^{264.40} (data not shown).




Fig. 9. The effect of ubiquitous expression of *wg* on *en* transcript distribution is still seen in a *N* null mutant background. All embryos are whole-mount stainings of *en* transcripts. (A) P[*hs-wg*] embryo with no heat shock. The *en* stripes are normal in appearance. (B) P[*hs-wg*] embryo after three 20 minute heat shocks (37°C) during early embryogenesis. The *en* stripes have expanded posteriorly, to about twice their normal width as previously described (Noordermeer et al., 1992). (C) *N*⁵⁴¹⁹ null mutant after the three heat shocks. The stripes are somewhat ragged, but still present at the normal width. (D) *N*⁵⁴¹⁹; P[*hs-wg*] embryo after heat-shock treatment. The stripes have broadened as they do in a *N*⁺ background. All embryos were mid-stage 10 and the same results were obtained using the *N*^{264.40} allele. *N* null embryos were created and identified as described in Materials and Methods.

wg expression is subject to positive autoregulation in the embryo (Bejsovec and Wieschaus, 1993; Hooper, 1994; Yoffe et al., 1995) and recent evidence suggests that this occurs through a distinct signaling mechanism (Hooper, 1994; Manoukian et al., 1995). Some discrepancies exist between the two reports, but Manoukian et al. (1995) provide strong evidence that *wg* autoregulation requires *porc* but not *dsh*, *zw3* and *arm*. They suggest a model where *porc* functions only in *wg* autoregulation and the other three genes in *wg* paracrine functions.

Our results in the eye indicate that, at least in the eye, *porc* is required for *wg* paracrine signaling. While we could clearly see *sev* enhancer-driven *wg* expression in cone cells and photoreceptors, we found no expression in the proneural clusters, the targets of *wg* action (Fig. 2). The endogenous *wg* gene was not required for the P[*sev-wg*]-dependent bristle inhibition (Fig. 4C), ruling out a paracrine-autocrine circuit. Our results indicating a role for *porc* in paracrine *wg* signaling are consistent with the observation that secretion or diffusion of *wg* protein is blocked in *porc* mutant embryos (Siegfried et al., 1994; van den Heuvel et al., 1993a).

Overexpression of *dsh* can mimic the action of *wg* in the eye (Fig. 6) as has been shown previously in the wing (Axelrod et al., 1996) and in cultured cells (Yanagawa et al., 1995). In the wing, this effect of *dsh* required *wg*. This does not appear to be the case in the eye (Fig. 6E). This is an important point because it speaks as to whether *dsh* can completely bypass the requirement for *wg* or whether overexpression of *dsh* simply potentiates *wg* signaling. It may be that there is residual *wg* activity left in our experiments (we could only rear the animals for 6 hours at the restrictive temperature before induction of *dsh*; longer times killed the organism before disc eversion). Another possibility is that a much higher threshold of *wg* activity is needed to transform wing blade to wing margin than

is needed to inhibit eye bristles. The data of Axelrod et al. (1996) show that the transformation of identity is more penetrant closest to the normal wing margin, where *wg* is expressed. Thus, overexpression of *dsh* in the wing blade may not easily reach the necessary level of signaling to trigger the change in cell fate. In the eye, *dsh* is able (at 3 hours APF) to inhibit bristles in the middle of the eye (far from endogenous *wg* expression) just as efficiently as bristles closer to the periphery. That *dsh* can bypass the need for *wg* is also supported by the cell culture experiments (Yanagawa et al., 1995) where no detectable *wg* protein was observed under conditions where *dsh* could stabilize arm protein. In addition, Park et al. (1996) have recently shown that overexpression of *dsh* in the embryo can induce *wg* targets in a *wg* null background.

Is *N* required for *wg* signaling?

On the basis of genetic interactions between mutations in the two genes, the *N* protein was proposed to be a receptor (or part of a receptor complex) for *wg* (Couso and Martinez Arias, 1994). In the eye, we also observed strong genetic interactions between *wg* and *N* (Fig. 7). However, the interpretation of these experiments are complicated, since *N* is known to affect bristle development independently of *wg*, and because, for technical reasons, we could not completely remove *N* activity to determine whether *wg* signaling could still occur. Likewise, the previously published genetic interactions involve animals where *wg* and *N* activities are only partially removed (many of the experiments were done with double heterozygotes of various *wg* and *N* alleles), and are therefore subject to the same limits of interpretation.

Unlike the eye, *wg* signaling in the complete absence of *N* activity can be assayed in the embryonic epidermis until just after germ-band extension is complete (mid stage 10), right before the absence of *N* causes most of the epidermis to delaminate and become neuroblasts. We found no significant change in the expression of *wg* and *en* in *N* null mutants at this time (Fig. 8), even though their expression fades at early stage 10 in *wg* mutants and mutants in *dsh* or *arm* (Manoukian et al., 1995; Van den Heuvel et al., 1993b). In addition, the effect of overexpression of *wg* on the *en* stripes is still seen in a *N* mutant background (Fig. 9). Couso and Martinez Arias (1994) reported that the *en* stripes were affected in about half the *N* mutants they examined, but they used hyperplasia of the nervous system as their method for determining which embryos were *N* mutants. This happens after mid-stage 10, thus any effect on the stripes may be a secondary consequence of the epidermis falling apart. Therefore, we conclude that in *N* mutant embryos, *wg* signaling occurs normally, at least with regard to the two markers we assayed.

A similar conclusion with regards to *N-wg* interactions has been reached in the wing (Rulifson and Blair, 1995). They showed that *wg* could still regulate *ac* expression in homozygous clones for a *N* null allele. These mutant clones should completely lack *N*, barring prolonged perdurance of the *N* protein. Of equal importance is their finding that *N* activity is required for *wg* expression at the wing margin (see also Diaz-Benjumea and Cohen, 1995; Doherty et al., 1996). This means that all of the genetic interactions between *wg* and *N* in the wing can potentially be explained by a reduction in *N* activity causing a reduction in the amount of *wg* signal, not the ability of *wg* to signal.

Another link between *wg* and *N* has been proposed by Axelrod et al. (1996), who have presented evidence that *dsh* protein can bind to and inhibit *N* activity in the wing imaginal disc. They suggest that part of the ability of *wg* to induce bristles in the wing is achieved by inhibition of *N* through *dsh*. Such an antagonistic relationship does not appear to be occurring in the eye since *wg*, *dsh* and *N* all inhibit bristle formation, although we can not rule out a mechanism where *wg* and *dsh* activate *N* to inhibit *ac* expression.

A subtle role for *N* in transducing the *wg* signal cannot be entirely ruled out. However, our results and those of Rulifson and Blair (1995) argue that in tissues where the direct test can be done, i.e., can *wg* signaling occur in cells that lack *N* protein, *N* is not required. A better candidate for a *wg* receptor is the product of the *Drosophila frizzled2* gene, which can bind to *wg* and transduce the *wg* signal in cultured cells (Bhanot et al. 1996). *N* showed no activity in this *wg*-binding assay. In the absence of any biochemical data suggesting that the proteins interact, the simplest models for *wg* signal transduction should exclude a direct role for *N*.

Special acknowledgment and thanks to Monty Laskosky, for superb operation of the SEM. We would also like to thank Dr Mike Simon for the pSEW construct, Drs Shu-wen Wang, Mark Muskavitch, Esther Siegfried, Todd Laverty, Juan Pablo Couso, Alfonso Martínez Arias, Kathy Matthews and the Bloomington Stock Center for various fly stocks, and especially Drs. Jeff Axelrod and Norbert Perrimon for providing the P[*hs-dsh*] flies prior to publication. Thanks also to all the researchers who provided antibodies (see Materials and Methods) and to Drs Sofia Lopes da Silva, Harsh Thaker, Andreas Wodarz, Diane Spillane and Derek Lessing for critical reading of the manuscript. Sectioning of the P[*sev-wg*] eyes was performed by Sue Kaech, to whom we are grateful. We would like to thank Dr Matt Scott for use of his confocal microscope, and Dr Andreas Wodarz for instruction on its proper use. Mike Ollman and Brent Wilson helped with Fig. 5B. These studies were supported by the Howard Hughes Medical Institute, of which R. N. is an investigator and K. M. C. is an associate, and by a grant from the USAMRAMC, Grant number DAMD17-94-J-4351.

REFERENCES

- Artavanis-Tsakonas, S., Matsuno, K. and Fortini, M. E. (1995). Notch signaling. *Science* **268**, 225-232.
- Axelrod, J. D., Matsuno, K., Artavanis-Tsakonas, S. and Perrimon, N. (1996). Interaction between Wingless and Notch signaling pathways mediated by Dishevelled. *Science* **271**, 1826-1832.
- Baker, N. E. (1988). Embryonic and imaginal requirements for wingless, a segment-polarity gene in *Drosophila*. *Dev. Biol.* **125**, 96-108.
- Baylies, M. K., Martínez Arias, A. and Bate, M. (1995). *wingless* is required for the formation of a subset of muscle founder cells during *Drosophila* embryogenesis. *Development* **121**, 3829-3837.
- Bejsovec, A. and Martínez-Arias, A. (1991). Roles of *wingless* in patterning the larval epidermis of *Drosophila*. *Development* **113**, 471-485.
- Bejsovec, A. and Wieschaus, E. (1993). Segment polarity gene interactions modulate epidermal patterning in *Drosophila* embryos. *Development* **119**, 501-517.
- Bejsovec, A. and Wieschaus, E. (1995). Signaling activities of the *Drosophila wingless* gene are separately mutable and appear to be transduced at the cell surface. *Genetics* **139**, 309-320.
- Bhanot, P., Brink, M., Harryman Samos, C., Hsieh, J.-C., Wang, Y., Macke, J. P., Andrew, D., Nathans, J. and Nusse, R. (1996). A new member of the *frizzled* gene family in *Drosophila* functioning as a wingless receptor. *Nature* (in press).
- Bier, E., Vaessin, H., Shepherd, S., Lee, K., McCall, K., Barbel, S., Ackermann, L., Carretto, R., Vemura, T., Grell, E., Jan, L. Y. and Jan, Y. N. (1989). Searching for pattern and mutation in the *Drosophila* genome with a P-lacZ vector. *Genes Dev.* **3**, 1273-1287.
- Blair, S. S. (1994). A role for the segment polarity gene *shaggy-zeste white 3* in the specification of regional identity in the developing wing of *Drosophila*. *Dev. Biol.* **162**, 229-244.
- Blochlinger, K., Jan, L. Y. and Jan, Y. N. (1993). Postembryonic patterns of expression of *cut*, a locus regulating sensory organ identity in *Drosophila*. *Development* **117**, 441-450.
- Bodmer, R., Carretto, R. and Jan, Y. N. (1989). Neurogenesis of the peripheral nervous system in *Drosophila* embryos: DNA replication patterns and cell lineages. *Neuron* **3**, 21-32.
- Bowtell, D. D. L., Kimmel, B. E., Simon, M. A. and Rubin, G. M. (1989). Regulation of the complex pattern of sevenless expression in the developing *Drosophila* eye. *Proc. Natl. Acad. Sci. USA* **86**, 6245-6249.
- Bradley, R. S. and Brown, A. M. (1990). The proto-oncogene *int-1* encodes a secreted protein associated with the extracellular matrix. *EMBO J.* **9**, 1569-1575.
- Brown, N. L., Sattler, C. A., Markey, D. R. and Carroll, S. B. (1991). *hairly* gene function in the *Drosophila* eye: normal expression is dispensable but ectopic expression alters cell fates. *Trends Genet.* **8**, 202-208.
- Cagan, R. L. and Ready, D. F. (1989a). The emergence of order in the *Drosophila* pupal retina. *Dev. Biol.* **136**, 346-362.
- Cagan, R. L. and Ready, D. F. (1989b). Notch is required for successive cell decisions in the developing *Drosophila* retina. *Genes Dev.* **3**, 1099-112.
- Campos-Ortega, J. A. (1993) Early neurogenesis in *Drosophila melanogaster*. In *The development of Drosophila melanogaster*. (ed. M. Bate and A. Martínez-Arias). pp. 1091-1129. Cold Spring Harbor: Cold Spring Harbor Laboratory.
- Campos-Ortega, J. A. and Hartenstein, V. (1985) *The Embryonic Development of Drosophila melanogaster*. Berlin: Springer-Verlag.
- Campuzano, S. and Modolell, J. (1992). Patterning of the *Drosophila* nervous system – the achaete-scute gene complex. *Trends Genet.* **8**, 202-208.
- Chou, T. B. and Perrimon, N. (1992). Use of a yeast site-specific recombinase to produce female germline chimeras in *Drosophila*. *Genetics* **131**, 643-653.
- Chu-Lagraff, Q. and Doe, C. (1993). Neuroblast specification and formation regulated by wingless in the *Drosophila* CNS. *Science* **261**, 1594-1597.
- Couso, J. P., Bate, M. and Martínez Arias, A. (1993). A *wingless*-dependent polar coordinate system in *Drosophila* imaginal discs. *Science* **259**, 484-489.
- Couso, J. P., Bishop, S. A. and Martínez Arias, A. (1994). The wingless signalling pathway and the patterning of the wing margin in *Drosophila*. *Development* **120**, 621-636.
- Couso, J. P. and Martínez Arias, A. M. (1994). Notch is required for wingless signaling in the epidermis of *Drosophila*. *Cell* **79**, 259-272.
- Cubas, P., de Celis, J.-F., Campuzano, S. and Modolell, J. (1991). Proneural clusters of *achaete-scute* expression and the generation of sensory organs in the *Drosophila* imaginal wing disc. *Genes Dev.* **5**, 996-1008.
- Cubas, P. and Modolell, J. (1992). The Extramacrochaetae Gene Provides Information for Sensory Organ Patterning. *EMBO J.* **11**, 3385-3393.
- Diaz-Benjumea, F. J. and Hafen, E. (1994). The sevenless signalling cassette mediates *Drosophila* EGF receptor function during epidermal development. *Development* **120**, 569-578.
- Diaz-Benjumea, F. J. and Cohen, S. M. (1994). *wingless* acts through the shaggy/zeste-white 3 kinase to direct dorsal-ventral axis formation in the *Drosophila* leg. *Development* **120**, 1661-1670.
- Diaz-Benjumea, F. J. and Cohen, S. M. (1995). Serrate signals through notch to establish a wingless-dependent organizer at the dorsal/ventral compartment boundary of the *Drosophila* wing. *Development* **121**, 4215-4225.
- Dietrich, U. and Campos-Ortega, J. A. (1984). The expression of neurogenic loci in imaginal epidermal cells of *Drosophila melanogaster*. *J. Neurogenetics* **1**, 315-332.
- DiNardo, S., Sher, E., Heemskerk-Jorgens, J., Kassis, J. and O'Farrell, P. (1988). Two-tiered regulation of spatially patterned *engrailed* gene expression during *Drosophila* embryogenesis. *Nature* **332**, 604-609.
- Doherty, D., Feger, G., Younger-Shepherd, S., Jan, L. Y. and Jan, Y. N. (1996). Delta is a ventral to dorsal signal complementary to Serrate, another Notch ligand, in *Drosophila* wing formation. *Genes Dev.* **10**, 421-434.
- Dominguez, I., Itoh, K. and Sokol, S. Y. (1995). Role of glycogen synthase kinase 3 beta as a negative regulator of dorsoventral axis formation in *Xenopus* embryos. *Proc. Natl. Acad. Sci. USA* **92**, 8498-8502.
- Dougan, S. and Dinardo, S. (1992). *Drosophila wingless* generates cell type diversity among engrailed expressing cells. *Nature* **360**, 347-350.
- Fehon, R. G., Kooh, P. J., Rebay, I., Regan, C. L., Xu, T., Muskavitch, M. A. T. and Artavanis-Tsakonas, S. (1990). Molecular interactions between

- the protein products of the neurogenic loci *Notch* and *Delta*, two EGF-homologous genes in *Drosophila*. *Cell* **61**, 523-534.
- Fortini, M., Simon, M. and Rubin, G.** (1992). Signalling by the sevenless protein tyrosine kinase is mimicked by Ras1 activation. *Nature* **355**, 559-561.
- Fradkin, L. G., Noordermeer, J. N. and Nusse, R.** (1995). The *Drosophila* Wnt protein Dwtnt-3 is a secreted glycoprotein localized on the axon tracts of the embryonic CNS. *Dev. Biol.* **168**, 202-213.
- Frasch, M., Hoey, T., Rushlow, C., Doyle, H. and Levine, M.** (1987). Characterization and localization of the even-skipped protein of *Drosophila*. *EMBO J.* **6**, 749-759.
- Ghysen, A., Damblychaudiere, C., Jan, L. Y. and Jan, Y. N.** (1993). Cell interactions and gene interactions in peripheral neurogenesis. *Genes Dev.* **7**, 723-733.
- Gómez-Skarmeta, J. L., Rodriguez, I., Martínez, C., Culi, J., Ferrés-Marcó, D., Beamonte, D. and Modolell, J.** (1995). *Cis*-regulation of *achaete* and *scute*: shared enhancer-like elements drive their coexpression in proneural clusters of the imaginal discs. *Genes Dev.* **9**, 1869-1882.
- González, F., Swales, L., Bejsovec, A., Skaer, H. and Martínez-Arias, A.** (1991). Secretion and movement of *wingless* protein in the epidermis of the *Drosophila* embryo. *Mech. Dev.* **35**, 43-54.
- Grossniklaus, U., Kurth-Pearson, R. and Gehring, W. J.** (1992). The *Drosophila* *sloppy paired* locus encodes two proteins involved in segmentation that show homology to mammalian transcription factors. *Genes Dev.* **6**, 1030-1051.
- Hartenstein, A. Y., Rugendorff, A., Tepass, U. and Hartenstein, V.** (1992). The Function of the Neurogenic Genes During Epithelial Development in the *Drosophila* Embryo. *Development* **116**, 1203-1220.
- Hartenstein, V. and Posakony, J. W.** (1989). Development of adult sensilla on the wing and notum of *Drosophila melanogaster*. *Development* **107**, 389-405.
- He, X., Saint-Jeannet, J.-P., Woodgett, J. R., Varmus, H. E. and Dawid, I. B.** (1995). Glycogen synthase kinase-3 and dorsoventral patterning in *Xenopus* embryos. *Nature* **374**, 617-622.
- Heasman, J., Crawford, A., Goldstone, K., Garner-Hamrick, P., Gumbiner, B., McCrea, P., Kintner, C., Noro, C. Y. and Wylie, C.** (1994). Overexpression of cadherins and underexpression of β -catenin inhibit dorsal mesoderm induction in early *Xenopus* embryos. *Cell* **79**, 791-803.
- Hing, H. K., Sun, X. and Artavanis-Tsakonas, S.** (1994). Modulation of wingless signaling by notch in *Drosophila*. *Mech. Dev.* **47**, 261-268.
- Hiroimi, Y. and Gehring, W. J.** (1987). Regulation and function of the *Drosophila* segmentation gene *fushi tarazu*. *Cell* **50**, 963-74.
- Hooper, J. E.** (1994). Distinct pathways for autocrine and paracrine Wingless signalling in *Drosophila* embryos. *Nature* **372**, 461-464.
- Hoppler, S. and Bienz, M.** (1995). Two different thresholds of wingless signalling with distinct developmental consequences in the *Drosophila* midgut. *EMBO J.* **14**, 5016-5026.
- Huang, F. and Dambly-Chaudière, C.** (1991). The emergence of sense organs in the wing disc of *Drosophila*. *Development* **111**, 1087-1095.
- Jan, Y. N. and Jan, L. Y.** (1993a). HLH proteins, fly neurogenesis and vertebrate myogenesis. *Cell* **75**, 827-830.
- Jan, Y. N. and Jan, L. Y.** (1993b). The peripheral nervous system. In *The Development of Drosophila melanogaster*. (ed. M. Bate and A. Martínez Arias). pp. 1207-1244. Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
- Kemler, R.** (1993). From cadherins to catenins: cytoplasmic protein interactions and regulation of cell adhesion. *Trends Genetics* **9**, 317-321.
- Kennedy, M. B.** (1995). Origin of PDZ (DHR, GLGF) domains [letter]. *Trends Biochem. Sci.* **20**, 350.
- Kimmel, B. E., Heberlein, U. and Rubin, G. M.** (1990). The homeo domain protein *rough* is expressed in a subset of cells in the developing *Drosophila* eye where it can specify photoreceptor cell subtype. *Genes Dev.* **4**, 712-727.
- Klingensmith, J., Noll, E. and Perrimon, N.** (1989). The segment polarity phenotype of *Drosophila* involves differential tendencies toward transformation and cell death. *Dev. Biol.* **134**, 130-145.
- Klingensmith, J.** (1993). Genetic dissection of an intercellular signaling pathway in *Drosophila* pattern formation. Harvard.
- Klingensmith, J., Nusse, R. and Perrimon, N.** (1994). The *Drosophila* segment polarity gene *dishevelled* encodes a novel protein required for response to the *wingless* signal. *Genes Dev.* **8**, 118-130.
- Klingensmith, J. and Nusse, R.** (1994). Signaling by wingless in *Drosophila*. *Dev. Biol.* **166**, 396-414.
- Lindsley, D. L. and Zimm, G. G.** (1992) *The Genome of Drosophila melanogaster*. San Diego: Academic Press, Inc.
- Ma, C. Y. and Moses, K.** (1995). Wingless and patched are negative regulators of the morphogenetic furrow and can affect tissue polarity in the developing *Drosophila* compound eye. *Development* **121**, 2279-2289.
- Manoukian, A. S. and Krause, H. M.** (1992). Concentration-Dependent Activities of the even-skipped Protein in *Drosophila* Embryos. *Genes Dev.* **6**, 1740-1751.
- Manoukian, A. S., Yoffe, K. B., Wilder, E. L. and Perrimon, N.** (1995). The porcupine gene is required for wingless autoregulation in *Drosophila*. *Development* **121**, 4037-4044.
- Martínez, C. and Modolell, J.** (1991). Cross-regulatory interactions between the proneural *achaete* and *scute* genes of *Drosophila*. *Science* **251**, 1485-1487.
- Martínez-Arias, A., Baker, N. E. and Ingham, P. W.** (1988). Role of segment polarity genes in the definition and maintenance of cell states in the *Drosophila* embryo. *Development* **103**, 157-170.
- Muskavitch, M. A. T.** (1994). Delta-Notch signaling and *Drosophila* cell fate choice. *Dev. Biol.* **166**, 415-430.
- Noordermeer, J., Johnston, P., Rijsewijk, F., Nusse, R. and Lawrence, P.** (1992). The consequences of ubiquitous expression of the wingless gene in the *Drosophila* embryo. *Development* **116**, 711-719.
- Noordermeer, J., Klingensmith, J., Perrimon, N. and Nusse, R.** (1994). *dishevelled* and *armadillo* act in the *wingless* signalling pathway in *Drosophila*. *Nature* **367**, 80-83.
- Nusse, R. and Varmus, H. E.** (1992). Wnt Genes. *Cell* **69**, 1073-1087.
- Papkoff, J. and Schryver, B.** (1990). Secreted int-1 protein is associated with the cell surface. *Mol. Cell. Biol.* **10**, 2723-2730.
- Park, M., Wu, X., Golden, K., Axelrod, J. D. and Bodmer, R.** (1996). The *wingless* signaling pathway is directly involved in *Drosophila* heart development. *Dev. Biol.* in press.
- Parody, T. R. and Muskavitch, M. A. T.** (1993). The pleiotropic function of delta during postembryonic development of *Drosophila melanogaster*. *Genetics* **135**, 527-539.
- Peifer, M. and Wieschaus, E.** (1990). The segment polarity gene *armadillo* encodes an evolutionarily conserved and functionally modular protein that is the *Drosophila* homolog of human plakoglobin. *Cell* **63**, 1167-1178.
- Peifer, M., Rauskolb, C., Williams, M., Riggleman, B. and Wieschaus, E.** (1991). The segment polarity gene *armadillo* interacts with the *wingless* signaling pathway in both embryonic and adult pattern formation. *Development* **111**, 1029-1043.
- Peifer, M.** (1993). The product of the *Drosophila* segment polarity gene *armadillo* is part of a multi-protein complex resembling the vertebrate adherens junction. *J. Cell Sci.* **105**, 993-1000.
- Peifer, M., Pai, L.-M. and Casey, M.** (1994a). Phosphorylation of the *Drosophila* adherens junction protein armadillo: roles for wingless signal and zeste-white 3 kinase. *Dev. Biol.* **166**, 543-566.
- Peifer, M., Sweeton, D., Casey, M. and Wieschaus, E.** (1994b). *wingless* signal and zeste-white 3 kinase trigger opposing changes in the intracellular distribution of armadillo. *Development* **120**, 369-380.
- Perrimon, N. and Mahowald, A. P.** (1987). Multiple functions of segment polarity genes in *Drosophila*. *Dev. Biol.* **119**, 587-600.
- Perrimon, N., Engstrom, L. and Mahowald, A. P.** (1989). Zygotic lethals with specific maternal effect phenotypes in *Drosophila melanogaster*. I. Loci on the X chromosome. *Genetics* **121**, 333-352.
- Perrimon, N. and Smouse, D.** (1989). Multiple functions of a *Drosophila* homeotic gene, zeste-white 3, during segmentation and neurogenesis. *Dev. Biol.* **135**, 287-305.
- Phillips, R. G. and Whittle, J. R. S.** (1993). *wingless* expression mediates determination of peripheral nervous system elements in late stages of *Drosophila* wing disc development. *Development* **118**, 427-438.
- Pierce, S. B. and Kimelman, D.** (1995). Regulation of Spemann organizer formation by the intracellular kinase Xgsk-3. *Development* **121**, 755-765.
- Riggleman, B., Schedl, P. and Wieschaus, E.** (1990). Spatial expression of the *Drosophila* segment polarity gene *armadillo* is post-transcriptionally regulated by *wingless*. *Cell* **63**, 549-560.
- Rijsewijk, F., Schuermann, M., Wagenaar, E., Parren, P., Weigel, D. and Nusse, R.** (1987). The *Drosophila* homolog of the mouse mammary oncogene int-1 is identical to the segment polarity gene *wingless*. *Cell* **50**, 649-657.
- Rothbacher, U., Laurent, M. N., Blitz, I. L., Watabe, T., Marsh, J. L. and Cho, K. W. Y.** (1995). Functional conservation of the wnt signaling pathway revealed by ectopic expression of *Drosophila* *dishevelled* in *Xenopus*. *Dev. Biol.* **170**, 717-721.
- Rubin, G. M. and Spradling, A. C.** (1982). Genetic transformation of *Drosophila* with transposable element vectors. *Science* **218**, 348-53.

- Ruel, L., Bourouis, M., Heitzler, P., Pantescio, V. and Simpson, P. (1993a). Drosophila shaggy Kinase and Rat Glycogen Synthase Kinase-3 Have Conserved Activities and Act Downstream of Notch. *Nature* **362**, 557-560.
- Ruel, L., Pantescio, V., Lutz, Y., Simpson, P. and Bourouis, M. (1993b). Functional Significance of a Family of Protein Kinases Encoded at the shaggy Locus in Drosophila. *EMBO J.* **12**, 1657-1669.
- Rulifson, E. J. and Blair, S. S. (1995). Notch regulates wingless expression and is not required for reception of the paracrine wingless signal during wing margin neurogenesis in Drosophila. *Development* **121**, 2813-2824.
- Siegfried, E., Chou, T. B. and Perrimon, N. (1992). wingless signaling acts through zeste-white 3, the Drosophila homolog of glycogen synthase kinase-3, to regulate engrailed and establish cell fate. *Cell* **71**, 1167-1179.
- Siegfried, E. and Perrimon, N. (1994). Drosophila wingless: a paradigm for the function and mechanism of wnt signaling. *BioEssays* **16**, 395-404.
- Siegfried, E., Wilder, E. L. and Perrimon, N. (1994). Components of wingless signalling in drosophila. *Nature* **367**, 76-80.
- Simon, M. A., Bowtell, D. D. L., Dodson, G. S., Laverty, T. R. and Rubin, G. M. (1991). Ras1 and a putative guanine nucleotide exchange factor perform crucial steps in signaling by the sevenless protein tyrosine kinase. *Cell* **67**, 701-716.
- Simpson, P. (1990). Lateral inhibition and the development of the sensory bristles of the adult peripheral nervous system of Drosophila. *Development* **109**, 509-519.
- Skeath, J. B. and Carroll, S. B. (1991). Regulation of *achaete-scute* gene expression and sensory organ pattern formation in the *Drosophila* wing. *Genes Dev.* **5**, 984-995.
- Sokol, S. Y., Klingensmith, J., Perrimon, N. and Itoh, K. (1995). Dorsalizing and neuralizing properties of Xdsh, a maternally expressed *Xenopus* homolog of *dishevelled*. *Development* **121**, 1637-1647.
- Stambolic, V. and Woodgett, J. R. (1994). Mitogen inactivation of glycogen synthase kinase-3 β in intact cells via serine 9 phosphorylation. *Biochem. J.* **303**, 701-704.
- Struhl, G. and Basler, K. (1993). Organizing activity of wingless protein in Drosophila. *Cell* **72**, 527-540.
- Tautz, D. and Pfeiffle, C. (1989). A non-radioactive in situ hybridization method for the localization of specific RNAs in *Drosophila* embryos reveals translational control of the segmentation gene *hunchback*. *Chromosoma* **98**, 81-85.
- Theisen, H., Purcell, J., Bennett, M., Kansagara, D., Syed, A. and Marsh, J. L. (1994). *dishevelled* is required during wingless signaling to establish both cell polarity and cell identity. *Development* **120**, 347-360.
- Tomlinson, A., Bowtell, D. D., Hafen, E. and Rubin, G. M. (1987). Localization of the sevenless protein, a putative receptor for positional information, in the eye imaginal disc of Drosophila. *Cell* **51**, 143-50.
- Treisman, J. E. and Rubin, G. M. (1995). wingless inhibits morphogenetic furrow movement in the drosophila eye disc. *Development* **121**, 3519-3527.
- Van den Heuvel, M., Nusse, R., Johnston, P. and Lawrence, P. A. (1989). Distribution of the *wingless* gene product in *Drosophila* embryos: A protein involved in cell-cell communication. *Cell* **59**, 739-749.
- Van den Heuvel, M., Harryman-Samos, C., Klingensmith, J., Perrimon, N. and Nusse, R. (1993a). Mutations in the segment polarity genes *wingless* and *porcupine* impair secretion of the wingless protein. *EMBO J.* **12**, 5293-5302.
- Van den Heuvel, M., Klingensmith, J., Perrimon, N. and Nusse, R. (1993b). Cell patterning in the drosophila segment: engrailed and wingless antigen distributions in segment polarity mutant embryos. *Development* **1993 Supplement** 105-114.
- Van Doren, M., Bailey, A. M., Esnayra, J., Ede, K. and Posakony, J. W. (1994). Negative regulation of proneural gene activity: hairy is a direct transcriptional repressor of *achaete*. *Genes Dev.* **8**, 2729-2742.
- Van Doren, M., Powell, P. A., Pasternak, D., Singson, A. and Posakony, J. W. (1992). Spatial regulation of proneural gene activity: auto- and cross-activation of *achaete* is antagonized by *extramacrochaetae*. *Genes Dev.* **6**, 2592-2605.
- Van Leeuwen, F., Harryman Samos, C. and Nusse, R. (1994). Biological activity of soluble wingless protein in cultured Drosophila imaginal disc cells. *Nature* **368**, 342-344.
- Vincent, J.-P. and Lawrence, P. A. (1994). Drosophila *wingless* sustains *engrailed* expression only in adjoining cells: evidence from mosaic embryos. *Cell* **77**, 909-915.
- Wilder, E. L. and Perrimon, N. (1995). Dual functions of *wingless* in the *Drosophila* leg imaginal disc. *Development* **121**, 477-488.
- Wu, X. S., Golden, K. and Bodmer, R. (1995). Heart development in Drosophila requires the segment polarity gene *wingless*. *Dev. Biol.* **169**, 619-628.
- Xu, T. and Rubin, G. M. (1993). Analysis of genetic mosaics in developing and adult drosophila tissues. *Development* **117**, 1223-1237.
- Yanagawa, S., Van Leeuwen, F., Wodarz, A., Klingensmith, J. and Nusse, R. (1995). The Dishevelled protein is modified by Wingless signaling in Drosophila. *Genes Dev.* **9**, 1087-1097.
- Yoffe, K. B., Manoukian, A. S., Wilder, E. L., Brand, A. H. and Perrimon, N. (1995). Evidence for *engrailed*-independent *wingless* autoregulation in *Drosophila*. *Dev. Biol.* **170**, 636-650.

(Accepted 15 June 1996)