

Delta-Notch signalling and the patterning of sensory cell differentiation in the zebrafish ear: evidence from the *mind bomb* mutant

Catherine Haddon, Yun-Jin Jiang, Lucy Smithers and Julian Lewis*

Vertebrate Development Laboratory, Imperial Cancer Research Fund, PO Box 123, Lincoln's Inn Fields, London WC2A 3PX, UK

*Author for correspondence (e-mail: j.lewis@icrf.icnet.uk)

Accepted 1 September; published on WWW 9 November 1998

SUMMARY

Mechanosensory hair cells in the sensory patches of the vertebrate ear are interspersed among supporting cells, forming a fine-grained pattern of alternating cell types. Analogies with *Drosophila* mechanosensory bristle development suggest that this pattern could be generated through lateral inhibition mediated by Notch signalling. In the zebrafish ear rudiment, homologues of *Notch* are widely expressed, while the *Delta* homologues *deltaA*, *deltaB* and *deltaD*, coding for Notch ligands, are expressed in small numbers of cells in regions where hair cells are soon to differentiate. This suggests that the *delta*-expressing cells are nascent hair cells, in agreement with findings for *Delta1* in the chick. According to the lateral inhibition hypothesis, the nascent hair cells, by expressing Delta protein, would inhibit their neighbours from becoming hair cells, forcing them to be supporting cells instead. The zebrafish *mind bomb* mutant has abnormalities in the central nervous system, somites, and elsewhere, diagnostic of a failure of

Delta-Notch signalling: in the CNS, it shows a neurogenic phenotype accompanied by misregulated *delta* gene expression. Similar misregulation of *delta* genes is seen in the ear, along with misregulation of a *Serrate* homologue, *serrateB*, coding for an alternative Notch ligand. Most dramatically, the sensory patches in the *mind bomb* ear consist solely of hair cells, which are produced in great excess and prematurely; at 36 hours post fertilization, there are more than ten times as many as normal, while supporting cells are absent. A twofold increase is seen in the number of otic neurons also. The findings are strong evidence that lateral inhibition mediated by Delta-Notch signalling controls the pattern of sensory cell differentiation in the ear.

Key words: Delta, Notch, Serrate, Ear, *mind bomb*, Hair cell, Lateral inhibition, Zebrafish

INTRODUCTION

The mechanosensory receptor cells of the inner ear – the hair cells – lie in specialised patches in the inner ear epithelium. Each such sensory patch consists of a fine-grained mixture of hair cells and supporting cells. An individual hair cell within a patch is generally surrounded by supporting cells, which isolate it from the next hair cell (Goodyear and Richardson, 1997); in some cases, as in the mammalian organ of Corti or the avian basilar papilla, the arrangement is remarkably regular, with the hair cells interspersed among the supporting cells in a precise periodic spacing pattern. Hair cells and supporting cells arise from a common precursor (Stone and Cotanche, 1994; Weisleder et al., 1995; Jones and Corwin, 1996). The challenge, therefore, is to explain what causes cells that are sisters or cousins, sitting side by side in the same environment, to differentiate in contrasting ways.

Lateral inhibition provides one possible mechanism: each cell that begins to differentiate as a hair cell might deliver a signal to its immediate neighbours in the developing sensory patch, inhibiting them from differentiating in the same way, with the result that they differentiate as supporting cells instead

(Corwin et al., 1991; Lewis, 1991). Studies in *Drosophila*, in *C. elegans*, and more recently in vertebrates have shown that signalling of this sort controls cell differentiation in several different tissues, where it is mediated by the transmembrane protein Notch, acting as signal receptor, and the transmembrane protein Delta, acting as ligand (Artavanis-Tsakonas et al., 1995; Lewis, 1996). In the central nervous system, for example, nascent neural cells express Delta and thereby inhibit their neighbours, which express Notch, from simultaneously differentiating along a neural pathway (Campos-Ortega, 1993; Chitnis et al., 1995). The lateral inhibition is thought to operate competitively because of a feedback loop that tends to amplify any initial difference between neighbouring cells: a cell that expresses Delta more strongly activates Notch more strongly in its neighbours; and activation of Notch in these cells inhibits not only their differentiation, but also their expression of Delta, thereby reducing their ability to deliver lateral inhibition in return (Heitzler and Simpson, 1991; Ghysen et al., 1993; Sternberg, 1993; Wilkinson et al., 1994; Chitnis, 1995). In vertebrates, this type of regulation occurs at a transcriptional level: blockade of Delta-Notch signalling by the dominant-negative

construct Δ^{dn} leads to a dramatic up-regulation of endogenous *Delta* expression, while forced overexpression of exogenous *Delta* has the opposite effect (Chitnis et al., 1995; Haddon et al., 1998). *Delta*-Notch-mediated cell-cell interactions with this type of feedback regulation of *Delta* are sufficient in principle to generate a regular spacing pattern of cell types (Collier et al., 1996).

The sensory bristles of *Drosophila* provide one of the best-characterised examples of lateral inhibition mediated by the Notch signalling pathway (Hartenstein and Posakony, 1990; Ghysen et al., 1993; Parks and Muskavitch, 1993; Jan and Jan, 1995; Zeng et al., 1998): the four cell types that form the bristle derive from a single sensory mother cell, and signalling via Notch, with *Delta* and the *Delta*-related protein *Serrate* acting in parallel as ligands, is necessary to cause them to adopt their different fates. In the accompanying paper (Adam et al., 1998), we point out a detailed parallel between insect bristles and the sensory patches of the vertebrate inner ear. In particular, we show in the chick embryo that *Delta*, *Serrate* and *Notch* homologues are expressed in the ear with a timing and pattern suggesting that they mediate lateral inhibition in the developing sensory patches: *C-Delta1* is expressed in scattered cells that appear to be nascent hair cells, as expected if lateral inhibition from each hair cell serves to force its neighbours to be supporting cells rather than hair cells. *C-Serrate1*, meanwhile, is expressed more uniformly within the sensory patch, and its product may act there as an additional ligand for Notch.

For further evidence that lateral inhibition indeed controls the differentiation of hair cells and supporting cells, we have turned to the ear of the zebrafish (Haddon and Lewis, 1996). We have previously identified four fish homologues of *Delta*, called *deltaA*, *deltaB*, *deltaC* and *deltaD*, and analysed their role in governing genesis of primary neurons in the early central nervous system: *deltaA*, *deltaB* and *deltaD* are all expressed strongly in the nascent neurons, all deliver lateral inhibition, and all are subject to feedback regulation of the type described above (Dornseifer et al., 1997; Appel and Eisen, 1998; Haddon et al., 1998). Because the ear develops relatively late, it is difficult to test the function of these genes in the ear by the simple method of RNA injection into an early blastomere. Zebrafish mutants, however, provide an alternative approach.

In both the Tübingen and the Boston zebrafish mutagenesis screens, mutant embryos were encountered that show a neurogenic phenotype, analogous to that seen in *Drosophila* embryos with mutations in the *Delta*-Notch signalling pathway (Jiang et al., 1996; Schier et al., 1996). The mutant gene was named *white tail* (*wit*) in Tübingen, and *mind bomb* (*mib*) in Boston, but complementation testing has shown that the mutations are allelic. *mind bomb* takes precedence as the official gene name. In the neural plate of *mib* mutants, neurons are produced in the normal regions, but in greatly excessive numbers, and contiguous with one another instead of being separated by intervening non-neuronal cells (Jiang et al., 1996); the pattern is similar to that seen in embryos where *Delta*-Notch signalling is blocked by injection of RNA coding for the dominant-negative *Delta* construct Δ^{dn} (Appel and Eisen, 1998; Haddon et al., 1998). *mib* mutations also cause abnormalities in other tissues where *Delta*-Notch signalling is thought to have a role, notably the somites, which likewise

show disturbances of segmentation that match those seen in embryos where *Delta*-Notch signalling is known to be blocked or defective (Conlon et al., 1995; Hrabé de Angelis et al., 1997; Jen et al., 1997). All these observations suggest that *mib* codes for a component that is required for *Delta*-Notch signalling.

If the *mib* mutation corresponds to a failure of *Delta*-Notch signalling, and *Delta*-Notch signalling is needed to control the hair-cell/supporting-cell decision, we should expect to see a disturbance in the ratio of hair cells to supporting cells in the sensory patches of the mutant ear. In this paper we test this prediction.

MATERIALS AND METHODS

Fish rearing and embryo culture

Zebrafish embryos were obtained by natural spawnings and maintained at 28.5°C in system water. Several wild-type strains were used, but comparisons between mutant and wild-type phenotypes are based in all cases on sibling embryos produced from matings between heterozygotes. The allele of *mind bomb* described in this paper, referred to throughout simply as *mib*, is the *ta52b* (*white tail*) allele from Tübingen (Jiang et al., 1996). It is the strongest of the five known alleles, as judged by the severity of the defects in pigmentation and somite segmentation, and may thus be a null mutation. Embryos were staged according to Kimmel et al. (1995); embryonic ages are given in hours post fertilization (hpf) at 28.5°C

Semi-thin resin sections

Embryos were fixed in BT fix (Westerfield, 1995) at 4°C overnight, embedded in Araldite (Agar Scientific Ltd), sectioned at 2 µm and stained with toluidine blue.

Cloning and sequencing of *Serrate* homologues

A fragment of a *Serrate* homologue, cloned by PCR as described by Haddon et al. (1998), was used to screen a 15- to 19-hour zebrafish λ ZAP II cDNA library (a gift from Bruce Appel), yielding clones corresponding to the zebrafish gene we have called *serrateB*. By probing the same library with a C-terminal fragment of the chick *C-Serrate1* gene (Myat et al., 1996), we discovered an additional zebrafish *Serrate* homologue, *serrateA*. cDNAs were excised from the vector using Stratagene's Rapid Excision Kit. Double-stranded sequencing was performed in both directions using an ABI PRISM system. Sequence comparisons were made using the Wisconsin GCG Gap program.

In situ hybridisation

Digoxigenin RNA antisense probes were made using Stratagene's RNA transcription kit. Whole-mount in situ hybridisation was essentially as described by Oxtoby and Jowett (1993); in situ hybridisation on cryosections followed Strähle et al. (1994) with minor modifications. Probes for *delta* genes were as described by Haddon et al. (1998). *serrateB* was linearised with *Xba*I and transcribed with T7.

Whole-mount phalloidin and antibody staining

To visualise and count hair bundles, fixed intact embryos were stained with FITC-phalloidin, and in some cases counterstained with the red fluorescent nuclear dye 7AAD, and viewed by confocal microscopy, as described by Haddon and Lewis (1996).

To identify neurons, whole-mount embryos were stained with anti-*islet-1* antibody (1:500) with HRP detection, as in Hammerschmidt and Nüsslein-Volhard (1993). This antibody recognises a LIM homeobox-containing nuclear protein expressed by early differentiating neurons (Korzh et al., 1993).

Mounting and imaging

For photography, live embryos were anaesthetised in 0.2–0.5 mM MS222 (Sigma) and immobilised in 1.2% low-melting point agarose (BioRad). Fixed and stained embryos were mounted in glycerol under a coverslip supported at its corners by high-vacuum grease, or were dehydrated in methanol and cleared in 2:1 benzyl benzoate: benzyl alcohol solution before mounting. Images were edited using Adobe Photoshop.

Single-cell labelling and fate mapping

Cells within the ear were labelled by iontophoretic injection. Clark 1.2 mm glass capillaries with an internal filament were pulled to make injection needles. Needles were back-filled by capillary action with lysinated rhodamine dextran (M_r 10 \times 10³; Molecular Probes; 5% in 200 mM KCl). Embryos were anaesthetised and mounted in low-melting point agarose, surrounded by Ringer's saline. The needle was positioned using a micromanipulator whilst monitoring with fluorescence optics, until contact was made in the ear epithelium. An oscillating voltage was then applied to the needle using a microelectrode amplifier (BioLogic); any cell touching the needle tip became labelled. The position of labelled cells was immediately recorded using a confocal microscope (BioRad MRC600). Embryos were then released and re-mounted in agarose 24 hours later, and again after 48 hours, to determine the location of the labelled cell or cells, using the confocal microscope as before.

RESULTS

Expression of *delta* genes foreshadows hair-cell differentiation in the normal ear

Differentiated hair cells, identifiable by their stereociliary bundles, are first seen in the zebrafish ear at about 24 hours post fertilization (hpf); they arise in two small patches, at the anterior and posterior ends of the otocyst respectively (Haddon and Lewis, 1996; Riley et al., 1997). These patches, corresponding to the maculae of the future utricle and saccule, then enlarge by addition of new hair cells and supporting cells, a process that probably continues throughout life (Platt, 1993; Haddon and Lewis, 1996).

We have used in situ hybridisation to examine how hair-cell production is related to the expression of *deltaA*, *deltaB* and *deltaD* in the ear epithelium. The expression patterns are similar for all three genes. Already at the 10-somite stage (14

hpf), when the otic placode is just becoming visible, expression is seen in two small groups of cells at its anteromedial and posteromedial ends, and this pattern persists over the next few hours as the placode becomes converted into a vesicle (Fig. 1, upper panels).

To discover the fate of cells at the early sites of *delta* expression in the ear, iontophoretic injections of fluorescent dextran were made into cells at these two sites, and the positions of the labelled cells were followed by confocal microscopy over the next 48 hours, as shown in Fig. 2. As the ear enlarges, cells at the labelled sites become positioned within the two sensory maculae: anteromedial cells end up in the anteroventral (utricle) macula, posteromedial cells end up in the posteromedial (saccular) macula.


In situ hybridization on sections through the ear at later stages reveals scattered *delta*-expressing cells in the thickened ventral areas of epithelium corresponding to the developing sensory patches (Fig. 3). Thus the pattern of expression of *delta* genes foreshadows both the pattern of production of the very first hair cells, and that of the hair cells that are added later, suggesting that the cells expressing the genes in the sites described above may be prospective hair cells. In other systems, *delta* expression is transient, switching off as overt differentiation begins (Muskavitch, 1994; Chitnis et al., 1995; Henrique et al., 1995; Adam et al., 1998). The same seems to be true here: expression is generally not seen in the mature differentiated regions of the developing sensory patches, although *deltaB*-expressing cells in the hair-cell layer are encountered occasionally (Fig. 3C) (similarly, *deltaB* persists for a little while in differentiating neurons; Haddon et al., 1998).

Expression of *serrateB* at first resembles that of the *delta* genes and then becomes restricted to the hair-cell layer

In chick and rodent embryos, the developing sensory patches in the ear are sites of expression of homologues not only of *Delta*, but also of the *Delta*-related gene *Serrate* (Hayashi et al., 1996; Lindsell et al., 1996; Myat et al., 1996). Is the same true in the zebrafish?

We have found two zebrafish *Serrate* homologues (see Materials and Methods) which we call *serrateA* (*serA*) and

Fig. 1. Expression of *deltaA*, *deltaB*, *deltaD* and *serrateB* in wild-type embryos (WT, top row) and *mib* mutant embryos (bottom row) at placode (14 hpf) and early otocyst (18 hpf) stages of ear development, shown by whole-mount in situ hybridization. The photographs are dorsal views of the ear rudiments and the hindbrain region between them; arrows indicate sites of *delta* and *serrate* gene expression at the anterior and posterior ends of the otic placode/otocyst. Anterior is to the left. Note the intensified and expanded expression of the genes at these sites in *mib* mutants – most obvious for *deltaD* and *serrateB*, but evident also for *deltaA* and *deltaB*. The band of *deltaD*-expressing cells very strikingly seen in the 14 hpf *mib* specimen extending between the anterior and posterior poles of the otocyst may correspond to the region from which neuroblasts will later delaminate.






Fig. 2. Tracing cell fate by labelling with lysinated rhodamine dextran (LRDX). In this example, a single cell injected with LRDX at the anterior site of *delta* gene expression in an early otocyst (18 hpf; A) ends up in the anteroventral macula, where it gives rise to a hair cell plus a supporting cell (identified by their shape and location in the epithelium; B and C). The pictures are superpositions of fluorescence and transmission-optics images taken with a confocal microscope in the living embryo at the time of injection and 24 and 48 hours afterwards. The specimen shown is one of approximately 80 that were used to build up a fate map (Haddon, 1997). In other specimens, labelled cells at the same original site end up in the same location. ao, anterior otolith; po, posterior otolith; am, anterior macula; hc, hair cell.

serrateB (*serB*). We have determined the full cDNA sequence of one of these, *serrateB* (GenBank accession number AF090432) and analysed its expression pattern. In amino-acid sequence, SerrateB appears slightly more closely related to chick Serrate2 (63% identity) than to chick Serrate1 (55% identity). Preliminary comparisons, based on partial sequence data, indicate that SerrateA may be more closely related to Serrate1.

At early stages of ear development, before overt differentiation of hair cells, the expression pattern of *serrateB* appears similar to that of *deltaA*, *deltaB* and *deltaD*: there are two patches of expression, at the anterior and posterior ends of the ear rudiment (Fig. 1). At later stages, however, up to at least 4 days (Fig. 3), expression of *serrateB*, in contrast with *delta* gene expression, persists in the differentiated hair cells.


Overexpression of *delta* genes in *mib* indicates a failure of Delta-Notch signalling

If the *mib* mutation causes a failure of Delta-Notch signalling, as hypothesized, we might expect to see abnormalities in the pattern of *delta* gene

Fig. 3. (A,B) Expression of *deltaA*, *deltaB* and *deltaD* in wild-type ears at 30 and 42 hpf, shown by in situ hybridisation on cryosections cut transverse to the main body axis. Arrows indicate scattered cells expressing *delta* genes in the thickened ventral wall of the otocyst. Beneath this epithelium, in the 30 hpf *deltaA* and *deltaB* sections, one can see *delta* gene expression in cells of the statoacoustic ganglion. (C) Expression of *deltaB* in a cell in the hair-cell layer of a sensory patch (the anteroventral macula) at 42 hpf; such *deltaB*-expressing hair cells are few and far between, because expression of the gene fades rapidly as a hair cell matures. (D) Persistent and extensive expression of *serrateB*, seen at 4 days in the hair-cell layer of the posteromedial and anteroventral maculae (arrows).

expression, because of the feedback regulation of *delta* genes, discussed in the Introduction: we know in the central nervous system, at least, that an artificial blockade of activity in the Notch pathway by exogenous dominant-negative Delta leads to a striking increase of endogenous *delta* expression (Haddon et al., 1998). Fig. 1 shows that in the early *mib* ear rudiment the *delta* genes are indeed expressed more intensely and in an increased number of cells. *serrateB* likewise shows increased expression (Fig. 1), strongly suggesting that it too is normally subject to inhibition by Notch activity.

The same figure also shows, incidentally, a similar upregulation of *deltaA* and *deltaB* expression in the *mib* neural tube, and the same phenomenon is seen for all three genes at earlier stages in the neural plate, during genesis of primary neurons (Fig. 4). This is further strong evidence that *mib* has a defect in the Delta-Notch signalling pathway. In wild-type embryos, artificial overexpression of *deltaA*, *B* or *D* inhibits production of primary neurons (Dornseifer et al., 1997; Appel and Eisen, 1998; Haddon et al., 1998); in *mib*, all three genes are overexpressed, and yet primary neurons are produced in excess (Jiang et al., 1996; Schier et al., 1996). The *mib* cells therefore must be refractory to the effects of the *delta* gene products.






Fig. 4. Expression of *deltaA*, *B* and *D* in the neural plate in wild-type and *mib* embryos at 12 hpf (6-7 somites); a part of the trunk region is shown in dorsal view, as in Fig. 1. The mutation causes increased *delta* expression but also increased numbers of primary neurons, showing that the mutant cells are refractory to the inhibitory effect of Delta-Notch signalling.

In *mib*, hair cells are produced in great excess at the expense of supporting cells

The *mib* ear shows several abnormalities in its gross structure, as noted by Malicki et al. (1996). At 60 hpf, the ear is smaller than normal, the semicircular canal system is imperfectly formed, and the two otoliths have failed to enlarge (Fig. 5A,B). These appear as tiny granules, similar to those seen in the 24-hpf ear in association with the first-formed hair cells. Since the otoliths overlie the sensory maculae and are thought to be created, in part at least, from the secretions of supporting cells in the maculae (Fermin and Igarashi, 1986; Riley and Grunwald, 1996), this defect hints at a defect in the sensory patches themselves.

To reveal the underlying pattern of cell differentiation in the sensory patches, we examined the distribution of hair cells and supporting cells by labelling with fluorescent phalloidin (to reveal the actin-rich stereocilia) and anti-acetylated-tubulin (to reveal both the kinocilium and the hair cell cytoplasm; data not shown). In addition, some specimens were fixed, embedded in resin and analysed by conventional semi-thin sectioning.

In the wild-type zebrafish, hair-cell nuclei are found in the apical half of the thick sensory epithelium, sitting above a layer of supporting-cell nuclei, as shown in Fig. 5C,E,G for specimens at 28-48 hpf. The stereociliary bundles of the hair cells can be seen projecting into the ear lumen towards the overlying otolith. In *mib* mutants, the pattern is radically altered (Fig. 5D,F,H). All the cell nuclei in the sensory patch lie in the same layer, and all of them belong to hair cells, clearly identifiable by their hair bundles. Supporting cells are absent, and the number of hair cells is very much larger than normal. Thus hair cells appear to have been produced at the expense of supporting cells, just as primary neurons are overproduced at the expense of other cell types in the *mib* CNS.




Fig. 5. Wild-type and *mib* ears compared. (A,B) Lateral views of live embryos (Nomarski optics) at 60 hpf. Note that the ears are smaller and deformed in *mib* and that the otoliths are drastically reduced, reflecting a deficit in the sensory patches. (C-D) Semi-thin Araldite sections stained with toluidine blue, showing the altered pattern of cell differentiation. (E-H) Confocal images of whole-mount specimens stained with fluorescent phalloidin (green) to reveal actin-rich hair bundles and with 7AAD (red nuclear staining) as a counterstain. (E,F) Optical sections passing transversely through sensory patches; (G,H) En-face views of sensory patches. Supporting cells are missing in *mib*, while hair cells are present in great excess. (I,J) Stages in the extrusion of hair cells from the otic epithelium in *mib*; staining is as in E-H. The hair cells eventually degenerate and disappear. hc, hair cell; sc, supporting cell; o, otolith; hb, hair bundle; n, neuron.

To assess precisely how many extra hair cells develop in the mutant and over what time period, the numbers of hair bundles (seen by phalloidin staining) were counted (Fig. 6). In the wild




Fig. 6. Total numbers of macular hair cells in wild-type and *mib* mutant ears as a function of age. Error bars represent standard error of the mean. Since cristae never develop in *mib*, the hair cells of cristae are not included in the total of hair cells in the wild-type ear.

type, hair cell numbers increase rapidly at first, reaching a combined total for the two maculae of 30–40 at 48 hpf, and then rise more slowly, at a mean rate of about 11 hair cells per day. In *mib*, the early production is dramatically increased, so that by 36 hpf there are already an average of 110 – fifteen times as many as in the wild type at the same stage. By 48 hpf, however, the numbers in *mib* have begun to drop, and by 60 hpf there are no hair cells left in the ear epithelium. The process leading to their loss is shown in Fig. 5I,J: the mutant patches of hair cells bulge out from the ear epithelium into the underlying mesenchyme, pinch off from the ear vesicle and eventually disappear, presumably being devoured by macrophages. Hair cells without supporting cells evidently cannot survive for long.

In a normal embryo, additional patches of hair cells appear later, at about 60 hpf, forming the cristae. These may originate from the same ventral regions of thickened prospective sensory epithelium that contain the early rudiments of the utricular and saccular maculae (Iwasaki, 1937; Becerra and Anadón, 1993; Haddon and Lewis, 1996). In the *mib* ear, no such late-developing hair cells are seen: from 60 hpf onwards, hair cells are completely lacking, suggesting that the initial burst of overproduction has exhausted the supply of precursors for production of hair cells at later stages.

The number of neurons in the statoacoustic ganglion is doubled in *mib*

Our studies in the chick have suggested that Delta-Notch signalling may regulate not only the hair-cell/supporting-cell decision, but also the segregation of neuroblasts from the otic epithelium and the production of neurons from these neuroblasts in the statoacoustic ganglion. In normal zebrafish embryos, neuroblasts appear to delaminate from the ventral wall of the otocyst between 22 and 36 hours after fertilization, and differentiated neurons begin to be visible in the ganglion by 24–30 hours (Haddon and Lewis, 1996). Thus, in contrast with the chick, neurogenesis and sensory-patch differentiation




Fig. 7. Dorsal view of ears plus hindbrain at 26 and 36 hpf, stained with islet-1 antibody using HRP detection, showing the statoacoustic ganglion (arrows) containing many more neurons in *mib* than in wild type. The outline of the right ear is indicated by a dotted line. The statoacoustic ganglion at early stages forms a continuous mass with the anterior lateral line ganglion and the VIIIth cranial nerve ganglion. As a rough guide, however, one can take the statoacoustic ganglion cells to be those that lie at and posterior to the anterior end of the otocyst; we followed this rule in making the counts described in the text.

occur concurrently in the zebrafish. The neurogenic region of the epithelium seems to overlap partially with the sites of the sensory patches, although the precise relationship remains to be determined. As in the chick, neurogenesis is associated with *delta* gene expression, which is visible both in regions of the otocyst that may correspond to sites of neurogenesis (see Fig. 1, legend) and in the developing statoacoustic ganglion (see Fig. 3A).

To see whether this pattern of gene expression reflects a role for Delta-Notch signalling in regulating otic neurogenesis, we have compared the numbers of neurons produced in wild-type and *mib* ears, using islet1 antibody staining as a neuronal marker (Fig. 7). For wild-type embryos at 26 hpf, we estimate from whole mounts that the number of islet-1-positive cells in the statoacoustic ganglion is 19.7 ± 6.5 (mean \pm s.d., $n=22$ ears); for *mib* embryos at that stage, it is 40.3 ± 6.2 (mean \pm s.d., $n=18$ ears). This twofold early excess in *mib* strongly suggests that normal otic neurogenesis is regulated at an early step by lateral inhibition, in agreement with the hypothesis.

DISCUSSION

We have argued that the *mib* mutation blocks Delta-Notch signalling, making cells deaf to lateral inhibition. The molecular nature of the *mib* gene remains to be discovered: genetic linkage experiments have so far allowed us only to exclude a number of possibilities, including the known zebrafish *Notch* homologues (Y.-J. J., unpublished). Meanwhile, the mutant provides a way to test the role of the Notch signalling pathway in the various tissues of the body. In

particular, it allows us to test the hypothesis that lateral inhibition mediated by Delta-Notch signalling controls production of the fine-grained pattern of hair cells and supporting cells in the sensory patches of the ear (Corwin et al., 1991; Lewis, 1991; Adam et al., 1998).

In *mib* mutants, all the cells in each patch differentiate as hair cells, indicating that Delta-Notch signalling is needed in normal development to force the neighbours of each hair cell to be supporting cells rather than hair cells. The absence of late-developing hair cells in the mutant suggests that Delta-Notch signalling is also needed to keep some cells competent to act as precursors for future hair cells, i.e. to maintain the sensory-patch stem-cell population that normally would persist to generate hair cells throughout life. This role is similar to the one we have demonstrated for Delta-Notch signalling in the chick retina (Henrique et al., 1997).

A caveat must, however, be stated. Homologues of the alternative Notch ligand *Serrate* are also expressed in the developing sensory patches of the ear, and the *mib* mutation may well affect their action as well as that of Delta. The misregulation of *serrateB* in *mib* (see Fig. 1) strengthens the suggestion that it may have a role in the control of hair cell differentiation in conjunction with the *delta* genes. Its persistence in the differentiated hair cells raises the possibility that it might play a part in stabilising the mature pattern of differentiation by keeping the supporting cells inhibited after *delta* expression has faded. Although the individual functions of the various *Delta* and *Serrate* homologues remain to be disentangled, the present findings further strengthen the parallel with bristle development in *Drosophila*, where signalling via Notch is required for cell diversification and Delta and Serrate act in parallel to deliver the signal (Zeng et al., 1998). It seems that information from *Drosophila* can indeed give insights into the development of the ear.

The finding that Notch signalling is necessary for genesis of the mixture of hair cells and supporting cells does not, of course, exclude a role for other factors in controlling the pattern. Molecules such as Numb, distributed unequally through asymmetric cell division, can bias the outcome of Delta-Notch-mediated interactions (Guo et al., 1996); cell movements can refine the spatial arrangement of the two cell types once they have been generated (Goodyear and Richardson, 1997).

The abnormalities in the gross structure of the *mib* ear may reflect additional roles for Notch signalling in the ear, or may be secondary to the defect in the sensory patches. The sensory patches are, for example, sites of expression of signalling molecules of the BMP family, which may be required for normal growth and global patterning (Oh et al., 1996; Takemura et al., 1996). It is also possible that the early reduction in the size of the otic vesicle in *mib* could reflect overproduction of delaminating neuroblasts at the expense of epithelial cells.

We have found a twofold increase in neuron production in the *mib* ear (see Fig. 7), but the precise role of Delta-Notch signalling in zebrafish ear neurogenesis remains to be clarified. In *Drosophila* sensory bristle development, extreme loss-of-function mutations in the Notch signalling pathway cause all the progeny of each sensory mother cell to develop as neurons, and no bristle shaft cells are produced (Hartenstein and Posakony, 1990). If neurons and hair cells in the fish ear derive

in a similar way from a common ancestor, one might expect that in *mib* there should be a massive overproduction of neurons and no production of hair cells. There are many possible reasons why we do not see this. Neuroblasts and sensory epithelial cells may, for example, have separate origins in the otocyst; or their segregation, though dependent on Delta-Notch signalling, may be only mildly affected by the *mib* mutation, because of genetic redundancy.

Our findings in the ear are not only of interest from the point of view of ear development. Intimate mixtures of cells of contrasting types are produced from the progeny of common precursors in a wide range of vertebrate tissues, from the gut lining to the bone marrow, and it is still a mystery how such mixtures are generated. Homologues of Notch and its ligands are, however, known to be expressed in many of these sites (Gridley, 1997). The sensory patches of the ear provide one of the few examples in the vertebrate body where we can now point with reasonable confidence to a mechanism for driving such fine-grained patterns of cell diversification. The insights we gain from the ear may thus help us to understand other tissues.

We thank Jenny Corrigan for cryosectioning, Suresh Jesuthasan for advice on microinjection, Vladimir Korzh for the gift of islet-1 antibody, Stephen Massey and Tobias Simmonds for looking after the fish, David Ish-Horowicz and Alastair Morrison for comments on the manuscript, and past and present members of the lab for discussions. The work was supported by the Imperial Cancer Research Fund and an EMBO fellowship to Y.-J. J.

REFERENCES

- Adam, J., Myat, A., Le Roux, I., Eddison, M., Henrique, D., Ish-Horowicz, D. and Lewis, J. (1998). Cell fate choices and the expression of Notch, Delta and Serrate homologues in the chick inner ear: parallels with *Drosophila* sense-organ development. *Development* **125**, 4645-4654.
- Appel, B. and Eisen, J. S. (1998). Regulation of neuronal specification in the zebrafish spinal cord by Delta function. *Development* **125**, 371-380.
- Artavanis-Tsakonas, S., Matsuno, K. and Fortini, M. E. (1995). Notch signalling. *Science* **267**, 225-232.
- Becerra, M. and Anadón, R. (1993). Development of the inner ear of the brown trout (*Salmo trutta fario*): I. Gross morphology and sensory cell proliferation. *J. Morphol.* **216**, 209-223.
- Campos-Ortega, J. A. (1993). Early neurogenesis in *Drosophila melanogaster*. In *The Development of Drosophila melanogaster* (ed. M. Bate and A. Martínez-Arías), Vol. 2, pp. 1091-1129. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Chitnis, A., Henrique, D., Lewis, J., Ish-Horowicz, D. and Kintner, C. (1995). Primary neurogenesis in *Xenopus* embryos regulated by a homologue of the *Drosophila* neurogenic gene *Delta*. *Nature* **375**, 761-766.
- Chitnis, A. B. (1995). The role of Notch in lateral inhibition and cell fate specification. *Mol. Cell. Neurosci.* **6**, 311-321.
- Collier, J. R., Monk, N. A. M., Maini, P. K. and Lewis, J. H. (1996). Pattern formation by lateral inhibition with feedback: a mathematical model of Delta-Notch intercellular signalling. *J. Theor. Biol.* **183**, 429-446.
- Conlon, R. A., Reaume, A. G. and Rossant, J. (1995). Notch1 is required for the coordinate segmentation of somites. *Development* **121**, 1533-1545.
- Corwin, J. T., Jones, J. E., Katayama, A., Kelley, M. W. and Warchol, M. E. (1991). Hair cell regeneration: the identities of progenitor cells, potential triggers and instructive cues. *Ciba Found. Symp.* **160**, 103-130.
- Dornseifer, P., Takke, C. and Campos-Ortega, J. A. (1997). Overexpression of a zebrafish homologue of the *Drosophila* neurogenic gene *Delta* perturbs differentiation of primary neurons and somite development. *Mech. Dev.* **63**, 159-171.
- Fermin, C. D. and Igarashi, M. (1986). Review of statoconia formation in birds and original research in chicks (*Gallus domesticus*). *Scan. Electron Microsc.* **4**, 1649-1665.

- Ghysen, A., Dambly-Chaudiere, C., Jan, L. Y. and Jan, Y.-N. (1993). Cell interactions and gene interactions in peripheral neurogenesis. *Genes Dev.* **7**, 723-733.
- Goodyear, R. and Richardson, G. (1997). Pattern formation in the basilar papilla: evidence for cell rearrangement. *J. Neurosci.* **17**, 6289-6301.
- Gridley, T. (1997). Notch signaling in vertebrate development and disease. *Mol. Cell. Neurosci.* **9**, 103-8.
- Guo, M., Jan, L. Y. and Jan, Y. N. (1996). Control of daughter cell fates during asymmetric division: interaction of Numb and Notch. *Neuron* **17**, 27-41.
- Haddon, C. and Lewis, J. (1996). Early ear development in the embryo of the zebrafish, *Danio rerio*. *J. Comp. Neurol.* **365**, 113-128.
- Haddon, C., Smithers, L., Schneider-Maunoury, S., Coche, T., Henrique, D. and Lewis, J. (1998). Multiple *delta* genes and lateral inhibition in zebrafish primary neurogenesis. *Development* **125**, 359-370.
- Haddon, C. M. (1997). *The Development of the Zebrafish Ear and a Quest for Genes Involved in Sensory Patterning* (Ph. D. Thesis, Open University).
- Hammerschmidt, M. and Nüsslein-Volhard, C. (1993). The expression of a zebrafish gene homologous to *Drosophila snail* suggests a conserved function in invertebrate and vertebrate gastrulation. *Development* **119**, 1107-1118.
- Hartenstein, V. and Posakony, J. W. (1990). A dual function of the *Notch* gene in *Drosophila* sensillum development. *Dev. Biol.* **142**, 13-30.
- Hayashi, H., Mochii, M., Kodama, R., Hamada, Y., Mizuno, N., Eguchi, G. and Tachi, C. (1996). Isolation of a novel chick homolog of *Serrate* and its coexpression with *C-Notch-1* in chick development. *Int. J. Dev. Biol.* **40**, 1089-1096.
- Heitzler, P. and Simpson, P. (1991). The choice of cell fate in the epidermis of *Drosophila*. *Cell* **64**, 1083-1092.
- Henrique, D., Adam, J., Myat, A., Chitnis, A., Lewis, J. and Ish-Horowitz, D. (1995). Expression of a *Delta* homologue in prospective neurons in the chick. *Nature* **375**, 787-790.
- Henrique, D., Hirsinger, E., Adam, J., Le Roux, L., Pourquie, O., Ish-Horowitz, D. and Lewis, J. (1997). Maintenance of neuroepithelial progenitor cells by *Delta*-*Notch* signaling in the embryonic chick retina. *Curr. Biol.* **7**, 661-670.
- Hrabé de Angelis, M., McIntyre, J. and Gossler, A. (1997). Maintenance of somite borders in mice requires the *Delta* homologue *DIII*. *Nature* **386**, 717-721.
- Iwasaki, I. (1937). Entwicklungsgeschichtliche Untersuchungen über das häutige Labyrinth der Knochenfische. *Jap. J. Med. Sci. I. (Anat.)* **6**, 301-419.
- Jan, Y. N. and Jan, L. Y. (1995). Maggot's hair and bug's eye: role of cell interactions and intrinsic factors in cell fate specification. *Neuron* **14**, 1-5.
- Jen, W. C., Wettstein, D., Turner, D., Chitnis, A. and Kintner, C. (1997). The *Notch* ligand, X-Delta-2, mediates segmentation of the paraxial mesoderm in *Xenopus* embryos. *Development* **124**, 1169-1178.
- Jiang, Y.-J., Brand, M., Heisenberg, C.-P., Beuchle, D., Furutani-Seiki, M., Kelsh, R. N., Warga, R. M., Granato, M., Haffter, P., Hammerschmidt, M., et al. (1996). Mutations affecting neurogenesis and brain morphology in the zebrafish, *Danio rerio*. *Development* **123**, 205-216.
- Jones, J. E. and Corwin, J. T. (1996). Regeneration of sensory cells after laser ablation in the lateral line system: hair cell lineage and macrophage behavior revealed by time-lapse video microscopy. *J. Neurosci.* **16**, 649-62.
- Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F. (1995). Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203**, 253-310.
- Korzh, V., Edlund, T. and Thor, S. (1993). Zebrafish primary neurons initiate expression of the LIM homeodomain protein *Isl-1* at the end of gastrulation. *Development* **118**, 417-425.
- Lewis, J. (1991). Rules for the production of sensory cells. *Regeneration of Vertebrate Sensory Cells/Ciba Symp* **160**, 25-39.
- Lewis, J. (1996). Neurogenic genes and vertebrate neurogenesis. *Curr. Opin. Neurobiol.* **6**, 3-10.
- Lindsell, C. E., Boulter, J., diSibio, G., Gossler, A. and Weinmaster, G. (1996). Expression patterns of *Jagged*, *Delta1*, *Notch1*, *Notch2*, and *Notch3* genes identify ligand-receptor pairs that may function in neural development. *Mol. Cell. Neurosci.* **8**, 14-27.
- Malicki, J., Schier, A. F., Solnica-Krezel, L., Stemple, D. L., Neuhaus, S. C. F., Stainier, D. Y. R., Abdelilah, S., Rangini, Z., Zwartkruis, F. and Driever, W. (1996). Mutations affecting development of the zebrafish ear. *Development* **123**, 275-283.
- Muskavitch, M. A. (1994). *Delta*-*Notch* signaling and *Drosophila* cell fate choice. *Dev. Biol.* **166**, 415-430.
- Myat, A., Henrique, D., Ish-Horowitz, D. and Lewis, J. (1996). A chick homologue of *Serrate* and its relationship with *Notch* and *Delta* homologues during central neurogenesis. *Dev. Biol.* **174**, 233-247.
- Oh, S. H., Johnson, R. and Wu, D. K. (1996). Differential expression of bone morphogenetic proteins in the developing vestibular and auditory sensory organs. *J. Neurosci.* **16**, 6463-6475.
- Oxtoby, E. and Jowett, T. (1993). Cloning of the zebrafish *krox-20* gene (*krx-20*) and its expression during hindbrain development. *Nucleic Acids Res.* **21**, 1087-1095.
- Parks, A. L. and Muskavitch, M. A. (1993). *Delta* function is required for bristle organ determination and morphogenesis in *Drosophila*. *Dev. Biol.* **157**, 484-496.
- Platt, C. (1993). Zebrafish inner ear sensory surfaces are similar to those in goldfish. *Hearing Res.* **65**, 133-140.
- Riley, B. B. and Grunwald, D. J. (1996). A mutation in zebrafish affecting a localized cellular function required for normal ear development. *Dev. Biol.* **179**, 427-435.
- Riley, B. B., Zhu, C., Janetopoulos, C. and Auderheide, K. J. (1997). A critical period of ear development controlled by distinct populations of ciliated cells in the zebrafish. *Dev. Biol.* **191**, 191-201.
- Schier, A. F., Neuhaus, S. C. F., Harvey, M., Malicki, J., Solnica-Krezel, L., Stainier, D. Y. R., Zwartkruis, F., Abdelilah, S., Stemple, D. L., Rangini, Z., Yang, H. and Driever, W. (1996). Mutations affecting the development of the embryonic zebrafish brain. *Development* **123**, 165-178.
- Sternberg, P. W. (1993). Falling off the knife edge. *Curr. Biol.* **3**, 763-765.
- Stone, J. S. and Cotanche, D. A. (1994). Identification of the timing of S phase and the patterns of cell proliferation during hair cell regeneration in the chick cochlea. *J. Comp. Neurol.* **341**, 50-67.
- Strähle, U., Blader, P., Adam, J. and Ingham, P. W. (1994). A simple and efficient procedure for non-isotopic *in situ* hybridisation to sectioned material. *Trends Genet.* **10**, 75-76.
- Takemura, T., Sakagami, M., Takebayashi, K., Umamoto, M., Nakase, T., Takaoka, K., Kubo, T., Kitamura, Y. and Nomura, S. (1996). Localization of bone morphogenetic protein-4 messenger-RNA in developing mouse cochlea. *Hearing Res.* **95**, 26-32.
- Weisleder, P., Tsue, T. T. and Rubel, E. W. (1995). Hair cell replacement in avian vestibular epithelium: supporting cell to type I hair cell. *Hearing Res.* **82**, 125-133.
- Westerfield, M. (1995). *The Zebrafish Book*. Eugene: University of Oregon Press.
- Wilkinson, H. A., Fitzgerald, K. and Greenwald, I. (1994). Reciprocal changes in expression of the receptor *lin-12* and its ligand *lag-2* prior to commitment in a *C. elegans* cell fate decision. *Cell* **79**, 1187-1198.
- Zeng, C., Younger-Shepherd, S., Jan, L. Y. and Jan, Y. N. (1998). *Delta* and *Serrate* are redundant *Notch* ligands required for asymmetric cell divisions within the *Drosophila* sensory organ lineage. *Genes Dev.* **12**, 1086-1091.