

Lsh controls silencing of the imprinted *Cdkn1c* gene

Tao Fan¹, John P. Hagan², Serguei V. Kozlov², Colin L. Stewart² and Kathrin Muegge^{1,*}

¹Laboratory of Molecular Immunoregulation, Basic Research Program, SAIC-Frederick, National Cancer Institute, Frederick, MD 21702-1201, USA

²Cancer and Developmental Biology Laboratory, National Cancer Institute, Frederick, MD 21702-1201, USA

*Author for correspondence (e-mail: muegge@ncifcrf.gov)

Accepted 26 November 2004

Development 132, 635–644

Published by The Company of Biologists 2005

doi:10.1242/dev.01612

Summary

Epigenetic regulation, such as DNA methylation plays an important role in the control of imprinting. Lsh, a member of the SNF2 family of chromatin remodeling proteins, controls DNA methylation in mice. To investigate whether Lsh affects imprinting, we examined CpG methylation and allelic expression of individual genes in Lsh-deficient embryos. We report here that loss of Lsh specifically alters expression of the *Cdkn1c* gene (also known as *p57^(Kip2)*) but does not interfere with maintenance of imprints at the *H19*, *Igf2*, *Igf2r*, *Zac1* and *Meg9* genes. The reactivation of the silenced paternal *Cdkn1c* allele correlates closely with a loss of CpG methylation at the 5' DMR at the *Cdkn1c* promoter, whereas KvDMR1 and DMRs of other imprinted

genes were not significantly changed. Chromatin immunoprecipitations demonstrate a direct association of Lsh with the 5' DMR at the *Cdkn1c* promoter, but not with Kv DMR1 or other imprinted loci. These data suggest that methylation of the 5' DMR plays an important role in the imprinting of the *Cdkn1c* gene. Furthermore, it suggests that Lsh is not required for maintenance of imprinting marks in general, but is only crucial for imprinting at distinct genomic sites.

Key words: Chromatin structure, DNA methylation, Gene imprinting, Lsh

Introduction

Imprinting is an epigenetic mechanism that regulates mono-allelic expression in a parent-of-origin-dependent manner (Bartolomei and Tilghman, 1997; Reik and Walter, 2001; Feinberg et al., 2002; Delaval and Feil, 2004). Epigenetic modifications of chromatin label imprinted loci in a heritable fashion to ensure expression of either the maternal or paternal allele of the gene. Loss of imprinting leads to bi-allelic expression or complete silencing of a gene. As a consequence of aberrant gene expression, abnormal embryogenesis can occur that is frequently associated with overgrowth or undersizing of fetal or extra-embryonic tissue. Loss of imprinting also plays an important role in a number of human diseases (such as Beckwith-Wiedemann, Prader-Willi or Angelman syndrome) and is thought to play a role in cancer development (Paulsen and Ferguson-Smith, 2001; Feinberg et al., 2002).

Chromatin modifications that ensure the inheritance of the imprinting pattern are first erased during gametocyte development and then gender-specifically established either at late foetal stages or also after birth in female germ cells (Constancia et al., 1998; Verona et al., 2003; Delaval and Feil, 2004). Epigenetic modifications are based on DNA methylation or histone tail modifications that regulate the accessibility of chromatin. A great deal of evidences points to CpG methylation as a major epigenetic modification that commands the parental identity. Imprinted genes are frequently associated with differentially methylated regions (DMRs) that are established during gametocyte development and maintained with a high fidelity throughout embryogenesis

(Constancia et al., 1998; Mann et al., 2000). Mutational analysis of DMRs has supported their crucial role in the mechanism of imprinting (Elson and Bartolomei, 1997; Wutz et al., 1997). CpG methylation, for example, can shape chromatin structure by influencing histone modifications such as acetylation or methylation levels (Bird, 2002). This in turn can alter the accessibility of transcription factors to their appropriate binding site and thus control gene expression. CpG methylation may also directly interfere with the recruitment of DNA binding factors to their target sites. For example, CpG methylation abolishes binding of CTCF to the ICR (imprinting control center) of *H19*, a differentially methylated region that controls *H19* and *Igf2* (insulin growth factor 2) gene expression (Burgess-Beusse et al., 2002). The ICR is located about 2 to 4 kb upstream of the *H19* gene (Tremblay et al., 1997). Binding of CTCF in turn affects the organization of a 'chromatin boundary' that blocks the interaction of a downstream enhancer with the *Igf2* promoter (Simpson et al., 2002). Ultimately, CTCF binding controls *H19* gene expression and suppresses *Igf2* gene transcription. The importance of CpG methylation for imprinting has also been demonstrated in *Dnmt1^{-/-}* embryos that display reduced CpG methylation and loss of imprinting at multiple genomic loci (Li et al., 1993; Caspary et al., 1998).

Lsh (lymphoid specific helicase), a member of the SNF2/helicase family of chromatin remodeling proteins is an epigenetic regulator in mice (Jarvis et al., 1996; Geiman et al., 1998; Yan et al., 2003a; Huang et al., 2004). Since Lsh regulates DNA methylation (Dennis et al., 2001), we tested whether changes in CpG methylation levels in *Lsh^{-/-}* mice can

influence the mechanisms of gene imprinting. As *Lsh*^{-/-} mice die at birth (Geiman et al., 2001), we examined the maintenance of gene imprints in *Lsh*^{-/-} embryos at day 17.5 of gestation. Among six analyzed imprinted loci, mono-allelic expression was only disturbed in the *Cdkn1c* (*p57*^{Kip2}) gene by deletion of *Lsh*. The bi-allelic expression of the *Cdkn1c* gene was accompanied by a substantial reduction of CpG methylation at the 5' DMR of the *Cdkn1c* gene. Furthermore, *Lsh* is specifically associated with the promoter of *Cdkn1c* in wild-type cells. These results imply independent control mechanisms for imprinted genes and suggest a crucial role for *Lsh* and DNA methylation to execute the imprinting epigenetic program in a locus-specific manner.

Materials and methods

Mice

To generate DNA polymorphisms at imprinted loci, *Lsh*^{+/-} 129/SvEv (Geiman et al., 2001) mice were crossed with *CzechIII/Ei* mice to produce *Lsh*^{+/-} F1 hybrid mice. The *Lsh*^{+/-} F1 hybrid mice were backcrossed with *Lsh*^{+/-} 129/SvEv mice to produce F2 hybrid embryos that were analyzed for allele polymorphism at imprinting loci. To investigate *Zac1* gene imprinting, *Zac1/lacZ* knock-in mice were used. Generation of *Zac1/lacZ* knock-in animals expressing the bacterial β -galactosidase cDNA under the transcription control of the endogenous *Zac1* promoter has been performed by conventional gene targeting methodology and will be described elsewhere (S.V.K. and C.L.S., unpublished). The mutant allele has been kept on a 129/SvEv mouse background and proven to maintain the imprinting status both *in vivo* and in cultured primary MEFs (S.V.K. and C.L.S., unpublished). We crossed *Lsh*^{+/-} mice and *Zac1/lacZ* mice to produce *Lsh*^{+/-}*lacZ*^{+/-} F1 mice. These were backcrossed with *Lsh*^{+/-} mice. The F2 hybrid offsprings were examined for β -galactosidase activity. *Lsh* genotyping was done as reported before (Geiman et al., 2001), while the presence of the *lacZ* gene was determined by using primers as indicated in Table 1A. Cycling conditions were 3 minutes at 94°C, then 30 seconds at 94°C, 30 seconds at 58°C and 1 minute at 72°C for 30 cycles followed by a final 7-minute extension step at 72°C.

Polymorphisms analysis

DNA was isolated from whole embryos after removal of the head and the internal organs. Parental origin of alleles of heterozygous individuals was discriminated by exploiting polymorphisms between two mouse strains. PCR primers were used as indicated in Table 1B. PCR products were purified with the QIAquick PCR purification kit (Qiagen), and then digested with the indicated restriction enzymes. The fragments were separated on 0.8-2.0% agarose gels for RFLP analysis to identify the parental origin of the alleles. Undigested PCR products and cleaved products derived from 129/SvEv and *CzechIII/Ei* mice were used as controls.

RT-PCR analysis

Total RNA was isolated from day 17.5 embryo body tissues using RNazol (Tel-Test) and the expression patterns of individual genes were analyzed and compared between *Lsh*^{+/+} and *Lsh*^{-/-} embryos. Residual genomic DNA was removed from total RNA with a DNase I (Invitrogen) treatment. Reverse transcription was performed on 2 μ g of total RNA using the Mu-MLV reverse transcriptase kit (Ambion). Control reactions were prepared in parallel without reverse transcriptase. The different transcripts were amplified from 1/40 of the reverse transcription reaction in the presence of each of the specific primers. RFLP analysis was done as described above.

Histochemical staining for β -galactosidase activity

We investigated β -galactosidase gene expression by X-Gal (5-bromo-

4-chloro-3-indolyl- β -D-galactopyranoside) staining. Day 13.5 embryos were removed and rinsed with PBS. Embryos were fixed in fixative (0.2% (v/v) glutaraldehyde, 1.5% (v/v) formaldehyde, 5 mM EGTA (pH 8.0), 2 mM MgCl₂, in PBS) for 30 minutes. The samples were washed three times in buffer (2 mM MgCl₂, 0.01% (w/v) sodium deoxycholate, 0.02% (v/v) Nonidet P-40, in PBS). Staining was carried out at 30°C in the dark for 16 hours in the following buffer (1 mg/ml X-Gal, 5 mM K₄Fe(CN)₆, 5 mM K₃Fe(CN)₆, in PBS). For staining of tissue sections, embryos were immediately fixed in liquid nitrogen; 5 μ m thick coronal sections were mounted on a set of gelatin-coated glass slides such that serial sections could be used for X-Gal stains.

Bisulphite mutagenesis assay

An aliquot of 1 μ g of genomic DNA was subjected to bisulphite treatment using CpGenome DNA modification kit (Intergen Co) according to the manufacturer's instructions. Primers were generated to match the DMRs of the imprinting gene *H19*, *Igf2r*, *Zac1* and *Cdkn1c*. The converted DNA was subjected to methylation-specific nested PCR, using the following primers (Table 1C) and conditions: for *H19* ICR (Tremblay et al., 1997); for the *Igf2r* gene (Lucifero et al., 2002); for *Cdkn1c* (Yatsuki et al., 2002). For outside primers the following PCR conditions were used: cycling conditions were 3 minutes at 94°C, then 30 seconds at 94°C, 30 seconds at 58°C and 1 minute at 72°C for 35 cycles followed by a final 7-minute extension step at 72°C. For use of inside primers, 30 cycles were used. The PCR products were separated in 1.5% agarose gels and purified using the QIAEX II gel extraction kit (Qiagen). Amplified fragments were subcloned into the pGEMT-Easy vector (Promega). Independent clones for each fragment were sequenced by using the T7 primer.

Chromatin immunoprecipitation assay

In order to investigate the interaction between *Lsh* protein and chromatin, we performed chromatin immunoprecipitation assays using a stable transfected 3T3 cell that expressed Flag-tagged *Lsh* under an inducible promoter (Yan et al., 2003b). Zeocin-resistant cells were induced with 100 pM mifepristone for induction of the *Lsh* protein. Following cross-linking with culture medium containing 1% formaldehyde at 37°C for 10 minutes and washing twice with ice-cold PBS containing protease inhibitors (1 mM phenylmethyl sulfonyl fluoride (PMSF), 1 μ g/ml aprotinin and 1 μ g/ml peptatin A) cells were scraped off the dishes and pelleted. The cells were resuspended in SDS lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris-HCl pH 8.1) with protease inhibitors for 10 minutes on ice and sonicated four times for 30 seconds each at a power setting of 3.0 with the Sonicator 3000 (MISONIX) to get 100-1000 bp DNA fragments. The sample was centrifuged to remove cell debris and diluted ten-fold in ChIP dilution buffer (0.01% SDS, 1.1% Triton X-100, 1.2 mM EDTA, 16.7 mM, Tris-HCl pH 8.1, 167 mM NaCl, protease inhibitors). The supernatants were pre-cleared with 80 μ l of a mixture of salmon sperm DNA-protein A agarose slurry (Upstate Biotechnology). The slurry solution was centrifuged and the supernatants were incubated with 2 μ l of Flag M2 antibody (Sigma) or murine IgG1 as isotype control with rotation overnight at 4°C. Then, 80 μ l of salmon sperm DNA-protein A agarose slurry was added and incubated for 1 hour. The beads were washed several times, and the attached immune complexes were eluted with a buffer containing 1% SDS and 0.1 M NaHCO₃, followed by reverse-crosslinking at 65°C for 4 hours. DNA was purified by proteinase K digestion, phenol-chloroform extraction and ethanol precipitation. DNA was resuspended in 100 μ l of 1 \times TE and 4 μ l were used for PCR analysis. The amplification profile was designed for 30 cycles using the same primers described in the DMR polymorphism analysis (Table 1D).

Table 1. Oligonucleotides used in this study

Gene name	Accession number	Primers	Polymorphism(129 →Czech)	Restriction enzyme (129→Czech)
A Genotyping				
<i>lacZ</i>	L08936	Forward 5'-gtctcgttgctgcataaac-3' Reverse 5'-tcgtctgctcatccatgacc-3'		
B Polymorphisms between 129Sv and Czech mice				
<i>H19</i>	NM_02123	Forward 5'-caaagcaccctgactctgtttcc-3' Reverse 5'-gggcatgtgaacactttatgatggaac-3'	1631A→G	No site→ <i>Bgl</i> I
<i>Igf2</i>	NM_010514	Forward 5'-tcagtaatcgatattgggatccccagtgggaa-3' Reverse 5'-ccagtcacgatactcactgatggtctggac-3'	1225C→T	<i>Dde</i> I→no site
<i>Igf2r</i>	L22109	Forward 5'-ttacactgatggtgatgactgtggcagtg-3' Reverse 5'-tggcagggcccgagtttgactgac-3'	41T→C	<i>Bgl</i> III→no site
<i>Cdkn1c</i>	AF160190	Forward 5'-aggagccgtccatcacaatcag-3' Reverse 5'-cagagacctgctcaggacctgttc-3'	1684A→G	No site→ <i>Tfi</i> I
<i>Meg9</i>	AK013406	Forward 5'-caggtgacacgctgaattgg-3' Reverse 5'-ggtgtgacagctcctcagg-3'	124A→T	<i>Nhe</i> I→no site
C Primers for bisulfite sequencing				
<i>H19</i>	U19619	Outside forward 5'-gagtatttaggagtataagaatt-3' Outside reverse 5'-atcaaaaactaacataaacct-3' Inside forward 5'-gtaaggagattatgttttttgg-3' Inside reverse 5'-cctcattaatcccataactat-3'		
<i>Igf2r</i>	L06446	Outside forward 5'-ttagtggggtattttattgtatgg-3' Inside forward 5'-gtgtggtttttatgtatgttagg-3' Reverse 5'-aaatctcaaaaatacaaaactacac-3'		
<i>Cdkn1c</i>	AP001293	Forward 5'-aggatttagttgtagtagt-3' Inside reverse 5'-tttcaattcaacaacacc-3' Outside reverse 5'-tatectatccaactaaacc-3'		
<i>KvDMR1(IGF8a)</i>	AP001295	Outside forward 5'-ggttttagttaggaaggtag-3' Inside forward 5'-ggatgaggaaggtaggtttt-3' Reverse 5'-ctaactaataataacctcacc-3'		
<i>KvDMR1(IGF8b)</i>	AP001295	Outside forward 5'-gtgtgattttttggagag-3' Inside forward 5'-taagtgagtggttaggat-3' Reverse 5'-aatccccacacctaaattc-3'		
<i>Zac1</i>	AJ308559	Forward 5'-attgtatttagttgggtggg-3' Inside reverse 5'-ccaaattcaaaattatcacctc-3' Outside reverse 5'-attctccaaaattcttaaaatc-3'		
D Primers for polymorphism analysis of DMR and/or chip assay				
<i>H19</i>	U19619	Forward 5'-aaggaacatgctacattcac-3' Reverse 5'-ctgagatagctcttgagaac-3'		
<i>Igf2r</i>	L06446	Forward 5'-gtgtggcaccctcatgcatag-3' Reverse 5'-aggtatcctgagggtgcaaac-3'		
<i>Cdkn1c</i>	AP001293	Forward 5'-cagccacggtactgccaggac-3' Reverse 5'-cgggacctcctcagattagc-3'		
<i>KvDMR1(IGF8a)</i>	AP001295	Forward 5'-agggatgaggaaggtaggcc-3' Reverse 5'-ctggctgatgacctctcc-3'		
<i>KvDMR1(IGF8b)</i>	AP001295	Forward 5'-caagtgagtggcctaggac-3' Reverse 5'-cacctgaattccgagtcggc-3'		
<i>Zac1</i>	AJ308559	Forward 5'-attgtatttagttgggtggg-3' Reverse 5'-ccaaattcaaaattatcacctc-3'		

Results

The *Zac1* imprinting pattern is maintained in the absence of Lsh


Previously, we reported a global loss of CpG methylation in all tissues derived from *Lsh*^{-/-} mice (Dennis et al., 2001). The overall decrease of DNA methylation in Lsh-deficient tissue is about 50% in comparison to wild-type DNA. In order to determine whether Lsh can affect gene imprinting, we used the *Zac1/lacZ* transgenic mouse model. The *Zac1* (*Plagl1* – Mouse Genome Informatics) gene encodes a zinc finger transcription factor and is paternally expressed (Spengler et al., 1997). In the *Zac1/lacZ* knock-in strain of mice, one allele of the imprinted *Zac1* gene had been deleted by homologous recombination and replaced with the *lacZ* gene. Thus breeding of *Lsh* heterozygous mice with the *Zac1/lacZ* heterozygous mice allowed distinguishing between the parental alleles at the *Zac1*

locus. If the *lacZ* allele is inherited from the paternal site, the *lacZ* gene is expressed, whereas propagation from the maternal site silences the *lacZ* gene. Since the *Zac1* gene is expressed in multiple adult and embryonic tissues (Piras et al., 2000), this transgenic mouse model allows examination of aberrant imprinting control in different tissues using histochemical staining for β-galactosidase activity.

As shown in Fig. 1A, β-galactosidase activity was detected in the tail of wild-type and Lsh heterozygous mice, when the *lacZ* allele was propagated from the paternal side. Similarly, the *Lsh*^{-/-} embryos displayed normal paternal expression pattern of the *lacZ* allele. When the *lacZ* allele was propagated from the maternal side, *lacZ* activity was completely suppressed in wild-type tails as well as in tissue derived from *Lsh*^{-/-} embryos. Thus silencing of the maternal allele was not relieved in the absence of Lsh. In order to determine if Lsh deletion possibly resulted in a 'leaky' phenotype and could

Fig. 2. Genomic DNA polymorphism analysis.

(A) Schematic representation indicating the distinct imprinted genes examined. The filled boxes represent exons. The open boxes represent the position of differentially methylated regions as examined in this study. The lines under the graph indicate the position of the primers used for genomic DNA polymorphism analysis. The arrows indicate the position of the genomic polymorphisms. (B) Genomic DNA from F2 hybrids was amplified at specific regions using primers presented in Table 1B. PCR products were digested with *Bgl*I for *H19*, *Dde*I for *Igf2*, *Bgl*II for *Igf2r*, *Nhe*I for *Meg9*, and *Tfi*I for *Cdkn1c*. Parent-of-origin alleles are distinguished by the size of the DNA fragments generated by digestion and visualized by ethidium bromide stain after agarose gel electrophoresis. Mice that were homozygous for 129 alleles and homozygous for Czech alleles served as controls. Only embryo #2 but not embryo #1 showed appropriate polymorphisms at the maternally expressed *H19*, *Igf2r*, *Meg9*, and *Cdkn1c* genes and at the paternally expressed *Igf2* gene.




of Lsh. Thus neither did the DMR2 of the *Igf2r* gene exhibit loss of DNA methylation, nor was any change in the *Igf2r* gene expression pattern detectable, suggesting that this imprinted locus was not affected by Lsh deletion.

Next, we characterized the methylation status of the 5' DMR of the *Zac1* gene. The 5' DMR, which is about 1580 bp in length, contains a 420 bp-long CpG island that spans part of the promoter region and the first exon of the *Zac1* gene (Smith et al., 2002). This island shows differential methylation on the maternal allele. In wild-type embryos, we detected clones that were fully methylated at the DMR or completely unmethylated in the ratio 1:1, representing the maternal and paternal alleles (Fig. 4). In *Lsh*^{-/-} samples, a few unmethylated CpG sites sporadically occurred without any recognizable pattern. Since the *Zac1* gene remained imprinted in all examined tissues of the *Lsh*^{-/-} mice, the data suggested that limited demethylation was not sufficient to activate the silent allele.

The mouse *Igf2* and *H19* genes are located 70 kb apart on chromosome 7, and demonstrate reciprocal imprinting status

(Fig. 2A). Only the paternal allele of the *Igf2* gene and only the maternal allele of the *H19* gene are expressed. A DMR upstream of the *H19* gene is essential for their parental allele-specific expression (Tremblay et al., 1997). This region contains conserved CTCF-binding sites involved in the establishment of a 'chromatin boundary' that regulates the imprinted expression of *Igf2* and *H19*. Recently, we studied DNA methylation of the H19 locus at a 3.8 kb region comprising the ICR, examining about ten methylation-

Fig. 3. Effect of Lsh on expression of different imprinted genes. RT-PCR analysis for the indicated genes was performed using total RNA derived from F2 hybrid offspring. PCR products were subjected to RFLP analysis by using restriction enzymes listed in Table 1. Omission of reverse transcriptase and uncut PCR products served as controls.




Fig. 4. Methylation analysis of DMRs at different imprinted loci in the absence of Lsh. The methylation status of individual DNA alleles derived from *Lsh*^{+/+} and *Lsh*^{-/-} embryos are represented in the graph. The gene fragments were amplified for *H19*, *Igf2r*, *Zac1* DMRs from sodium bisulphite-modified DNA and subjected to sequence analysis. (Top) Open boxes correspond to DMRs of each gene. The arrow indicates the transcription start site. Each row corresponds to an individual sequenced strand of DNA, and each circle represents a CpG on the strand, red circles and blue circles indicate methylated and unmethylated sites, respectively. Maternal (M) and paternal (P) alleles were distinguished by a DNA polymorphism in the DMR sequence (*I29/Czech*: *Igf2*, 1654A/G, U19619; *Igf2r*, 851C/G, L06446). *Parent-of-origin determined by the methylation pattern.

sensitive HhaI sites by Southern analysis. Although hypomethylation of some HhaI sites was observed, their precise location could not be determined based on the high density of HhaI sites. Therefore a conclusion as to whether CpG hypomethylation was indeed affecting CTCF binding sites was not possible. Thus we examined in this study a 400 bp-long region of the ICR that comprised two CTCF binding sites. (Note: Previous Southern analysis had not shown any evidence of hypomethylation at the single HhaI site located within this region.) Almost all 16 CpG sites were methylated on the paternal allele, whereas the maternal allele was completely unmethylated (Fig. 4). No significant differences in the methylation pattern of the paternal allele comparing wild-type and *Lsh*^{-/-} tissues were found. Thus the similar methylation pattern in *Lsh* wild-type and *Lsh*-deleted samples corresponded with the lack of expression changes in the *Igf2* and *H19* genes.

Next, we examined the methylation changes at the *Cdkn1c* gene that is maternally expressed (Hatada and Mukai, 1995) (Fig. 5). DNA methylation analysis had identified several DMRs in the locus (Yatsuki et al., 2002). The DMR (KvDMR1) of the *Kcnq1ot1* (*Lit1*) promoter is located about 150 kb downstream of the *Cdkn1c* gene. The KvDMR1 gets methylated in oocytes and is unmethylated in sperm, and may represent an imprinting mark in this domain. Loss of methylation and silencing of the *Cdkn1c* gene has been implicated in patients with Beckman-Wiedeman syndrome

(Diaz-Meyer et al., 2003). A recent study reported that deletion of the KvDMR1 in mice results in reactivation of the silenced paternal *Cdkn1c* allele (Fitzpatrick et al., 2002). Two sites within the KvDMR1 were analyzed by bisulphite sequencing (known as ICG8a and ICG8b) (Yatsuki et al., 2002), but no significant methylation differences were detectable comparing *Lsh*^{+/+} and *Lsh*^{-/-} samples. In contrast, the CpG methylation pattern of the ICG5 site (Yatsuki et al., 2002) was greatly altered in the absence of Lsh (Fig. 5). The ICG5 site is contained within a 5' DMR located in the promoter upstream of the *Cdkn1c* gene and is largely methylated on the paternal allele in wild-type controls. However, *Lsh*^{-/-} samples had lost dramatically more than half of the cytosine methylation. Thus the substantial decrease of CpG methylation in the 5' DMR at the paternal allele is closely associated with loss of paternal silencing and bi-allelic expression of the *Cdkn1c* gene.

Lsh is physically associated with the DMR of the *Cdkn1* gene

In order to understand why Lsh dramatically affects the methylation state of the *Cdkn1c* gene, we investigated the possibility that Lsh directly interacts with specific sites of the *Cdkn1c* gene. Using the chromatin immunoprecipitation assay (ChIPs), we examined binding of Lsh to the distinct differentially methylated regions that we had previously examined for CpG methylation by bisulphite mutagenesis. Nuclei were prepared from stably transfected 3T3 cells that




Fig. 5. Methylation analysis at three DMRs of the *Cdkn1c* locus in the absence of Lsh. The methylation status was determined as outlined in Fig. 4. The arrow indicates the transcription start site. Each row corresponds to an individual sequenced strand of DNA, and each circle represents a CpG on the strand, red circles and blue circles indicate methylated and unmethylated sites, respectively. Maternal (M) and paternal (P) alleles were distinguished by a DNA polymorphism in the DMR sequences (129/Czech: 5' DMR *Cdkn1c* (ICG5) 34012 T/G; KvDMR1: ICG8a 101803 A/G and ICG8b 103181 G/C).

could be induced to express high levels of Lsh as a Flag-tagged protein (Yan et al., 2003b). Chromatin was precipitated with anti-Flag antibody and examined by PCR amplification for the presence of DMRs of the *H19*, *Igf2r*, *Zac1* or the *Cdkn1c* gene. As shown in Fig. 6, only sequences from the DMR of *Cdkn1c* were detected in the Flag-tagged precipitates. Furthermore, only the 5' DMR of the *Cdkn1c* promoter region, but not the KvDMR1 were associated with Lsh.

These data suggest a direct role for Lsh in DNA methylation and chromatin structure in the regulation of imprinting at specific sites such as the *Cdkn1c* gene.

Discussion

Epigenetic modifications such as DNA methylation are important mechanisms for control of gene imprinting. In this study we have evaluated the role of Lsh, a major regulator of CpG methylation levels in mice, for its effect on gene

imprinting. We found that Lsh affects only selected genes such as the maternally expressed *Cdkn1c* gene but had no impact on the majority of imprinted loci analyzed (the paternally expressed *Zac1*, *Igf2* or maternally expressed *H19*, *Igf2r* and *Meg9* genes). Thus independent control mechanisms for genomic imprinting exist and Lsh participates in the control in a locus-specific manner. Since *Cdkn1c* is a cell cycle inhibitor, bi-allelic expression may be in part responsible for the growth retardation observed in the whole *Lsh*^{-/-} embryo, cultured lymphocytes and embryonal fibroblasts derived from *Lsh*^{-/-} embryos (Geiman and Muegge, 2000; Geiman et al., 2001; Fan et al., 2003).

Various evidence points to DNA methylation as an important mechanism in genomic imprinting. For example treatment of cell cultures with the demethylating drug azacytidine abolishes the imprint of several genes (El Kharroubi et al., 2001). In addition, deletion of the major 'maintenance' methyltransferase gene *Dnmt1* leads to global demethylation and a loss of imprinting at




Fig. 6. Binding of Lsh to the DMR of *Cdkn1c*. Chromatin immunoprecipitation assays were performed in 3T3 cells that had stably integrated the Flag-Lsh expression plasmids. Specific antibodies directed against the Flag-tagged form of Lsh and murine control antibodies of the same isotype were used. Aliquots of chromatin taken before immunoprecipitation were used as 'Input' controls. The immunoprecipitates obtained were PCR amplified using the primers specific to the DMRs of indicated genes.

many loci (Li et al., 1993; Caspary et al., 1998). The close association of CpG hypomethylation at the 5' DMR of the promoter region, and the de-repression of the silent paternal *Cdkn1c* gene as reported in this study give further support for a functional role of DNA methylation in imprinting.

Two differentially methylated regions have been identified in the *Cdkn1c* gene: KvDMR1 located in an intron within the *Kcnql* gene (about 150 kb downstream of the *Cdkn1c* gene) and 5' DMR located in the *Cdkn1c* promoter region. The KvDMR1 is methylated on the maternal allele, whereas the 5' DMR is methylated on the silenced paternal allele. Furthermore the KvDMR1 site is critical for repression of the silenced paternal allele, as deletion of the unmethylated paternal KvDMR1 results in bi-allelic expression of the embryo (Fitzpatrick et al., 2002). In contrast, deletion of the methylated KvDMR1 on the maternal allele had no effect on *Cdkn1c* gene expression. However, CpG methylation must also play a role in *Cdkn1c* silencing, since genomic hypomethylation caused by deletion of the *Dnmt1* gene resulted in bi-allelic expression of *Cdkn1c* (Caspary et al., 1998). In this report we provide the first evidence that the methylation of the 5' DMR is critical for paternal silencing, implying that two mechanisms (suppression by the methylated 5' DMR as well as unmethylated KvDMR1) are critical for silencing of the paternal *Cdkn1c* allele.

Although DNA methylation is an important mechanism in the control of imprinting, not all loci are equally affected by methylation and other chromatin-modifying mechanisms are involved, too. For example, in *Dnmt1*^{-/-} mice, H19 and *Cdkn1c* are activated by DNA hypomethylation, whereas *Igf2*, *Igfr* and

Kvlqt1 are silenced and the *Mash2* gene appears unaffected (Li et al., 1993; Caspary et al., 1998). Likewise, uniparental murine embryonal fibroblasts that are treated with the demethylating drug azacytidine show loss of imprinting and de-repression of a few genes (*H19*, *Cdkn1c*, *Peg3*, *Zac1*), whereas other loci remain unaffected (*Grb10*, *Sgca*, *Snrpn*, *U2af1*) (El Kharroubi et al., 2001). Thus, distinct epigenetic mechanisms in addition to DNA methylation have been postulated in the control of imprinting. Furthermore, in some cases the establishment of imprinting may be independent of DNA methylation. For example, in *Dnmt3a/b*-deleted ES cells, CpG methylation is lost at several imprinted loci over time in culture (Chen et al., 2003). Re-introduction of *Dnmt3a/b* transgenes leads to proper remethylation of the paternal allele, but not the maternal allele. Therefore at least at some loci, the imprinted memory is apparently stored as epigenetic modification independent of CpG methylation. The 5' DMR (ICG5) of the *Cdkn1c* is another example, since it is not methylated in germ cells, but obtains differential methylation in somatic cells before day 7.5 of gestation.

Since most DMRs such as the KvDMR1 and other DMRs analyzed in this study, obtain their methylation pattern already in germ cells, they require only maintenance of methylation during embryogenesis. Our results therefore suggest that Lsh does not play a general role in maintenance of methylation, since deletion of the major maintenance methyltransferase *Dnmt1*, in contrast to *Lsh* deletion, leads to loss of methylation and imprinting at many loci (Li et al., 1993; Caspary et al., 1998). Instead, the specific effect of Lsh on the *Cdkn1c* gene is consistent with the hypothesis that Lsh may play a role in de novo methylation because only the 5' DMR of the *Cdkn1c* gene acquires methylation in the embryo and not in germ cells. Lsh may promote recruitment and association of de novo DNA methyltransferases (such as *Dnmt3a*) to specific sites in the genome, or alternatively may facilitate the DNA methyltransferase activity on nucleosomal targets by remodeling chromatin and giving greater accessibility to hidden CpG sites. The patchy loss of methylation at the 5' DMR of the *Cdkn1c* gene (Fig. 5) may reflect the inaccessibility of DNA methyltransferase in the absence of Lsh, to the central portion of the nucleosomal DNA as opposed to the edges that show greater probability of *Dnmt* exposure in vitro (Okuwaki and Verreault, 2004). However, in order to test the hypothesis that Lsh plays an important role in de novo methylation, different experimental systems are required. To examine a general role in de novo methylation, the acquisition of methylation patterns in integrated retroviral sequences in embryonal stem cells or in episomal constructs should be tested (Okano et al., 1999; Hsieh, 1999).

The human and mouse promoter regions of *Cdkn1c* share high homology, however, only the murine CpG islands have been reported to show differential methylation. Thus either the human *CDKN1C* gene may be independent of methylation (and Lsh), or alternatively, Lsh may participate in the imprinting control of the human gene, but independently of CpG methylation. We have previously shown that Lsh also controls post-translational modifications such as histone acetylation or methylation levels (Yan et al., 2003; Huang et al., 2004). However, further analysis of genomic imprinting control at the human *CDKN1C* gene has to be performed in order to determine the role of Lsh.

Loss of imprinting has been implicated in the origin of sporadic cancers and human inherited syndromes that are cancer prone (Reik and Walter, 2001; Paulsen and Ferguson-Smith, 2001; Feinberg et al., 2002). A subset of patients with Beckwith-Wiedemann syndrome that are prone to childhood malignancies show a functional mutation in the *Cdkn1c* gene. We report here the hypomethylation at the *Cdkn1c* promoter correlated with bi-allelic expression. Since *Cdkn1c* is a cell cycle inhibitor its role has been largely implicated as a tumor suppressor gene whose loss of function promotes growth and tumor progression. However, a number of tumors have been reported that do not show silencing, but instead show overexpression of the *Cdkn1c* gene (Hartmann et al., 2000; Lai et al., 2000; Ito et al., 2002). For example, a subset of patients with head and neck cancers, or patients with hepatoblastoma exhibit an upregulation of *Cdkn1c* gene expression with reactivation of the paternal allele, and frequent loss of heterozygosity of the maternal gene. Furthermore, some patients with Wilms tumor show paternal expression of *Cdkn1c* and loss of heterozygosity of the maternal region. Though *Cdkn1c* is a cell cycle inhibitor, it interacts with transcription factors (such as MyoD) and proteins of the c-Jun/stress-activated kinase pathway (Chang et al., 2003; Reynaud et al., 2000). Thus inhibition of the UV-or stress-induced apoptotic pathway *Cdkn1c* may contribute to cancer progression or therapy resistance of some tumors.

Investigating Lsh's unique contribution to the epigenetic regulation at distinct imprinted loci should help our understanding of the multiple mechanisms that control imprinting, and their role in pathogenetic processes such as cancer.

We thank Drs Joost Oppenheim and Nancy Colburn for critical review of the manuscript. We are grateful for the excellent technical assistance of Terry Stull and Rodney Wiles. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the US Government. This project has been funded in whole or part with Federal funds from the National Cancer Institute, National Institutes of Health, under Contract No. N01-C0-12400. Animal care was provided in accordance with the procedures outlined in the 'Guide for the Care and Use of Laboratory Animals' (NIH Publication No. 86-23, 1985).

References

- Bartolomei, M. S. and Tilghman, S. M. (1997). Genomic imprinting in mammals. *Annu. Rev. Genet.* **31**, 493-525.
- Bird, A. (2002). DNA methylation patterns and epigenetic memory. *Genes Dev.* **16**, 6-21.
- Burgess-Beusse, B., Farrell, C., Gaszner, M., Litt, M., Mutskov, V., Recillas-Targa, F., Simpson, M., West, A. and Felsenfeld, G. (2002). The insulation of genes from external enhancers and silencing chromatin. *Proc. Natl. Acad. Sci. USA* **99 Suppl.** 4, 16433-16437.
- Caspary, T., Cleary, M. A., Baker, C. C., Guan, X. J. and Tilghman, S. M. (1998). Multiple mechanisms regulate imprinting of the mouse distal chromosome 7 gene cluster. *Mol. Cell. Biol.* **18**, 3466-3474.
- Chang, T. S., Kim, M. J., Ryoo, K., Park, J., Eom, S. J., Shim, J., Nakayama, K. I., Nakayama, K., Tomita, M., Takahashi, et al. (2003). p57KIP2 modulates stress-activated signaling by inhibiting c-Jun NH2-terminal kinase/stress-activated protein Kinase. *J. Biol. Chem.* **278**, 48092-48098.
- Chen, T., Ueda, Y., Dodge, J. E., Wang, Z. and Li, E. (2003). Establishment and maintenance of genomic methylation patterns in mouse embryonic stem cells by Dnmt3a and Dnmt3b. *Mol. Cell Biol.* **23**, 5594-5605.
- Constancia, M., Pickard, B., Kelsey, G. and Reik, W. (1998). Imprinting mechanisms. *Genome Res.* **8**, 881-900.
- Delaval, K. and Feil, R. (2004). Epigenetic regulation of mammalian genomic imprinting. *Curr. Opin. Genet. Dev.* **14**, 188-195.
- Dennis, K., Fan, T., Geiman, T., Yan, Q. and Muegge, K. (2001). Lsh, a member of the SNF2 family, is required for genome-wide methylation. *Genes Dev.* **15**, 2940-2944.
- Diaz-Meyer, N., Day, C. D., Khatod, K., Maher, E. R., Cooper, W., Reik, W., Junien, C., Graham, G., Algar, E., Der Kaloustian, V. M. and Higgins, M. J. (2003). Silencing of CDKN1C (p57(KIP2)) is associated with hypomethylation at KvDMR1 in Beckwith-Wiedemann syndrome. *J. Med. Genet.* **40**, 797-801.
- El Kharroubi, A., Piras, G. and Stewart, C. L. (2001). DNA demethylation reactivates a subset of imprinted genes in uniparental mouse embryonic fibroblasts. *J. Biol. Chem.* **276**, 8674-8680.
- Elson, D. A. and Bartolomei, M. S. (1997). A 5' differentially methylated sequence and the 3'-flanking region are necessary for H19 transgene imprinting. *Mol. Cell Biol.* **17**, 309-317.
- Fan, T., Yan, Q., Huang, J., Austin, S., Cho, E., Ferris, D. and Muegge, K. (2003). Lsh-deficient murine embryonal fibroblasts show reduced proliferation with signs of abnormal mitosis. *Cancer Res.* **63**, 4677-4683.
- Feinberg, A. P., Cui, H. and Ohlsson, R. (2002). DNA methylation and genomic imprinting: insights from cancer into epigenetic mechanisms. *Semin. Cancer Biol.* **12**, 389-398.
- Fitzpatrick, G. V., Soloway, P. D. and Higgins, M. J. (2002). Regional loss of imprinting and growth deficiency in mice with a targeted deletion of KvDMR1. *Nat. Genet.* **32**, 426-431.
- Geiman, T. M. and Muegge, K. (2000). Lsh, an SNF2/helicase family member, is required for proliferation of mature T lymphocytes. *Proc. Natl. Acad. Sci. USA* **97**, 4772-4777.
- Geiman, T. M., Durum, S. K. and Muegge, K. (1998). Characterization of gene expression, genomic structure, and chromosomal localization of Hells (Lsh). *Genomics* **54**, 477-483.
- Geiman, T. M., Tessarollo, L., Anver, M. R., Kopp, J. B., Ward, J. M. and Muegge, K. (2001). Lsh, a SNF2 family member, is required for normal murine development. *Biochim. Biophys. Acta* **1526**, 211-220.
- Hartmann, W., Waha, A., Koch, A., Goodyer, C. G., Albrecht, S., von Schweinitz, D. and Pietsch, T. (2000). p57(KIP2) is not mutated in hepatoblastoma but shows increased transcriptional activity in a comparative analysis of the three imprinted genes p57(KIP2), IGF2, and H19. *Am. J. Pathol.* **157**, 1393-1403.
- Hatada, I. and Mukai, T. (1995). Genomic imprinting of p57KIP2, a cyclin-dependent kinase inhibitor, in mouse. *Nat. Genet.* **11**, 204-206.
- Hsieh, C. L. (1999). In vivo activity of murine de novo methyltransferases, Dnmt3a and Dnmt3b. *Mol. Cell Biol.* **19**, 8211-8218.
- Huang, J., Fan, T., Yan, Q., Zhu, H., Fox, S., Isaaq, H. J., Best, L., Gangi, L., Munroe, D. and Muegge, K. (2004). Lsh, an epigenetic guardian of repetitive elements. *Nucleic Acids Res.* **32**, 5019-5028.
- Ito, Y., Yoshida, H., Nakano, K., Kobayashi, K., Yokozawa, T., Hirai, K., Matsuzuka, F., Matsuura, N., Kuma, K. and Miyauchi, A. (2002). Expression of p57/Kip2 protein in normal and neoplastic thyroid tissues. *Int. J. Mol. Med.* **9**, 373-376.
- Jarvis, C. D., Geiman, T., Vila-Storm, M. P., Osipovich, O., Akella, U., Candeias, S., Nathan, I., Durum, S. K. and Muegge, K. (1996). A novel putative helicase produced in early murine lymphocytes. *Gene* **169**, 203-207.
- Lai, S., Goepfert, H., Gillenwater, A. M., Luna, M. A. and El Naggar, A. K. (2000). Loss of imprinting and genetic alterations of the cyclin-dependent kinase inhibitor p57KIP2 gene in head and neck squamous cell carcinoma. *Clin. Cancer Res.* **6**, 3172-3176.
- Li, E., Beard, C. and Jaenisch, R. (1993). Role for DNA methylation in genomic imprinting. *Nature* **366**, 362-365.
- Lucifero, D., Mertineit, C., Clarke, H. J., Bestor, T. H. and Trasler, J. M. (2002). Methylation dynamics of imprinted genes in mouse germ cells. *Genomics* **79**, 530-538.
- Mann, J. R., Szabo, P. E., Reed, M. R. and Singer-Sam, J. (2000). Methylated DNA sequences in genomic imprinting. *Crit. Rev. Eukaryot. Gene Expr.* **10**, 241-257.
- Okano, M., Bell, D. W., Haber, D. A. and Li, E. (1999). DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* **99**, 247-257.

- Okuwaki, M. and Verreault, A.** (2004). Maintenance DNA methylation of nucleosome core particles. *J. Biol. Chem.* **279**, 2904-2912.
- Paulsen, M. and Ferguson-Smith, A. C.** (2001). DNA methylation in genomic imprinting, development, and disease. *J. Pathol.* **195**, 97-110.
- Piras, G., El Kharroubi, A., Kozlov, S., Escalante-Alcalde, D., Hernandez, L., Copeland, N. G., Gilbert, D. J., Jenkins, N. A. and Stewart, C. L.** (2000). *Zac1* (Lot1), a potential tumor suppressor gene, and the gene for epsilon-sarcoglycan are maternally imprinted genes: identification by a subtractive screen of novel uniparental fibroblast lines. *Mol. Cell Biol.* **20**, 3308-3315.
- Reik, W. and Walter, J.** (2001). Genomic imprinting: parental influence on the genome. *Nat. Rev. Genet.* **2**, 21-32.
- Reynaud, E. G., Leibovitch, M. P., Tintignac, L. A., Pelpel, K., Guillier, M. and Leibovitch, S. A.** (2000). Stabilization of MyoD by direct binding to p57(Kip2). *J. Biol. Chem.* **275**, 18767-18776.
- Simpson, M., West, A. and Felsenfeld, G.** (2002). The insulation of genes from external enhancers and silencing chromatin. *Proc. Natl. Acad. Sci. USA* **99 Suppl. 4**, 16433-16437.
- Smith, R. J., Arnaud, P., Konfortova, G., Dean, W. L., Beechey, C. V. and Kelsey, G.** (2002). The mouse *Zac1* locus: basis for imprinting and comparison with human *ZAC*. *Gene* **292**, 101-112.
- Spengler, D., Villalba, M., Hoffmann, A., Pantaloni, C., Houssami, S., Bockaert, J. and Journot, L.** (1997). Regulation of apoptosis and cell cycle arrest by *Zac1*, a novel zinc finger protein expressed in the pituitary gland and the brain. *EMBO J.* **16**, 2814-2825.
- Taniguchi, T., Okamoto, K. and Reeve, A. E.** (1997). Human p57(KIP2) defines a new imprinted domain on chromosome 11p but is not a tumour suppressor gene in Wilms tumour. *Oncogene* **14**, 1201-1206.
- Tremblay, K. D., Duran, K. L. and Bartolomei, M. S.** (1997). A 5' 2-kilobase-pair region of the imprinted mouse *H19* gene exhibits exclusive paternal methylation throughout development. *Mol. Cell Biol.* **17**, 4322-4329.
- Verona, R. I., Mann, M. R. and Bartolomei, M. S.** (2003). Genomic imprinting: intricacies of epigenetic regulation in clusters. *Annu. Rev. Cell Dev. Biol.* **19**, 237-259.
- Wutz, A., Smrzka, O. W., Schweifer, N., Schellander, K., Wagner, E. F. and Barlow, D. P.** (1997). Imprinted expression of the *Igf2r* gene depends on an intronic CpG island. *Nature* **389**, 745-749.
- Yan, Q., Huang, J., Fan, T., Zhu, H. and Muegge, K.** (2003a). Lsh, a modulator of CpG methylation, is crucial for normal histone methylation. *EMBO J.* **22**, 5154-5162.
- Yan, Q., Cho, E., Lockett, S. and Muegge, K.** (2003b). Association of Lsh, a regulator of DNA methylation, with pericentromeric heterochromatin is dependent on intact heterochromatin. *Mol. Cell Biol.* **23**, 8416-8428.
- Yatsuki, H., Joh, K., Higashimoto, K., Soejima, H., Arai, Y., Wang, Y., Hatada, I., Obata, Y., Morisaki, H., Zhang, Z., Nakagawachi, T., Satoh, Y. and Mukai, T.** (2002). Domain regulation of imprinting cluster in *Kip2/Lit1* subdomain on mouse chromosome 7F4/F5: large-scale DNA methylation analysis reveals that DMR-Lit1 is a putative imprinting control region. *Genome Res.* **12**, 1860-1870.