

Left-right patterning: conserved and divergent mechanisms

Tetsuya Nakamura* and Hiroshi Hamada

Summary

The left-right (LR) asymmetry of visceral organs is fundamental to their function and position within the body. Over the past decade or so, the molecular mechanisms underlying the establishment of such LR asymmetry have been revealed in many vertebrate and invertebrate model organisms. These studies have identified a gene network that contributes to this process and is highly conserved from sea urchin to mouse. By contrast, some specific steps of the process, such as the symmetry-breaking event and situs-specific organogenesis, appear to have diverged during evolution. Here, we summarize the common and divergent mechanisms by which LR asymmetry is established in vertebrates.


Key words: Nodal, Cilia, Left-right asymmetry

Introduction

The establishment of left-right (LR) asymmetry during early embryogenesis is crucial for the correct positioning and morphogenesis of internal organs. This process involves a number of intricately regulated developmental mechanisms, some of which appear to be conserved among vertebrates. For some organisms, LR symmetry breaking takes place around the left-right organizer (LR organizer; the node in mice, Hensen's node in the chick, the gastrocoel roof plate in *Xenopus*, and Kupffer's vesicle in zebrafish). In these animals, except for the chick and pig, asymmetric fluid flow produced by rotation of multiple cilia on the LR organizer results in asymmetric gene expression around the LR organizer. Earlier asymmetries in the localization of some molecules are observed in *Xenopus* and chick embryos and contribute to the establishment of LR asymmetry, but the roles of these molecules in establishing LR asymmetry in other organisms still need to be established (Spéder et al., 2007; Vandenberg and Levin, 2010). Following the symmetry-breaking event, asymmetric information is transmitted to surrounding tissues and induces the asymmetric expression of *Nodal* and *Lefty*. *Nodal* and *Lefty*,

Developmental Genetics Group, Graduate School of Frontier Biosciences, Osaka University, Osaka 565-0871, Japan.

*Author for correspondence (t-nakamura@fbs.osaka-u.ac.jp)



(See poster insert)

members of the transforming growth factor β (TGF β) superfamily, act as diffusible activator and inhibitor molecules, respectively, and constitute a self-enhancement and lateral-inhibition (SELI) system, which is highly conserved among vertebrates and some non-vertebrates, and provides robustness to LR determination. Finally, asymmetric Nodal signals induce the left-sided expression of other genes, such as pituitary homeobox 2 (Pitx2), which in turn contribute to the LR asymmetric morphogenesis of various visceral organs. The cellular mechanisms underlying situs-specific organogenesis, by contrast, seem to vary among vertebrates.

Early asymmetries

In *Xenopus* embryos, several genes exhibit asymmetric patterns of gene expression at early stages of development (Kramer et al., 2002; Fukumoto et al., 2005). The earliest LR asymmetry in *Xenopus* embryos described to date is the localization of H⁺- and K⁺-dependent ATPase α mRNA and protein in the two-cell embryo (Aw et al., 2008); a few hours after fertilization, the amount of these mRNAs and protein is higher in one cell than in the other. The mechanisms responsible for the unequal distribution of H⁺/K⁺-ATPase α mRNA and protein are unknown but probably involve unidirectional transport by a motor protein along the cytoskeleton. In addition, directionality of the embryonic cytoskeleton itself, that is the position of the '+' ends and '-' ends, might play a role in moving the motor protein in one direction and further analysis of this role is expected. At the 32-cell stage, the neurotransmitter serotonin is localized asymmetrically in *Xenopus* embryos, and this LR asymmetric localization requires gap junctions and H⁺/K⁺-ATPase α (Fukumoto et al., 2005). Furthermore, pharmacological inhibition of serotonin receptors R3 and R4 and injection of mutant serotonin receptor mRNAs in *Xenopus* embryos result in heterotaxia (Fukumoto et al., 2005). These observations thus suggest that H⁺/K⁺-ATPase α may determine the distribution of serotonin through gap junctions in the *Xenopus* embryo, and that this might in turn influence LR patterning. Other molecules, such as PKC- γ and phosphorylated syndecan 2, have also been shown to be distributed asymmetrically in *Xenopus* embryos (Kramer et al., 2002). Furthermore, H⁺-V-ATPase is also indispensable for left-right asymmetry in *Xenopus* and chick (Adams et al., 2006). In the chick, asymmetric localization of H⁺/K⁺-ATPase α mRNA is apparent at Hensen's node and is required for the subsequent establishment of Sonic hedgehog (Shh) asymmetric gene expression (Aw et al., 2008). However, in general, such early asymmetries in mRNA or protein localization have not been studied in detail in other vertebrate embryos.

Symmetry-breaking events and fluid flow

In the mouse, gastrulation in a symmetric embryo results in the formation of the primitive streak at the future posterior side. The formation of the organizer (the node) at the anterior tip of the primitive streak leads to the breaking of symmetry as a consequence of the concerted rotation of cilia present in the organizer (Essner et al., 2005; Nonaka et al., 2005; Schweickert et al., 2007). These cilia rotate in a clockwise direction (when viewed from a ventral position) and subsequently generate a leftward flow of extra-embryonic fluid. The precise mechanism by which the fluid flow results in symmetry breaking is unknown, but it has been suggested that it transports small molecules or vesicular particles to the left side of the embryo (Hirokawa et al., 2006) or exerts mechanical pressure on the left wall of the organizer (McGrath et al., 2003; Tabin and Vogan, 2003).

Two types of cilia, motile and immotile, have been identified in the mouse node, with the latter type, which are present at the edge

of the node, having been proposed to sense mechanical force (McGrath et al., 2003). The basal body, an organelle located at the base of each cilium, is located on the posterior side of mouse node cells (Nonaka et al., 2005; Okada et al., 2005). This localization results in a posterior tilt of the cilium, because the surface of node cells is curved. As a consequence of the posterior tilt of the rotation axis of the node cilia, a stroke towards the left side is more efficient in generating fluid flow than is a stroke toward the right side (Cartwright et al., 2004; Nonaka et al., 2005; Okada et al., 2005).

The gastrocoel roof plate (GRP) in *Xenopus* embryos and Kupffer's vesicle (KV) in zebrafish and Medaka embryos are the ciliated organs of asymmetry generation in these organisms. Thus, the GRP and KV similarly possess motile cilia and their movement generates unidirectional flow. In *Xenopus*, fluid flow is necessary only in the left portion of the GRP (Vick et al., 2009), suggesting that the cells on the left side of this structure sense the signal, whether it be mechanical or chemical. In zebrafish, KV acts as the LR organizer but its architecture is different from that of the mouse node. Thus, motile cilia are present at the ventral and dorsal surface inside this vesicle, but they are preferentially found in the anterior region of KV. Although cilia in the anterior region are posteriorly tilted, the basal body is not significantly shifted towards the posterior side. Nonetheless, unidirectional flow (counterclockwise as observed from the dorsal side) is formed in the anterior-dorsal region of KV (Kreiling et al., 2007; Okabe et al., 2008).

The posterior localization of the basal body within node cells can be regarded as a type of planar cell polarity (PCP) and is regulated by the noncanonical Wnt signaling pathway in the mouse and *Xenopus*, and in zebrafish, the distribution of cilia inside KV is also regulated (Maisonneuve et al., 2009; Zhang and Levin, 2009; Antic et al., 2010; Borovina et al., 2010; Hashimoto et al., 2010; Song et al., 2010; Wang et al., 2011). Whereas Dishevelled proteins, mediators of Wnt signaling, are localized to the posterior end of node cells (Hashimoto et al., 2010), the PCP protein Prickle2 is found at the anterior end (Antic et al., 2010). However, the initial anterior-posterior cue responsible for polarizing of the distribution of these proteins is unknown.

Recent studies have shown that calcium might also play a role in the symmetry-breaking event. Imaging of intracellular Ca²⁺ with a Ca²⁺ indicator has revealed an asymmetry in Ca²⁺ signaling on the two sides of the vertebrate organizer (McGrath et al., 2003; Tanaka et al., 2005; Garic-Stankovic et al., 2008; Francescato et al., 2010). Furthermore, polycystin 2 (polycystic kidney disease 2; Pkd2) and polycystin 1 like 1 (polycystic kidney disease 1 like 1; Pkd11), which form Ca²⁺ channels, are required in the mouse for asymmetric *Nodal* expression in the lateral plate mesoderm (LPM) (Pennekamp et al., 2002; Field et al., 2011), and Ca²⁺/calmodulin-dependent kinase II (CamKII) is required in KV for LR patterning in zebrafish (Francescato et al., 2010). However, *curly up* (*pkd2*) zebrafish mutant shows bilateral expression of *southpaw* (Schottenfeld et al., 2007), which is different from the phenotype of the *Pkd2* mouse mutant, which lacks *Nodal* expression in the LPM. These observations implicate Ca²⁺ influx in the establishment of LR asymmetry, but the precise role of such influx remains unknown. Given that the gene for the *Nodal* inhibitor Cerberus like 2 (*Cerl2*; also known as *Coco*) is the earliest known target of the leftward fluid flow in *Xenopus* embryos (Schweickert et al., 2010), it is possible that Ca²⁺ may regulate the expression of this gene directly or indirectly.

In the chick embryo, symmetry breaking does not appear to involve cilia-generated flow. Instead, asymmetric cell migration

around the LR organizer is an early event that has been observed in birds (Cui et al., 2009; Gros et al., 2009); cells migrate from the right to the left around Hensen's node (anticlockwise), resulting in the generation of asymmetric gene expression around this structure. The directionality of this movement is determined by H^+/K^+ ATPase. Intriguingly, the pig embryo appears to lack cilia at the LR organizer (Gros et al., 2009), suggesting that it also uses a cilia-independent mechanism for breaking symmetry. Interestingly, it is known that some migrating cells, including neutrophils, possess intrinsic polarity, which could be generated by a subcellular chiral structure such as the centrosome (Xu et al., 2007; Wan et al., 2011), and it might be possible that such a chiral structure determines the direction of cell migration for LR patterning. In the mouse embryo, however, cells at the periphery of the node move symmetrically around the node (Yamanaka et al., 2007). Furthermore, *Nodal* expression at the node is indispensable for *Nodal* expression in the LPM of the mouse (Brennan et al., 2002). Therefore, symmetry breaking by directional migration might be a mechanism that is specific to only some organisms.

Asymmetric gene expression around the LR organizer

The cilia-generated flow detected in many vertebrates, or the directional cell migration observed in the chick embryo, induces asymmetric gene expression around the organizer. The gene for *Cerl2* (Dand5 – Mouse Genome Informatics), which encodes a Nodal antagonist, shows an asymmetric (R>L) expression pattern around the node of the mouse embryo. *Cerl2* thus inhibits the activity of Nodal preferentially on the right side and is required for normal LR patterning (Marques et al., 2004). In *Xenopus*, *Coco* is the ortholog of *Cerl2* and is the gene thought to respond earliest to leftward flow; it thus manifests asymmetric expression around the organizer and plays a role similar to that of *Cerl2* (Schweickert et al., 2010), although histone deacetylase activity at earlier stages might be involved in epigenetic silencing of *Nodal* (Carneiro et al., 2011). In zebrafish and Medaka, Charon (Dand5 – Zebrafish Information Network), an ortholog of mouse *Cerl2*, is asymmetrically expressed (R>L) in KV and is indispensable for left and right asymmetry (Hashimoto et al., 2004; Schneider et al., 2010; Hojo et al., 2007; Lopes et al., 2010).

Asymmetric signal generated in the node is transmitted to the surrounding tissues. Current data collectively suggest that, in the mouse, Nodal protein generated by perinodal cells at the node is transported via the internal route of the embryonic tissue, that is between the endoderm and mesoderm, to the left LPM where it induces its own expression (Oki et al., 2007; Kawasumi et al., 2011). Alternatively, calcium signals may be transmitted from the node to the LPM, given that a calcium signal is detected at the left margin of the node (McGrath et al., 2003). Furthermore, as a result of the R>L asymmetric expression of *Cerl2*, Nodal activity is preferentially inhibited on the right side. Indeed, the distribution of *Nodal* mRNA around the node shows a small asymmetry, whereas the activity of Nodal (as reflected by the level of Smad2/3 phosphorylation) is highly L>R asymmetric (Kawasumi et al., 2011).

In the chick embryo, the establishment of asymmetry around the node seems to involve different signaling pathways. Thus, at the right side of the node, ActivinB, which belongs to the TGF β superfamily, stimulates the expression of bone morphogenetic protein 4 (*Bmp4*), which in turn represses *Shh* expression (Levin et al., 1995; Monsoro-Burq and Le Douarin, 2001). At the left side of the node, where ActivinB is downregulated, *Shh* is induced,

which represses *Bmp4* expression and induces expression of *cNR-1* (a Nodal ortholog). This gene network establishes asymmetric gene expression around the node and the resulting information (probably asymmetric Nodal activity, as for the mouse) is transmitted to the left LPM where it activates *cNR-1* expression.

Asymmetric gene expression in the left lateral plate

After symmetry-breaking in the LR organizer, asymmetric gene expression arises in the lateral plate on the left side (Meno et al., 1999; Saijoh et al., 2005; Shiratori et al., 2006). In the mouse, *Nodal*, *Lefty2* and the transcription factor *Pitx2* are expressed in the LPM exclusively on the left side. Whereas Nodal functions as a left-side determinant, Lefty is a feedback inhibitor of Nodal. Nodal induces its own expression via a Nodal-responsive enhancer of *Nodal*, and the operation of this positive feedback loop underlies the expansion of the Nodal expression domain. Nodal also simultaneously induces the expression of *Lefty2*. This combination of positive and negative feedback loops constitutes a SELI system that is able to amplify a small difference between the left and right of the node into a robust difference in the LPM, because *Nodal* expression in the right LPM is inhibited by *Lefty* diffusing from the left LPM (Nakamura et al., 2006). Mathematical modeling indicates that the SELI system can establish correct LR patterning only if the diffusion velocity of *Lefty* is higher than that of Nodal. That *Lefty* travels faster than Nodal was suggested by an early study (Sakuma et al., 2002), but recent work has directly revealed such a difference in the diffusion velocities of *Lefty* and Nodal (Marjoram and Wright, 2011). The diffusion of *Lefty* and Nodal also appears to be facilitated by sulfated proteoglycans of the extracellular matrix (Oki et al., 2007; Marjoram and Wright, 2011).

The SELI system composed of Nodal and *Lefty* seems to be conserved among vertebrate species (Nakamura et al., 2006; Marjoram and Wright, 2011); however, other genes are also expressed asymmetrically in the LPM. In the chick embryo, bone morphogenetic protein (BMP) represses *Nodal* expression in the LPM, whereas BMP signaling positively upregulates CFC (a member of EGF-CFC: Cripto/FRL-1/Cryptic) expression, which is indispensable for Nodal positive feedback loop (Piedra and Ros, 2002). However, *caronte*, an inhibitor of BMP signaling that is expressed in the paraxial mesoderm on the right, induces *Nodal* expression in the left LPM by inhibiting BMP activity (Yokouchi et al., 1999). Similarly in the mouse embryo, chordin and noggin, inhibitors of BMP signaling, are expressed at a higher level in the left LPM than in the right LPM, suggesting that BMP signaling might repress *Nodal* expression on the right side (Mine et al., 2008). In addition, the transcriptional repressor *snail* is preferentially induced by fibroblast growth factor 8 (*Fgf8*) in the right LPM and represses *Nodal* expression in the chick embryo, whereas mutant mice that conditionally lack *Snail* exhibit bilateral *Nodal* expression (Boettger et al., 1999; Patel et al., 1999; Murray and Gridley, 2006). In the zebrafish embryo, a different mechanism appears to inhibit *Nodal* expression on the right side. Thus, inhibition of *Nodal* on the right side involves a BMP signal at the posterior tip of the LPM and *Lefty2* at the anterior side (*Lefty2* expression is present only in the anterior region of LPM in the zebrafish) (Bisgrove et al., 2000; Lenhart et al., 2011).

Asymmetric Nodal signaling in the left lateral plate induces expression of *Pitx2*, which encodes a transcription factor with a bicoid-type homeobox, in the same region. Whereas Nodal induces *Pitx2* expression transiently and asymmetrically by activating FoxH1, which is a transcription factor of the TGF β pathway and

binds to ASE (asymmetric enhancer), asymmetric *Pitx2* expression is maintained as a result of the activity of Nkx2-dependent enhancer after the transient activation by FoxH1. FoxH1 is responsible for inducing transcription of *Pitx2* whereas Nkx2 maintains *Pitx2* expression once it has been induced (Shiratori et al., 2001).

Asymmetric morphogenesis

Left-sided expression of *Pitx2* induces asymmetric morphogenesis of most of visceral organs, including the heart, lungs, stomach, intestine and kidneys. However, the cellular mechanisms responsible for asymmetric organogenesis in vertebrates remain largely unknown. Several studies have provided insight into the mechanism of gut looping. Looping of the chick gut is determined by LR asymmetries in the cellular architecture of the dorsal mesentery, the structure that connects the developing gut to the body wall (Davis et al., 2008; Kurpios et al., 2008; Plageman et al., 2011). Furthermore, the overlying epithelium shows LR asymmetries in cellular morphology that are under the control of *Pitx2*. Subsequent morphogenesis of gut looping can be explained by the relative growth between the gut tube and the anchoring dorsal mesentery (Savin et al., 2011). In the case of zebrafish, the differential migration of LPM cells on the left and right sides gives rise to gut looping (Horne-Badovinac et al., 2003). This asymmetric migration of LPM cells depends on Nodal signaling but is independent of the overlying endoderm, and it involves cell rearrangements associated with remodeling of the extracellular matrix (Yin et al., 2010). Whether such mechanisms exist in the mouse and in *Xenopus* is under investigation.

During morphogenesis of the heart, leftward looping of the aortic arch results from asymmetric regression of the sixth branchial arch, which is induced by asymmetric blood flow triggered by the spiral rotation of the outflow tract in the mouse embryo (Yashiro et al., 2007). In zebrafish, asymmetric migration of cardiac progenitor cells is responsible for this cardiac rotation. This asymmetric migration is regulated by BMP signaling, which is preferentially activated in the LPM on the left side and is controlled by early asymmetric Nodal signaling (Baker et al., 2008; de Campos-Baptista et al., 2008; Rohr et al., 2008; Smith et al., 2008). However, it remains unknown whether this migration depends on *Pitx2* or not. In fact, some of the asymmetric morphogenesis events are independent of *Pitx2*. For example, heart looping and embryonic turning remain normal in the mutant mouse lacking asymmetric *Pitx2* expression (Shiratori et al., 2006). Interestingly, recent work suggests that the direction of embryonic turning depends on blastomere compositions at the 8-cell stage (Gardner, 2010). This pivotal finding also suggests that a part of asymmetry in the mouse may in fact be determined prior to node formation. It remains to be determined whether the cellular mechanisms proposed for the morphogenesis of individual organs in individual model animals are shared among vertebrates.

Perspectives

In summary, multiple mechanisms contribute to the generation of LR asymmetry in vertebrates and some of these appear to be conserved. The gene network comprising *Nodal*, *Lefty* and *Pitx2*, for example, is highly conserved among animals. Asymmetric expression of *Nodal* also appears to be conserved among deuterostomes (Chea et al., 2005). In the deuterostome sea urchin embryo, for example, these genes are expressed asymmetrically, albeit exclusively on the right side (Duboc and Lepage, 2008), which is possibly owing to the inversion of the dorsal-ventral axis

during evolution of bilaterian animals. Nodal is also found in non-deuterostomes, for example in snails that belong to the Lophotrochozoa (Grande and Patel, 2009; Kuroda et al., 2009). *Lefty* is similarly conserved although its precise expression domains might vary.

By contrast, the cellular mechanisms underlying asymmetric morphogenesis have probably diverged among species. Furthermore, there is considerable diversity in initial symmetry breaking even among vertebrates; whereas many vertebrates rely on unidirectional flow generated by cilia, the chick and pig do not adopt this strategy, and both the frog and chick exhibit asymmetries prior to organizer formation. Furthermore, chiral blastomere arrangement controls LR asymmetry in snails (Kuroda et al., 2009), whereas H^+/K^+ -ATPase determines the asymmetry in *Ciona* (Shimeld and Levin, 2006). The different strategies deployed for symmetry breaking (reviewed by Spéder et al., 2007) may originate from the divergence of morphologies or developmental processes (or both) among organisms. A key challenge is to determine to what extent each step of the LR pathway is conserved among species. It will be interesting to know the precise mechanism of symmetry breaking in diverse organisms. Another challenge is to clarify the cellular mechanisms underlying asymmetric morphogenesis, including those occurring during heart looping, branching and turning, which are largely unknown at present. Further development of various approaches, genetic, cellular, biophysical and mathematical, will undoubtedly help to clarify these issues.

Acknowledgements

We thank A. Fukumoto, Y. Ikawa and H. Nisimura for technical assistance.

Funding

This study was supported by a Grant-in-Aid for Scientific Research on Innovative Areas and a Grant-in-Aid for Young Scientists (B) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan; by Core Research for Evolutional Science and Technology (CREST) of the Japan Science and Technology Corporation (JST); and by Global Center of Excellence (GCOE) of Osaka University.

Competing interests statement

The authors declare no competing financial interests.

Development at a Glance

A high-resolution version of the poster is available for downloading in the online version of this article at <http://dev.biologists.org/content/139/18/3257.full>

References

- Adams, D. S., Robinson, K. R., Fukumoto, T., Yuan, S., Albertson, R. C., Yelick, P., Kuo, L., McSweeney, M. and Levin, M. (2006). Early, H^+ -V-ATPase-dependent proton flux is necessary for consistent left-right patterning of non-mammalian vertebrates. *Development* **133**, 1657-1671.
- Antic, D., Stubbs, J. L., Suyama, K., Kintner, C., Scott, M. P. and Axelrod, J. D. (2010). Planar cell polarity enables posterior localization of nodal cilia and left-right axis determination during mouse and *Xenopus* embryogenesis. *PLoS ONE* **5**, e8999.
- Aw, S., Adams, D. S., Qiu, D. and Levin, M. (2008). H,K -ATPase protein localization and Kir4.1 function reveal concordance of three axes during early determination of left-right asymmetry. *Mech. Dev.* **125**, 353-372.
- Baker, K., Holtzman, N. G. and Burdine, R. D. (2008). Direct and indirect roles for Nodal signaling in two axis conversions during asymmetric morphogenesis of the zebrafish heart. *Proc. Natl. Acad. Sci. USA* **105**, 13924-13929.
- Bisgrove, B. W., Essner, J. J. and Yost, H. J. (2000). Multiple pathways in the midline regulate concordant brain, heart and gut left-right asymmetry. *Development* **127**, 3567-3579.
- Boettger, T., Wittler, L. and Kessel, M. (1999). FGF8 functions in the specification of the right body side of the chick. *Curr. Biol.* **9**, 277-280.
- Borovina, A., Superina, S., Voskas, D. and Ciruna, B. (2010). Vangl2 directs the posterior tilting and asymmetric localization of motile primary cilia. *Nat. Cell Biol.* **12**, 407-412.
- Brennan, J., Norris, D. P. and Robertson, E. J. (2002). Nodal activity in the node governs left-right asymmetry. *Genes Dev.* **16**, 2339-2344.

- Carneiro, K., Donnet, C., Rejtar, T., Karger, B. L., Barisone, G. A., Diaz, E., Kortagere, S., Lemire, J. M. and Levin, M. (2011). Histone deacetylase activity is necessary for left-right patterning during vertebrate development. *BMC Dev. Biol.* **11**, 29.
- Cartwright, J. H., Piro, O. and Tuval, I. (2004). Fluid-dynamical basis of the embryonic development of left-right asymmetry in vertebrates. *Proc. Natl. Acad. Sci. USA* **101**, 7234-7239.
- Chea, H. K., Wright, C. V. and Swalla, B. J. (2005). Nodal signaling and the evolution of deuterostome gastrulation. *Dev. Dyn.* **234**, 269-278.
- Cui, C., Little, C. D. and Rongish, B. J. (2009). Rotation of organizer tissue contributes to left-right asymmetry. *Anat. Rec. (Hoboken)* **292**, 557-561.
- Davis, N. M., Kurpios, N. A., Sun, X., Gros, J., Martin, J. F. and Tabin, C. J. (2008). The chirality of gut rotation derives from left-right asymmetric changes in the architecture of the dorsal mesentery. *Dev. Cell* **15**, 134-145.
- de Campos-Baptista, M. I., Holtzman, N. G., Yelon, D. and Schier, A. F. (2008). Nodal signaling promotes the speed and directional movement of cardiomyocytes in zebrafish. *Dev. Dyn.* **237**, 3624-3633.
- Duboc, V. and Lepage, T. (2008). A conserved role for the nodal signaling pathway in the establishment of dorso-ventral and left-right axes in deuterostomes. *J. Exp. Zool.* **310B**, 41-53.
- Essner, J. J., Amack, J. D., Nyholm, M. K., Harris, E. B. and Yost, H. J. (2005). Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left-right development of the brain, heart and gut. *Development* **132**, 1247-1260.
- Field, S., Riley, K. L., Grimes, D. T., Hilton, H., Simon, M., Powles-Glover, N., Siggers, P., Bogani, D., Greenfield, A. and Norris, D. P. (2011). Pkd11 establishes left-right asymmetry and physically interacts with Pkd2. *Development* **138**, 1131-1142.
- Francescato, L., Rothschild, S. C., Myers, A. L. and Tombes, R. M. (2010). The activation of membrane targeted CaMK-II in the zebrafish Kupffer's vesicle is required for left-right asymmetry. *Development* **137**, 2753-2762.
- Fukumoto, T., Kema, I. P. and Levin, M. (2005). Serotonin signaling is a very early step in patterning of the left-right axis in chick and frog embryos. *Curr. Biol.* **15**, 794-803.
- Gardner, R. L. (2010). Normal bias in the direction of fetal rotation depends on blastomere composition during early cleavage in the mouse. *PLoS ONE* **5**, e9610.
- Garic-Stankovic, A., Hernandez, M., Flentke, G. R., Zile, M. H. and Smith, S. M. (2008). A ryanodine receptor-dependent Ca²⁺ asymmetry at Hensen's node mediates avian lateral identity. *Development* **135**, 3271-3280.
- Grande, C. and Patel, N. H. (2009). Nodal signalling is involved in left-right asymmetry in snails. *Nature* **457**, 1007-1011.
- Gros, J., Feistel, K., Viebahn, C., Blum, M. and Tabin, C. J. (2009). Cell movements at Hensen's node establish left/right asymmetric gene expression in the chick. *Science* **324**, 941-944.
- Hashimoto, H., Rebagliati, M., Ahmad, N., Muraoka, O., Kurokawa, T., Hibi, M. and Suzuki, T. (2004). The Cerberus/Dan-family protein Charon is a negative regulator of Nodal signaling during left-right patterning in zebrafish. *Development* **131**, 1741-1753.
- Hashimoto, M., Shinohara, K., Wang, J., Ikeuchi, S., Yoshida, S., Meno, C., Nonaka, S., Takada, S., Hatta, K., Wynshaw-Boris, A. et al. (2010). Planar polarization of node cells determines the rotational axis of node cilia. *Nat. Cell Biol.* **12**, 170-176.
- Hirokawa, N., Tanaka, Y., Okada, Y. and Takeda, S. (2006). Nodal flow and the generation of left-right asymmetry. *Cell* **125**, 33-45.
- Hojo, M., Takashima, S., Kobayashi, D., Sumeragi, A., Shimada, A., Tsukahara, T., Yokoi, H., Narita, T., Jindo, T., Kage, T. et al. (2007). Right-elevated expression of charon is regulated by fluid flow in medaka Kupffer's vesicle. *Dev. Growth Differ.* **49**, 395-405.
- Horne-Badovinac, S., Rebagliati, M. and Stainier, D. Y. (2003). A cellular framework for gut-looping morphogenesis in zebrafish. *Science* **302**, 662-665.
- Kawasumi, A., Nakamura, T., Iwai, N., Yashiro, K., Saijoh, Y., Belo, J. A., Shiratori, H. and Hamada, H. (2011). Left-right asymmetry in the level of active Nodal protein produced in the node is translated into left-right asymmetry in the lateral plate of mouse embryos. *Dev. Biol.* **353**, 321-330.
- Kramer, K. L., Barnette, J. E. and Yost, H. J. (2002). PKC γ regulates syndecan-2 inside-out signaling during xenopus left-right development. *Cell* **111**, 981-990.
- Kreiling, J. A., Prabhat Williams, G. and Creton, R. (2007). Analysis of Kupffer's vesicle in zebrafish embryos using a cave automated virtual environment. *Dev. Dyn.* **236**, 1963-1969.
- Kuroda, R., Endo, B., Abe, M. and Shimizu, M. (2009). Chiral blastomere arrangement dictates zygotic left-right asymmetry pathway in snails. *Nature* **462**, 790-794.
- Kurpios, N. A., Ibañez, M., Davis, N. M., Lui, W., Katz, T., Martin, J. F., Izpisua Belmonte, J. C. and Tabin, C. J. (2008). The direction of gut looping is established by changes in the extracellular matrix and in cell:cell adhesion. *Proc. Natl. Acad. Sci. USA* **105**, 8499-8506.
- Lenhart, K. F., Lin, S. Y., Titus, T. A., Postlethwait, J. H. and Burdine, R. D. (2011). Two additional midline barriers function with midline lefty1 expression to maintain asymmetric Nodal signaling during left-right axis specification in zebrafish. *Development* **138**, 4405-4410.
- Levin, M., Johnson, R. L., Stern, C. D., Kuehn, M. and Tabin, C. (1995). A molecular pathway determining left-right asymmetry in chick embryogenesis. *Cell* **82**, 803-814.
- Lopes, S. S., Lourenço, R., Pacheco, L., Moreno, N., Kreiling, J. and Saúde, L. (2010). Notch signalling regulates left-right asymmetry through ciliary length control. *Development* **137**, 3625-3632.
- Maisonneuve, C., Guilleret, I., Vick, P., Weber, T., Andre, P., Beyer, T., Blum, M. and Constam, D. B. (2009). Bicaudal C, a novel regulator of Dvl signaling abutting RNA-processing bodies, controls cilia orientation and leftward flow. *Development* **136**, 3019-3030.
- Marjoram, L. and Wright, C. (2011). Rapid differential transport of Nodal and Lefty on sulfated proteoglycan-rich extracellular matrix regulates left-right asymmetry in *Xenopus*. *Development* **138**, 475-485.
- Marques, S., Borges, A. C., Silva, A. C., Freitas, S., Cordenonsi, M. and Belo, J. A. (2004). The activity of the Nodal antagonist Cerl-2 in the mouse node is required for correct L/R body axis. *Genes Dev.* **18**, 2342-2347.
- McGrath, J., Somlo, S., Makova, S., Tian, X. and Brueckner, M. (2003). Two populations of node monocilia initiate left-right asymmetry in the mouse. *Cell* **114**, 61-73.
- Meno, C., Gritsman, K., Ohishi, S., Ohfujii, Y., Heckscher, E., Mochida, K., Shimono, A., Kondoh, H., Talbot, W. S., Robertson, E. J. et al. (1999). Mouse Lefty2 and zebrafish antivin are feedback inhibitors of nodal signalling during vertebrate gastrulation. *Mol. Cell* **4**, 287-298.
- Mine, N., Anderson, R. M. and Klingensmith, J. (2008). BMP antagonism is required in both the node and lateral plate mesoderm for mammalian left-right axis establishment. *Development* **135**, 2425-2434.
- Monsoro-Burq, A. and Le Douarin, N. M. (2001). BMP4 plays a key role in left-right patterning in chick embryos by maintaining Sonic Hedgehog asymmetry. *Mol. Cell* **7**, 789-799.
- Murray, S. A. and Gridley, T. (2006). Snail family genes are required for left-right asymmetry determination, but not neural crest formation, in mice. *Proc. Natl. Acad. Sci. USA* **103**, 10300-10304.
- Nakamura, T., Mine, N., Nakaguchi, E., Mochizuki, A., Yamamoto, M., Yashiro, K., Meno, C. and Hamada, H. (2006). Generation of robust left-right asymmetry in the mouse embryo requires a self-enhancement and lateral-inhibition system. *Dev. Cell* **11**, 495-504.
- Nonaka, S., Yoshida, S., Watanabe, D., Ikeuchi, S., Goto, T., Marshall, W. F. and Hamada, H. (2005). De novo formation of left-right asymmetry by posterior tilt of nodal cilia. *PLoS Biol.* **3**, e268.
- Okabe, N., Xu, B. and Burdine, R. D. (2008). Fluid dynamics in zebrafish Kupffer's vesicle. *Dev. Dyn.* **237**, 3602-3612.
- Okada, Y., Takeda, S., Tanaka, Y., Izpisua Belmonte, J. C. and Hirokawa, N. (2005). Mechanism of nodal flow: a conserved symmetry breaking event in left-right axis determination. *Cell* **121**, 633-644.
- Oki, S., Hashimoto, R., Okui, Y., Shen, M. M., Mekada, E., Otani, H., Saijoh, Y. and Hamada, H. (2007). Sulfated glycosaminoglycans are necessary for Nodal signal transmission from the node to the left lateral plate in the mouse embryo. *Development* **134**, 3893-3904.
- Patel, K., Isaac, A. and Cooke, J. (1999). Nodal signalling and the roles of the transcription factors SnR and Pitx2 in vertebrate left-right asymmetry. *Curr. Biol.* **9**, 609.
- Pennekamp, P., Karcher, C., Fischer, A., Schweickert, A., Skryabin, B., Horst, J., Blum, M. and Dworniczak, B. (2002). The ion channel polycystin-2 is required for left-right axis determination in mice. *Curr. Biol.* **12**, 938-943.
- Piedra, M. E. and Ros, M. A. (2002). BMP signaling positively regulates Nodal expression during left right specification in the chick embryo. *Development* **129**, 3431-3440.
- Plageman, T. F., Jr, Zacharias, A. L., Gage, P. J. and Lang, R. A. (2011). Shroom3 and a Pitx2-N-cadherin pathway function cooperatively to generate asymmetric cell shape changes during gut morphogenesis. *Dev. Biol.* **357**, 227-234.
- Rohr, S., Otten, C. and Abdelilah-Seyfried, S. (2008). Asymmetric involution of the myocardial field drives heart tube formation in zebrafish. *Circ. Res.* **102**, e12-e19.
- Saijoh, Y., Oki, S., Tanaka, C., Nakamura, T., Adachi, H., Yan, Y. T., Shen, M. M. and Hamada, H. (2005). Two nodal-responsive enhancers control left-right asymmetric expression of Nodal. *Dev. Dyn.* **232**, 1031-1036.
- Sakuma, R., Ohnishi, Y.-i., Meno, C., Fujii, H., Juan, H., Takeuchi, J., Ogura, T., Li, E., Miyazono, K. and Hamada, H. (2002). Inhibition of Nodal signalling by Lefty mediated through interaction with common receptors and efficient diffusion. *Genes Cells* **7**, 401-412.
- Savin, T., Kurpios, N. A., Shyer, A. E., Florescu, P., Liang, H., Mahadevan, L. and Tabin, C. J. (2011). On the growth and form of the gut. *Nature* **476**, 57-62.
- Schneider, I., Schneider, P. N., Derry, S. W., Lin, S., Barton, L. J., Westfall, T. and Slusarski, D. C. (2010). Zebrafish Nkd1 promotes Dvl degradation and is required for left-right patterning. *Dev. Biol.* **348**, 22-33.
- Schottenfeld, J., Sullivan-Brown, J. and Burdine, R. D. (2007). Zebrafish curly up encodes a Pkd2 ortholog that restricts left-side-specific expression of southpaw. *Development* **134**, 1605-1615.

- Schweickert, A., Weber, T., Beyer, T., Vick, P., Bogusch, S., Feistel, K. and Blum, M. (2007). Cilia-driven leftward flow determines laterality in *Xenopus*. *Curr. Biol.* **17**, 60-66.
- Schweickert, A., Vick, P., Getwan, M., Weber, T., Schneider, I., Eberhardt, M., Beyer, T., Pachur, A. and Blum, M. (2010). The nodal inhibitor *Coco* is a critical target of leftward flow in *Xenopus*. *Curr. Biol.* **20**, 738-743.
- Shimeld, S. M. and Levin, M. (2006). Evidence for the regulation of left-right asymmetry in *Ciona intestinalis* by ion flux. *Dev. Dyn.* **235**, 1543-1553.
- Shiratori, H., Sakuma, R., Watanabe, M., Hashiguchi, H., Mochida, K., Sakai, Y., Nishino, J., Saijoh, Y., Whitman, M. and Hamada, H. (2001). Two-step regulation of left-right asymmetric expression of *Pitx2*: initiation by nodal signaling and maintenance by *Nkx2*. *Mol. Cell* **7**, 137-149.
- Shiratori, H., Yashiro, K., Shen, M. M. and Hamada, H. (2006). Conserved regulation and role of *Pitx2* in situs-specific morphogenesis of visceral organs. *Development* **133**, 3015-3025.
- Smith, K. A., Chocron, S., von der Hardt, S., de Pater, E., Soufan, A., Bussmann, J., Schulte-Merker, S., Hammerschmidt, M. and Bakkers, J. (2008). Rotation and asymmetric development of the zebrafish heart requires directed migration of cardiac progenitor cells. *Dev. Cell* **14**, 287-297.
- Song, H., Hu, J., Chen, W., Elliott, G., Andre, P., Gao, B. and Yang, Y. (2010). Planar cell polarity breaks bilateral symmetry by controlling ciliary positioning. *Nature* **466**, 378-382.
- Spéder, P., Petzoldt, A., Suzanne, M. and Noselli, S. (2007). Strategies to establish left/right asymmetry in vertebrates and invertebrates. *Curr. Opin. Genet. Dev.* **17**, 351-358.
- Tabin, C. J. and Vogan, K. J. (2003). A two-cilia model for vertebrate left-right axis specification. *Genes Dev.* **17**, 1-6.
- Tanaka, Y., Okada, Y. and Hirokawa, N. (2005). FGF-induced vesicular release of Sonic hedgehog and retinoic acid in leftward nodal flow is critical for left-right determination. *Nature* **435**, 172-177.
- Vandenberg, L. N. and Levin, M. (2010). Far from solved: a perspective on what we know about early mechanisms of left-right asymmetry. *Dev. Dyn.* **239**, 3131-3146.
- Vick, P., Schweickert, A., Weber, T., Eberhardt, M., Mencl, S., Shcherbakov, D., Beyer, T. and Blum, M. (2009). Flow on the right side of the gastrocoel roof plate is dispensable for symmetry breakage in the frog *Xenopus laevis*. *Dev. Biol.* **331**, 281-291.
- Wan, L. Q., Ronaldson, K., Park, M., Taylor, G., Zhang, Y., Gimble, J. M. and Vunjak-Novakovic, G. (2011). Micropatterned mammalian cells exhibit phenotype-specific left-right asymmetry. *Proc. Natl. Acad. Sci. USA* **108**, 12295-12300.
- Wang, G., Cadwallader, A. B., Jang, D. S., Tsang, M., Yost, H. J. and Amack, J. D. (2011). The Rho kinase *Rock2b* establishes anteroposterior asymmetry of the ciliated Kupffer's vesicle in zebrafish. *Development* **138**, 45-54.
- Xu, J., Van Keymeulen, A., Wakida, N. M., Carlton, P., Berns, M. W. and Bourne, H. R. (2007). Polarity reveals intrinsic cell chirality. *Proc. Natl. Acad. Sci. USA* **104**, 9296-9300.
- Yamanaka, Y., Tamplin, O. J., Beckers, A., Gossler, A. and Rossant, J. (2007). Live imaging and genetic analysis of mouse notochord formation reveals regional morphogenetic mechanisms. *Dev. Cell* **13**, 884-896.
- Yashiro, K., Shiratori, H. and Hamada, H. (2007). Haemodynamics determined by a genetic programme govern asymmetric development of the aortic arch. *Nature* **450**, 285-288.
- Yin, C., Kikuchi, K., Hochgreb, T., Poss, K. D. and Stainier, D. Y. (2010). *Hand2* regulates extracellular matrix remodeling essential for gut-looping morphogenesis in zebrafish. *Dev. Cell* **18**, 973-984.
- Yokouchi, Y., Vogan, K. J., Pearse, R. V., 2nd and Tabin, C. J. (1999). Antagonistic signaling by *Caronte*, a novel Cerberus-related gene, establishes left-right asymmetric gene expression. *Cell* **98**, 573-583.
- Zhang, Y. and Levin, M. (2009). Left-right asymmetry in the chick embryo requires core planar cell polarity protein *Vangl2*. *Genesis* **47**, 719-728.